

Gravity in three dimensions as a noncommutative gauge theory

George Manolakos

National Technical University of Athens

March 30, 2018

Joint work with: A. Chatzistavrakidis, L. Jonke, D. Jurman, P. Manousselis, G.
Zoupanos



[arXiv:1802.075]

Table of contents

1. *Gravity as gauge theory in 1+2 (and 1+3) dimensions*
2. *How to build gauge theories on noncommutative (nc) spaces*
3. *Gauge theory on 3-d nc spaces based on $SU(2)$ and $SU(1,1)$*
4. *Gravity as gauge theory on 3-d nc spaces (Euclidean - Lorentzian)*

Gravity in three dimensions as a gauge theory

The algebra

Witten '88

- ▶ 3-d Gravity: gauge theory of $\mathfrak{iso}(1,2)$ (Poincaré - isometry of M^3)
- ▶ Presence of Λ : dS or AdS algebras, i.e. $\mathfrak{so}(1,3), \mathfrak{so}(2,2)$
- ▶ Corresponding generators: $P_a, J_{ab}, a = 1, 2, 3$ (translations, LT)
- ▶ Satisfy the following CRs:

$$[J_{ab}, J_{cd}] = 4\eta_{[a[c}J_{d]b]}, \quad [P_a, J_{bc}] = 2\eta_{a[b}P_{c]}, \quad [P_a, P_b] = \Lambda J_{ab}$$

- ▶ CRs valid in *arbitrary* dim; particularly in 3-d:

$$[J_a, J_b] = \epsilon_{abc}J^c, \quad [P_a, J_b] = \epsilon_{abc}P^c, \quad [P_a, P_b] = \Lambda\epsilon_{abc}J^c$$

- ▶ After the redefinition: $J^a = \frac{1}{2}\epsilon^{abc}J_{bc}$

The gauging procedure

- ▶ Intro of a gauge field for each generator: e_μ^a, ω_μ^a (transl, LT)
- ▶ The Lie-valued 1-form gauge connection is:

$$A_\mu = e_\mu^a(x)P_a + \omega_\mu^a(x)J_a$$

- ▶ Transforms in the adjoint rep, according to the rule:

$$\delta A_\mu = \partial_\mu \epsilon + [A_\mu, \epsilon]$$

- ▶ The gauge transformation parameter is expanded as:

$$\epsilon = \xi^a(x)P_a + \lambda^a(x)J_a$$

- ▶ *Combining* the above \rightarrow transformations of the fields:

$$\begin{aligned}\delta e_\mu^a &= \partial_\mu \xi^a - \epsilon^{abc}(\xi_b \omega_{\mu c} + \lambda_b e_{\mu c}) \\ \delta \omega_\mu^a &= \partial_\mu \lambda^a - \epsilon^{abc}(\lambda_b \omega_{\mu c} + \Lambda \xi_b e_{\mu c})\end{aligned}$$

Curvatures and action

- ▶ Curvatures of the fields are given by:

$$R_{\mu\nu}(A) = 2\partial_{[\mu}A_{\nu]} + [A_{\mu}, A_{\nu}]$$

- ▶ Tensor $R_{\mu\nu}$ is also Lie-valued:

$$R_{\mu\nu}(A) = T_{\mu\nu}{}^a P_a + R_{\mu\nu}{}^a J_a$$

- ▶ *Combining* the above \rightarrow curvatures of the fields:

$$\begin{aligned} T_{\mu\nu}{}^a &= 2\partial_{[\mu}e_{\nu]}{}^a + 2\epsilon^{abc}\omega_{[\mu b}e_{\nu]c} \\ R_{\mu\nu}{}^a &= 2\partial_{[\mu}\omega_{\nu]}{}^a + \epsilon^{abc}(\omega_{\mu b}\omega_{\nu c} + \Lambda e_{\mu b}e_{\nu c}) \end{aligned}$$

- ▶ The Chern-Simons action functional of the Poincaré, dS or AdS algebra is found to be *identical* to the 3-d E-H action:

$$\mathcal{S}_{CS} = \frac{1}{16\pi G} \int \epsilon^{\mu\nu\rho} (e_{\mu}{}^a (\partial_{\nu}\omega_{\rho a} - \partial_{\rho}\omega_{\nu a}) + \epsilon_{abc} e_{\mu}{}^a \omega_{\nu}{}^b \omega_{\rho}{}^c + \frac{\Lambda}{3} \epsilon_{abc} e_{\mu}{}^c e_{\nu}{}^b e_{\rho}{}^c) \equiv \mathcal{S}_{EH}$$

Remarks on 4-d gravity

Utiyama '56, Kibble '61
MacDowell-Mansouri '77
Kibble-Stelle '85

- ▶ Vielbein formulation of GR: Gauging Poincaré algebra $\mathfrak{iso}(1,3)$
- ▶ Comprises ten generators: $P_a, J_{ab}, a = 1, \dots, 4$ (transl, LT)
- ▶ Satisfy the aforementioned CRs (for $\Lambda = 0$)
- ▶ Gauging in the same way leading to field transformations
- ▶ Curvatures are obtained accordingly
- ▶ Dynamics follow from the E-H action:

$$\mathcal{S}_{EH4} = \frac{1}{2} \int d^4x \epsilon^{\mu\nu\rho\sigma} \epsilon_{abcd} e_\mu^a e_\nu^b R_{\rho\sigma}{}^{cd}$$

- ▶ *There is no such action in gauge theory*

In 4-d gravity, only kinematics can be obtained by gauge theory

Gauge theories on noncommutative spaces

The nc framework

Szabo '01

- ▶ Late 40's: Nc structure of spacetime at small scales for an effective ultraviolet cutoff → control of divergences in qfts *Snyder '47*
- ▶ Ignored → success of renormalization programme
- ▶ Inspiration from qm: Operators instead of variables
- ▶ Nc spacetime defined by replacing coords x^i by Herm generators X^i of a nc algebra of functions, \mathcal{A} , obeying: $[X^i, X^j] = i\theta^{ij}$ *Connes '94, Madore '99*
- ▶ 80's: nc geometry revived after the generalization of diff structure *Connes '85, Woronowicz '87*
- ▶ Along with the definition of a generalized integration → Yang-Mills gauge theories on nc spaces *Connes-Rieffel '87*
- ▶ Evidence for spacetime nc from string theory due to min length

Nc gauge theories

- ▶ Operators $X_\mu \in \mathcal{A}$ satisfy the CR: $[X_\mu, X_\nu] = i\theta_{\mu\nu}$, $\theta_{\mu\nu}$ arbitrary
- ▶ Lie-type nc: $[X_\mu, X_\nu] = iC_{\mu\nu}{}^\rho X_\rho$
- ▶ Natural intro of nc gauge theories through *covariant nc coordinates*: $\mathcal{X}_\mu = X_\mu + A_\mu$ *Madore-Schraml-Schupp-Wess '00*
- ▶ Obeys a covariant gauge transformation rule: $\delta\mathcal{X}_\mu = i[\epsilon, \mathcal{X}_\mu]$
- ▶ A_μ transforms in analogy with the gauge connection:
 $\delta A_\mu = -i[X_\mu, \epsilon] + i[\epsilon, A_\mu]$, (ϵ - the gauge parameter)
- ▶ Definition of a (Lie-type) nc *covariant field strength tensor*:
 $F_{\mu\nu} = [\mathcal{X}_\mu, \mathcal{X}_\nu] - iC_{\mu\nu\rho}\mathcal{X}_\rho$
- ▶ Gauge theory could be abelian or nonabelian:
 - ▶ Abelian if ϵ is a function in \mathcal{A}
 - ▶ Nonabelian if ϵ is matrix valued ($\text{Mat}(\mathcal{A})$)

Non-Abelian case

▷ *In nonabelian case, where are the gauge fields valued?*

▶ Let us consider the CR of two elements of an algebra:

$$[\epsilon, A] = [\epsilon^A T^A, A^B T^B] = \frac{1}{2} \{\epsilon^A, A^B\} [T^A, T^B] + \frac{1}{2} [\epsilon^A, A^B] \{T^A, T^B\}$$

▶ *Not possible to restrict to a matrix algebra:*

last term neither *vanishes* in nc nor is an *algebra element*

▶ There are two options to overpass the difficulty:

▶ Consider the universal enveloping algebra

▶ Extend the generators and/or fix the rep so that the anticommutators close

▷ *We employ the second option*

3-d fuzzy spaces based on $SU(2)$ and $SU(1,1)$

The Euclidean case

- ▶ Euclidean case: 3-d fuzzy space based on $SU(2)$
- ▶ Fuzzy sphere, S_F^2 : Matrix approximation of ordinary sphere, S^2
Madore '92
- ▶ S^2 defined by coordinates of \mathbb{R}^3 modulo $\sum_{a=1}^3 x_a x^a = r^2$
- ▶ S_F^2 defined by three rescaled angular momentum operators, $X_i = \lambda J_i$, J_i the Lie algebra generators in a UIR of $SU(2)$. The X_i s satisfy:

$$[X_i, X_j] = i\lambda \epsilon_{ijk} X_k, \quad \sum_{i=1}^3 X_i X_i = \lambda^2 j(j+1) := r^2, \lambda \in \mathbb{R}, 2j \in \mathbb{N}$$

- ▶ Allowing X_i to live in *reducible* rep: obtain the nc \mathbb{R}_λ^3 , viewed as direct sum of S_F^2 with all possible radii (determined by $2j$)
Vitale-Wallet '13, Vitale '14
Hammou-Lagraa-Sheikh Jabbari '02
- ▶ \mathbb{R}_λ^3 : discrete foliation of \mathbb{R}^3 by multiple S_F^2 of different radii

The Lorentzian case

- ▶ In analogy: Lorentzian case: 3-d fuzzy space based on $SU(1, 1)$

D.J.-Steinacker '14

- ▶ Fuzzy hyperboloids, dS_F^2 , defined by three rescaled operators, $X_i = \lambda J_i$, (in appropriate irreps) satisfying:

$$[X_i, X_j] = i\lambda C_{ij}^k X_k, \quad \sum_{i,j} \eta_{ij} X_i X_j = \lambda^2 j(j-1),$$

- ▶ where C_{ij}^k are the structure constants of $\mathfrak{su}(1, 1)$
- ▶ Difference to previous case: Non-compact group, i.e. no finite-dim UIRs but infinite-dim
- ▶ Again, letting X_i live in (infinite-dim) *reducible* reps: Block diagonal form - each block being a dS_F^2
- ▶ 3-d Minkowski spacetime foliated with leaves being dS_F^2 of different radii

Gravity as gauge theory on 3-d fuzzy spaces

The Lorentzian case

Aschieri-Castellani '09

- ▶ Consideration of the foliated M^3 with $\lambda > 0$
- ▶ Relevant isometry group: $SO(3, 1)$
- ▶ Consider the corresponding spin group:
 $SO(3, 1) \cong Spin(3, 1) = SL(2, \mathbb{C})$
- ▶ Anticommutators *do not close* \rightarrow Fix at spinor rep generated by:
$$\sum_{AB} = \frac{1}{2}\gamma_{AB} = \frac{1}{4}[\gamma_A, \gamma_B], A = 1, \dots, 4$$
- ▶ Satisfying the CRs and aCRs:
$$[\gamma_{AB}, \gamma_{CD}] = 8\eta_{[A[C\gamma_{D]B]}, \quad \{\gamma_{AB}, \gamma_{CD}\} = 4\eta_{C[B\eta_{A]D}\mathbb{1} + 2i\epsilon_{ABCD}\gamma_5$$
- ▶ Inclusion of γ_5 and identity in the algebra \rightarrow extension of $SL(2, \mathbb{C})$ to $GL(2, \mathbb{C})$ with set of generators: $\{\gamma_{AB}, \gamma_5, i\mathbb{1}\}$

SO(3) notation

- ▶ In $SO(3)$ notation: $\gamma_{a4} \equiv \gamma_a$ and $\tilde{\gamma}^a \equiv \epsilon^{abc}\gamma_{bc}$, with $a = 1, 2, 3$
- ▶ The CRs and aCRs are now written:

$$\begin{aligned}[\tilde{\gamma}^a, \tilde{\gamma}^b] &= -4\epsilon^{abc}\tilde{\gamma}_c, & [\gamma_a, \tilde{\gamma}_b] &= -4\epsilon_{abc}\gamma^c, & [\gamma_a, \gamma_b] &= \epsilon_{abc}\tilde{\gamma}^c, & [\gamma^5, \gamma^{AB}] &= 0 \\ \{\tilde{\gamma}^a, \tilde{\gamma}^b\} &= -8\eta^{ab}\mathbb{1}, & \{\gamma_a, \tilde{\gamma}^b\} &= 4i\delta_a^b\gamma_5, & \{\gamma_a, \gamma_b\} &= 2\eta_{ab}\mathbb{1}, \\ \{\gamma^5, \gamma^a\} &= i\tilde{\gamma}_a, & \{\tilde{\gamma}^5, \gamma^a\} &= -4i\gamma_a\end{aligned}$$

- ▶ Proceed with the gauging of $GL(2, \mathbb{C})$
- ▶ Determine the covariant coordinate: $\mathcal{X}_\mu = X_\mu + \mathcal{A}_\mu$
 $\mathcal{A}_\mu = \mathcal{A}_\mu^i(X_a) \otimes T^i$ the $\mathfrak{gl}(2, \mathbb{C})$ -valued gauge connection

- ▶ Gauge connection expands on the generators as:

$$\mathcal{A}_\mu = e_\mu^a(X) \otimes \gamma_a + \omega_\mu^a(X) \otimes \tilde{\gamma}_a + A_\mu(X) \otimes i\mathbb{1} + \tilde{A}_\mu(X) \otimes \gamma_5$$

- ▶ Gauge parameter, ϵ , expands similarly:

$$\epsilon = \xi^a(X) \otimes \gamma_a + \lambda^a(X) \otimes \tilde{\gamma}_a + \epsilon_0(X) \otimes i\mathbb{1} + \tilde{\epsilon}_0(X) \otimes \gamma_5$$

Kinematics

- Covariant transf rule: $\delta\mathcal{X}_\mu = [\epsilon, \mathcal{X}_\mu] \rightarrow$ transf of the gauge fields:

$$\delta e_\mu^a = -i[X_\mu + A_\mu, \xi^a] - 2\{\xi_b, \omega_{\mu c}\}\epsilon^{abc} - 2\{\lambda_b, e_{\mu c}\}\epsilon^{abc} + i[\epsilon_0, e_\mu^a] - 2i[\lambda^a, \tilde{A}_\mu] - 2i[\tilde{\epsilon}_0, \omega_\mu^a]$$

$$\delta\omega_\mu^a = -i[X_\mu + A_\mu, \lambda^a] + \frac{1}{2}\{\xi_b, e_{\mu c}\}\epsilon^{abc} - 2\{\lambda_b, \omega_{\mu c}\}\epsilon^{abc} + i[\epsilon_0, \omega_\mu^a] + \frac{i}{2}[\xi^a, \tilde{A}_\mu] + \frac{i}{2}[\tilde{\epsilon}_0, e_\mu^a]$$

$$\delta A_\mu = -i[X_\mu + A_\mu, \epsilon_0] - i[\xi_a, e_\mu^a] + 4i[\lambda_a, \omega_\mu^a] - i[\tilde{\epsilon}_0, \tilde{A}_\mu]$$

$$\delta\tilde{A}_\mu = -i[X_\mu + A_\mu, \tilde{\epsilon}_0] + 2i[\xi_a, \omega_\mu^a] + 2i[\lambda_a, e_\mu^a] + i[\epsilon_0, \tilde{A}_\mu]$$

- Abelian limit: $e_\mu^a = \omega_\mu^a = \tilde{A}_\mu = 0$: $\delta A_\mu = -i[X_\mu, \epsilon_0] + i[\epsilon_0, A_\mu] \rightarrow$ trans rule of a nc Maxwell gauge field
- Commutative limit: Y-M and gravity fields disentangle and inner derivation becomes $[X_\mu, f] \rightarrow -i\partial_\mu f$:

$$\delta e_\mu^a = -\partial_\mu \xi^a - 4\xi_b \omega_{\mu c} \epsilon^{abc} - 4\lambda_b e_{\mu c} \epsilon^{abc}$$

$$\delta\omega_\mu^a = -\partial_\mu \lambda^a + \xi_b e_{\mu c} \epsilon^{abc} - 4\lambda_b \omega_{\mu c} \epsilon^{abc}$$

- After the redefinitions: $\gamma_a \rightarrow \frac{2i}{\sqrt{\Lambda}} P_a$, $\tilde{\gamma}_a \rightarrow -4J_a$, $4\lambda^a \rightarrow \lambda^a$,
 $\xi^a \frac{2i}{\sqrt{\Lambda}} \rightarrow -\xi^a$, $e_\mu^a \rightarrow \frac{\sqrt{\Lambda}}{2i} e_\mu^a$, $\omega_\mu^a \rightarrow -\frac{1}{4}\omega_\mu^a \rightarrow$ 3-d gravity

Curvatures

- ▶ Definition of curvature:

$$\mathcal{R}_{\mu\nu} = [\mathcal{X}_\mu, \mathcal{X}_\nu] - i\lambda C_{\mu\nu}{}^\rho \mathcal{X}_\rho$$

- ▶ Curvature tensor can be expanded in the $GL(2, \mathbb{C})$ generators:

$$\mathcal{R}_{\mu\nu} = T_{\mu\nu}^a \otimes \gamma_a + R_{\mu\nu}^a \otimes \tilde{\gamma}_a + F_{\mu\nu} \otimes i\mathbb{1} + \tilde{F}_{\mu\nu} \otimes \gamma_5$$

- ▶ The expressions of the various tensors are:

$$T_{\mu\nu}^a = i[X_\mu + A_\mu, e_\nu^a] - i[X_\nu + A_\nu, e_\mu^a] - 2\{e_{\mu b}, \omega_{\nu c}\}\epsilon^{abc} - 2\{\omega_{\mu b}, e_{\nu c}\}\epsilon^{abc} - 2i[\omega_\mu^a, \bar{A}_\nu] + 2i[\omega_\nu^a, \bar{A}_\mu] - i\lambda C_{\mu\nu}{}^\rho e_\rho^a$$

$$R_{\mu\nu}^a = i[X_\mu + A_\mu, \omega_\nu^a] - i[X_\nu + A_\nu, \omega_\mu^a] - 2\{\omega_{\mu b}, \omega_{\nu c}\}\epsilon^{abc} + \frac{1}{2}\{e_{\mu b}, e_{\nu c}\}\epsilon^{abc} + \frac{i}{2}[e_\mu^a, \bar{A}_\nu] - \frac{i}{2}[e_\nu^a, \bar{A}_\mu] - i\lambda C_{\mu\nu}{}^\rho \omega_\rho^a$$

$$F_{\mu\nu} = i[X_\mu + A_\mu, X_\nu + A_\nu] - i[e_\mu^a, e_{\nu a}] + 4i[\omega_\mu^a, \omega_{\nu a}] - i[\bar{A}_\mu, \bar{A}_\nu] - i\lambda C_{\mu\nu}{}^\rho (X_\rho + A_\rho)$$

$$\tilde{F}_{\mu\nu} = i[X_\mu + A_\mu, \bar{A}_\nu] - i[X_\nu + A_\nu, \bar{A}_\mu] + 2i[e_\mu^a, \omega_{\nu a}] + 2i[\omega_\mu^a, e_{\nu a}] - i\lambda C_{\mu\nu}{}^\rho \bar{A}_\rho$$

- ▶ Commutative limit: *Coincidence* with the expressions of 3-d gravity after applying the redefinitions

The Euclidean case

- ▶ Isometry group: $SO(4) \cong Spin(4) = SU(2) \times SU(2)$
- ▶ Anticommutators *do not close* \rightarrow Extension to $U(2) \times U(2)$
- ▶ Each $U(2)$: four 4x4 matrices as generators:

$$J_a^L = \begin{pmatrix} \sigma_a & 0 \\ 0 & 0 \end{pmatrix}, \quad J_a^R = \begin{pmatrix} 0 & 0 \\ 0 & \sigma_a \end{pmatrix}, \quad J_0^L = \begin{pmatrix} \mathbb{1} & 0 \\ 0 & 0 \end{pmatrix}, \quad J_0^R = \begin{pmatrix} 0 & 0 \\ 0 & \mathbb{1} \end{pmatrix}$$

- ▶ Identification of the correct nc dreibein and spin connection fields:

$$P_a = \frac{1}{2}(J_a^L - J_a^R), \quad M_a = \frac{1}{2}(J_a^L + J_a^R), \quad \mathbb{1} = J_0^L + J_0^R, \quad \gamma_5 = J_0^L - J_0^R$$

- ▶ Calculations give the CRs and aCRs
- ▶ Gauging proceeds in the same way as before - only difference the signature of metric

- ▶ The action we propose is Chern-Simons type:

$$\mathcal{S} = \frac{1}{g^2} \text{Trtr} \left(\frac{i}{3} C^{\mu\nu\rho} \mathcal{X}_\mu \mathcal{X}_\nu \mathcal{X}_\rho - \frac{\lambda}{2} \mathcal{X}_\mu \mathcal{X}^\mu \right)$$

- ▶ Tr: Trace over matrices X ; tr: Trace over the algebra
- ▶ The action can be written as:

$$\mathcal{S} = \frac{1}{6g^2} \text{Trtr}(iC^{\mu\nu\rho} \mathcal{X}_\mu \mathcal{R}_{\nu\rho}) + \mathcal{S}_\lambda$$

- ▶ where $\mathcal{S}_\lambda = -\frac{\lambda}{6g^2} \text{Trtr}(\mathcal{X}_\mu \mathcal{X}^\mu)$
- ▶ Using the explicit form of the algebra trace:

$$\text{Tr}C^{\mu\nu\rho} (e_{\mu a} T_{\nu\rho}^a - 4\omega_{\mu a} R_{\nu\rho}^a - (X_\mu + A_\mu) F_{\nu\rho} + \tilde{A}_\mu \tilde{F}_{\nu\rho})$$

- ▶ Commutative limit: First two term *identical* to 3-d gravity (after redefinition)

Summary

- ▶ 3-d gravity described as gauge theory
- ▶ Translation to nc regime (gauge theories through cov. coord.)
- ▶ 3-d nc spacetimes built from $SU(2)$ and $SU(1,1)$
- ▶ Gauge their symmetry groups
- ▶ Transformations of fields - Curvatures - Action
- ▶ 3-d gravity recovered at comm limit

Future plans

- ▶ Further analysis of the Lorentzian case space structure
- ▶ Variation - equations of motion
- ▶ Move to the realistic 4-d case

THE END