

# Model Building with F-Theory GUTs

J. Callaghan, S. King, G. Leontaris, G. Ross

James Callaghan

University of Southampton

12/09/2011

# Outline

- Introduction and motivation
- Local F-theory models
- 'Dictionary' between different GUT theories
- Model which stabilises the proton and forbids R-parity violating operators

# Introduction and Motivation

# Introduction and Motivation

- A major motivation for string theory is that it provides a consistent formulation of quantum gravity; the effects of which are expected to become important at the Planck scale,  $M_{Planck} \sim 10^{19} GeV$

# Introduction and Motivation

- A major motivation for string theory is that it provides a consistent formulation of quantum gravity; the effects of which are expected to become important at the Planck scale,  $M_{Planck} \sim 10^{19} GeV$
- It seems very hard to predict anything about particle physics from such theories, because of the large number of consistent solutions to the string theory equations of motion

# Introduction and Motivation

- A major motivation for string theory is that it provides a consistent formulation of quantum gravity; the effects of which are expected to become important at the Planck scale,  $M_{Planck} \sim 10^{19} GeV$
- It seems very hard to predict anything about particle physics from such theories, because of the large number of consistent solutions to the string theory equations of motion
- In order to facilitate the search for realistic models, we can adopt the selection criteria of **unification** and **decoupling**

# Introduction and Motivation

- A major motivation for string theory is that it provides a consistent formulation of quantum gravity; the effects of which are expected to become important at the Planck scale,  $M_{Planck} \sim 10^{19} GeV$
- It seems very hard to predict anything about particle physics from such theories, because of the large number of consistent solutions to the string theory equations of motion
- In order to facilitate the search for realistic models, we can adopt the selection criteria of **unification** and **decoupling**
- **Decoupling** refers to the existence of a limit where in principle  $\frac{M_{GUT}}{M_{Planck}} \rightarrow 0$  (of course this limit is not taken in practice)

# Introduction and Motivation

- A major motivation for string theory is that it provides a consistent formulation of quantum gravity; the effects of which are expected to become important at the Planck scale,  $M_{Planck} \sim 10^{19} GeV$
- It seems very hard to predict anything about particle physics from such theories, because of the large number of consistent solutions to the string theory equations of motion
- In order to facilitate the search for realistic models, we can adopt the selection criteria of **unification** and **decoupling**
- **Decoupling** refers to the existence of a limit where in principle  $\frac{M_{GUT}}{M_{Planck}} \rightarrow 0$  (of course this limit is not taken in practice)
- A class of models which have these properties are **F-theory GUTs**



# What is an F-Theory GUT

# What is an F-Theory GUT

- F-theory is a 12d strongly coupled formulation of type IIB superstring theory


# What is an F-Theory GUT

- F-theory is a 12d strongly coupled formulation of type IIB superstring theory
- The 12 dimensional theory can be broken up into the 4 large spacetime dimensions, 6 internal dimensions, and 2 additional dimensions which encode the variation of the real and imaginary parts of the axio-dilaton

# What is an F-Theory GUT

- F-theory is a 12d strongly coupled formulation of type IIB superstring theory
- The 12 dimensional theory can be broken up into the 4 large spacetime dimensions, 6 internal dimensions, and 2 additional dimensions which encode the variation of the real and imaginary parts of the axio-dilaton
- The formal language of **global** F-theory models is that of an elliptically fibred Calabi-Yau fourfold,  $X$ , with a threefold base,  $B_3$

# Moebius Strip- Example of a Fibre Bundle





- The form of the elliptic fibration is

$$y^2 + \alpha_1 xy + \alpha_3 y = x^3 + \alpha_2 x^2 + \alpha_4 x + \alpha_6$$

- $(x,y)$  are coordinates on the torus fibre, and the  $\alpha_j$  are functions of the base coordinates

- The form of the elliptic fibration is

$$y^2 + \alpha_1 xy + \alpha_3 y = x^3 + \alpha_2 x^2 + \alpha_4 x + \alpha_6$$

- $(x,y)$  are coordinates on the torus fibre, and the  $\alpha_i$  are functions of the base coordinates
- When the discriminant vanishes  $\rightarrow$  singularity  $\rightarrow$  D7 branes
- The type of singularity is determined by the order of vanishing



- The form of the elliptic fibration is

$$y^2 + \alpha_1 xy + \alpha_3 y = x^3 + \alpha_2 x^2 + \alpha_4 x + \alpha_6$$

- $(x,y)$  are coordinates on the torus fibre, and the  $\alpha_i$  are functions of the base coordinates
- When the discriminant vanishes  $\rightarrow$  singularity  $\rightarrow$  D7 branes
- The type of singularity is determined by the order of vanishing
- 

$$SU(5) \rightarrow \alpha_1 = b_5, \alpha_2 = b_4 z, \alpha_3 = b_3 z^2, \alpha_4 = b_2 z^3, \alpha_6 = b_0 z^5$$

$$E_6 \rightarrow \alpha_1 = b_5 z, \alpha_2 = b_4 z^2, \alpha_3 = b_3 z^2, \alpha_4 = b_2 z^3, \alpha_6 = b_0 z^5$$

- The form of the elliptic fibration is

$$y^2 + \alpha_1 xy + \alpha_3 y = x^3 + \alpha_2 x^2 + \alpha_4 x + \alpha_6$$

- $(x,y)$  are coordinates on the torus fibre, and the  $\alpha_i$  are functions of the base coordinates
- When the discriminant vanishes  $\rightarrow$  singularity  $\rightarrow$  D7 branes
- The type of singularity is determined by the order of vanishing
- 

$$SU(5) \rightarrow \alpha_1 = b_5, \alpha_2 = b_4 z, \alpha_3 = b_3 z^2, \alpha_4 = b_2 z^3, \alpha_6 = b_0 z^5$$

$$E_6 \rightarrow \alpha_1 = b_5 z, \alpha_2 = b_4 z^2, \alpha_3 = b_3 z^2, \alpha_4 = b_2 z^3, \alpha_6 = b_0 z^5$$

- To get a local model, we assign scaling dimensions to the coordinates, and discard irrelevant terms, eg.

$$E_6 : y^2 = x^3 + b_3 y z^2 + b_2 x z^3 + b_0 z^5$$

# Local F-Theory Models

# Local F-Theory Models

- The idea of **local** F-theory models however, is to focus on the submanifold  $S$ , where the **GUT symmetry**  $G_S$  is localised

# Local F-Theory Models


- The idea of **local** F-theory models however, is to focus on the submanifold  $S$ , where the **GUT symmetry**  $G_S$  is localised
- Intersections of the gauge brane with other 7-branes wrapping surfaces  $S_i$  and supporting gauge groups  $G_i$  gives rise to **matter curves**  $\Sigma_i = S \cap S_i$ . Along the matter curves, the local symmetry group is enhanced to  $G_{\Sigma_i} \supset G_S \times G_i$

# Local F-Theory Models

- The idea of **local** F-theory models however, is to focus on the submanifold  $S$ , where the **GUT symmetry**  $G_S$  is localised
- Intersections of the gauge brane with other 7-branes wrapping surfaces  $S_i$  and supporting gauge groups  $G_i$  gives rise to **matter curves**  $\Sigma_i = S \cap S_i$ . Along the matter curves, the local symmetry group is enhanced to  $G_{\Sigma_i} \supset G_S \times G_i$
- When matter curves intersect at a point in the internal geometry, there is a further enhancement, and there is a Yukawa coupling induced

# Local F-Theory Models

- The idea of **local** F-theory models however, is to focus on the submanifold  $S$ , where the **GUT symmetry**  $G_S$  is localised
- Intersections of the gauge brane with other 7-branes wrapping surfaces  $S_i$  and supporting gauge groups  $G_i$  gives rise to **matter curves**  $\Sigma_i = S \cap S_i$ . Along the matter curves, the local symmetry group is enhanced to  $G_{\Sigma_i} \supset G_S \times G_i$
- When matter curves intersect at a point in the internal geometry, there is a further enhancement, and there is a Yukawa coupling induced
- **Different view**: larger gauge group, broken by a position dependent VEV for an adjoint Higgs field

$B_3 \sim \text{gravity}$ 



# The Role of Exceptional Gauge Symmetries

# The Role of Exceptional Gauge Symmetries


- In F-theory, the fact that  $g_s$  is of order one provides us with the presence of **exceptional symmetry groups**-  $E_6$ ,  $E_7$  and  $E_8$

# The Role of Exceptional Gauge Symmetries

- In F-theory, the fact that  $g_s$  is of order one provides us with the presence of **exceptional symmetry groups**-  $E_6$ ,  $E_7$  and  $E_8$
- In F-theory GUTs, all the interactions of particles are assumed to arise from a point of  $E_8$  in the internal geometry

# The Role of Exceptional Gauge Symmetries

- In F-theory, the fact that  $g_s$  is of order one provides us with the presence of **exceptional symmetry groups**-  $E_6$ ,  $E_7$  and  $E_8$
- In F-theory GUTs, all the interactions of particles are assumed to arise from a point of  $E_8$  in the internal geometry



# An SU(5) Example

# An $SU(5)$ Example

- We take the gauge symmetry on the surface  $S$  to be  $G_S = SU(5)$

# An $SU(5)$ Example

- We take the gauge symmetry on the surface  $S$  to be  $G_S = SU(5)$
- 

$$E_8 \supset SU(5) \times SU(5)_\perp$$

$$248 \rightarrow (24, 1) + (1, 24) + (10, 5) + (\bar{5}, 10) + (\bar{10}, \bar{5}) + (5, \bar{10})$$

# An $SU(5)$ Example

- We take the gauge symmetry on the surface  $S$  to be  $G_S = SU(5)$
- 

$$E_8 \supset SU(5) \times SU(5)_\perp$$

$$248 \rightarrow (24, 1) + (1, 24) + (10, 5) + (\bar{5}, 10) + (\bar{10}, \bar{5}) + (5, \bar{10})$$

- The equations for the curves in terms of the weights  $t_i$  ( $i = 1, \dots, 5$ ,  $\sum t_i = 0$ ), of the 5 representation of  $SU(5)$  are

$$\Sigma_{10} : t_i = 0$$

$$\Sigma_5 : -t_i - t_j = 0, i \neq j$$

$$\Sigma_1 : \pm(t_i - t_j) = 0, i \neq j$$



# An $SU(5)$ Example

- We take the gauge symmetry on the surface  $S$  to be  $G_S = SU(5)$
- 

$$E_8 \supset SU(5) \times SU(5)_\perp$$

$$248 \rightarrow (24, 1) + (1, 24) + (10, 5) + (\bar{5}, 10) + (\bar{10}, \bar{5}) + (5, \bar{10})$$

- The equations for the curves in terms of the weights  $t_i$  ( $i = 1, \dots, 5$ ,  $\sum t_i = 0$ ), of the 5 representation of  $SU(5)$  are

$$\Sigma_{10} : t_i = 0$$

$$\Sigma_5 : -t_i - t_j = 0, i \neq j$$

$$\Sigma_1 : \pm(t_i - t_j) = 0, i \neq j$$

- The fibration coefficients  $b_k$  are the elementary symmetric polynomials in the weights, of degree  $k$

# An $SU(5)$ Example

- We take the gauge symmetry on the surface  $S$  to be  $G_S = SU(5)$
- 

$$E_8 \supset SU(5) \times SU(5)_\perp$$

$$248 \rightarrow (24, 1) + (1, 24) + (10, 5) + (\bar{5}, 10) + (\bar{10}, \bar{5}) + (5, \bar{10})$$

- The equations for the curves in terms of the weights  $t_i$  ( $i = 1, \dots, 5$ ,  $\sum t_i = 0$ ), of the 5 representation of  $SU(5)$  are

$$\Sigma_{10} : t_i = 0$$

$$\Sigma_5 : -t_i - t_j = 0, i \neq j$$

$$\Sigma_1 : \pm(t_i - t_j) = 0, i \neq j$$

- The fibration coefficients  $b_k$  are the elementary symmetric polynomials in the weights, of degree  $k$
- Non linear  $\rightarrow$  **Monodromies**

# An Example

# An Example

Monodromy must be at least  $Z_2$  for a tree level top yukawa:  $t_1 \leftrightarrow t_2$

## An Example

Monodromy must be at least  $Z_2$  for a tree level top yukawa:  $t_1 \leftrightarrow t_2$

Field	$SU(5) \times SU(5)_\perp$	$SU(5)_\perp$ component
$Q_3, U_3^c, I_3^c$	$(10, 5)$	$t_{1,2}$
$Q_2, U_2^c, I_2^c$	$(10, 5)$	$t_3$
$Q_1, U_1^c, I_1^c$	$(10, 5)$	$t_4$
$D_3^c, L_3$	$(\bar{5}, 10)$	$t_3 + t_5$
$D_2^c, L_2$	$(\bar{5}, 10)$	$t_1 + t_3$
$D_1^c, L_1$	$(\bar{5}, 10)$	$t_1 + t_4$
$H_u$	$(5, \bar{10})$	$-t_1 - t_2$
$H_d$	$(\bar{5}, 10)$	$t_1 + t_4$
$\theta_{ij}$	$(1, 24)$	$t_i - t_j$

**Table:** Matter Fields and their assignments

# Family Symmetries

# Family Symmetries

- The up quark Yukawa operators  $Q_i U_j^c H_u$  require the following flavons to balance the charges

- $$Q_i U_j^c H_u \begin{pmatrix} \theta_{14}^2 & \theta_{13}\theta_{14} & \theta_{14} \\ \theta_{13}\theta_{14} & \theta_{13}^2 & \theta_{13} \\ \theta_{14} & \theta_{13} & 1 \end{pmatrix} \rightarrow Y^u \sim \begin{pmatrix} \epsilon^6 & 3\epsilon^5 & \epsilon^3 \\ 3\epsilon^5 & 9\epsilon^4 & 3\epsilon^2 \\ \epsilon^3 & 3\epsilon^2 & 1 \end{pmatrix}$$

- The choice  $\theta_{13} = 3\epsilon^2$ ,  $\theta_{14} = \epsilon^3$  has been made for the VEVs

# Family Symmetries

- The up quark Yukawa operators  $Q_i U_j^c H_u$  require the following flavons to balance the charges

- $$Q_i U_j^c H_u \begin{pmatrix} \theta_{14}^2 & \theta_{13}\theta_{14} & \theta_{14} \\ \theta_{13}\theta_{14} & \theta_{13}^2 & \theta_{13} \\ \theta_{14} & \theta_{13} & 1 \end{pmatrix} \rightarrow Y^u \sim \begin{pmatrix} \epsilon^6 & 3\epsilon^5 & \epsilon^3 \\ 3\epsilon^5 & 9\epsilon^4 & 3\epsilon^2 \\ \epsilon^3 & 3\epsilon^2 & 1 \end{pmatrix}$$

- The choice  $\theta_{13} = 3\epsilon^2$ ,  $\theta_{14} = \epsilon^3$  has been made for the VEVs
- The down quark Yukawa operators  $Q_i D_j^c H_d$  require the following flavons to balance the charges

- $$Q_i D_j^c H_d \begin{pmatrix} \theta_{54}\theta_{34} & \theta_{54} & \theta_{14} \\ \theta_{54} & \theta_{53} & \theta_{13} \\ \theta_{31}\theta_{54} + \theta_{34}\theta_{51} & \theta_{23} & \theta_{23} \end{pmatrix} \rightarrow Y^d \sim \begin{pmatrix} 0 & \epsilon^3 & \epsilon^3 \\ \epsilon^3 & \epsilon^2 & 3\epsilon^2 \\ 0 & 0 & 1 \end{pmatrix}$$

- The additional choices  $\theta_{53} = \epsilon^2$ ,  $\theta_{54} = \epsilon^3$ ,  $\theta_{31} = 0$  have been made



# Three Different Pictures

# Three Different Pictures

- We can take the gauge symmetry  $G_S$  on the seven brane wrapping a surface  $S$  to be  $E_6$ ,  $SO(10)$  or  $SU(5)$ . We have made complete the connections between these three pictures

# Three Different Pictures

- We can take the gauge symmetry  $G_S$  on the seven brane wrapping a surface  $S$  to be  $E_6$ ,  $SO(10)$  or  $SU(5)$ . We have made complete the connections between these three pictures



$$\begin{aligned}
 E_8 &\supset E_6 \times SU(3)_\perp \\
 &\rightarrow SO(10) \times U(1)_\psi \times SU(3)_\perp \\
 &\rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi \times SU(3)_\perp \\
 E_8 &\supset SO(10) \times SU(4)_\perp \\
 &\rightarrow SU(5) \times U(1)_\chi \times SU(4)_\perp \\
 E_8 &\supset SU(5) \times SU(5)_\perp
 \end{aligned}$$

# Three Different Pictures

- We can take the gauge symmetry  $G_S$  on the seven brane wrapping a surface  $S$  to be  $E_6$ ,  $SO(10)$  or  $SU(5)$ . We have made complete the connections between these three pictures



$$\begin{aligned}
 E_8 &\supset E_6 \times SU(3)_\perp \\
 &\rightarrow SO(10) \times U(1)_\psi \times SU(3)_\perp \\
 &\rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi \times SU(3)_\perp \\
 E_8 &\supset SO(10) \times SU(4)_\perp \\
 &\rightarrow SU(5) \times U(1)_\chi \times SU(4)_\perp \\
 E_8 &\supset SU(5) \times SU(5)_\perp
 \end{aligned}$$

- We assume the breakings proceed via **fluxes**

# Three Different Pictures

- We can take the gauge symmetry  $G_S$  on the seven brane wrapping a surface  $S$  to be  $E_6$ ,  $SO(10)$  or  $SU(5)$ . We have made complete the connections between these three pictures



$$\begin{aligned}
 E_8 &\supset E_6 \times SU(3)_\perp \\
 &\rightarrow SO(10) \times U(1)_\psi \times SU(3)_\perp \\
 &\rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi \times SU(3)_\perp \\
 E_8 &\supset SO(10) \times SU(4)_\perp \\
 &\rightarrow SU(5) \times U(1)_\chi \times SU(4)_\perp \\
 E_8 &\supset SU(5) \times SU(5)_\perp
 \end{aligned}$$

- We assume the breakings proceed via **fluxes**
- $U(1)$ s from perpendicular group: chirality of **complete** GUT reps

# Three Different Pictures

- We can take the gauge symmetry  $G_S$  on the seven brane wrapping a surface  $S$  to be  $E_6$ ,  $SO(10)$  or  $SU(5)$ . We have made complete the connections between these three pictures



$$\begin{aligned}
 E_8 &\supset E_6 \times SU(3)_\perp \\
 &\rightarrow SO(10) \times U(1)_\psi \times SU(3)_\perp \\
 &\rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi \times SU(3)_\perp \\
 E_8 &\supset SO(10) \times SU(4)_\perp \\
 &\rightarrow SU(5) \times U(1)_\chi \times SU(4)_\perp \\
 E_8 &\supset SU(5) \times SU(5)_\perp
 \end{aligned}$$

- We assume the breakings proceed via **fluxes**
- $U(1)$ s from perpendicular group: chirality of **complete** GUT reps
- Fluxes **inside** GUT group

# Flux breaking

# Flux breaking

Flux can be used to break down to the SM, leading to splitting equations

$$10 = \begin{cases} \text{Rep.} & \# \\ n_{3,2}^1 - n_{\bar{3},2}^1 & : M_{10} \\ n_{3,1}^1 - n_{\bar{3},1}^1 & : M_{10} - N \\ n_{1,1}^1 - n_{\bar{1},1}^1 & : M_{10} + N \end{cases} \quad 5 = \begin{cases} \text{Rep.} & \# \\ n_{3,1}^1 - n_{\bar{3},1}^1 & : M_5 \\ n_{1,2}^1 - n_{\bar{1},\bar{2}}^1 & : M_5 + N \end{cases}$$



# Flux breaking

Flux can be used to break down to the SM, leading to splitting equations

$$10 = \begin{cases} \text{Rep.} & \# \\ n_{3,2}^1 - n_{\bar{3},2}^1 & : M_{10} \\ n_{3,1}^1 - n_{\bar{3},1}^1 & : M_{10} - N \\ n_{1,1}^1 - n_{\bar{1},1}^1 & : M_{10} + N \end{cases} \quad 5 = \begin{cases} \text{Rep.} & \# \\ n_{3,1}^1 - n_{\bar{3},1}^1 & : M_5 \\ n_{1,2}^1 - n_{\bar{1},\bar{2}}^1 & : M_5 + N \end{cases}$$

The Ms and Ns are integers given by the flux dotted with the [homology](#) class of the matter curve.

The flux associated with M respects the GUT structure, and so is a flux in the perpendicular U(1)s.

The flux associated with N is a hypercharge flux, and leads to incomplete SU(5) multiplets.

$E_6$ Origin of $SU(5)$ Matter	$U(1)_i$	homology	$U(1)_Y$ -flux	$U(1)$ -flux
$27_{t_1} \supset 16_{t_1} \supset 10_M$	$t_{1,2}$	$\eta - 2c_1 - \chi$	$-N$	$M_{10_1}$
$27_{t_3} \supset 16_{t_3} \supset 10_2$	$t_3$	$-c_1 + \chi_5$	$N_5$	$M_{10_2}$
$78 \supset 16_{t_4} \supset 10_3$	$t_4$	$-c_1 + \chi_7$	$N_7$	$M_{10_3}$
$78 \supset 45_{t_5} \supset 10_4$	$t_5$	$-c_1 + \chi_9$	$N_9$	$M_{10_4}$
$27_{t_3} \supset 10_{t_3} \supset 5_{H_u}$	$-t_1 - t_2$	$-c_1 + \chi$	$N$	$M_{5_{h_u}}$
$27_{t_1} \supset 10_{t_1} \supset 5_1$	$-t_{1,2} - t_3$	$\eta - 2c_1 - \chi$	$-N$	$M_{5_1}$
$27_{t_1} \supset 10_{t_1} \supset 5_2$	$-t_{1,2} - t_4$	$\eta - 2c_1 - \chi$	$-N$	$M_{5_2}$
$27_{t_1} \supset 16_{t_1} \supset 5_3$	$-t_{1,2} - t_5$	$\eta - 2c_1 - \chi$	$-N$	$M_{5_3}$
$27_{t_3} \supset 10_{t_3} \supset 5_4$	$-t_3 - t_4$	$-c_1 + \chi - \chi_9$	$N - N_9$	$M_{5_4}$
$27_{t_3} \supset 16_{t_3} \supset 5_5$	$-t_3 - t_5$	$-c_1 + \chi - \chi_7$	$N - N_7$	$M_{5_{h_d}}$
$78 \supset 16_{t_4} \supset 5_6$	$-t_4 - t_5$	$-c_1 + \chi - \chi_5$	$N - N_5$	$M_{5_6}$

**Table:** Field representation content under  $SU(5) \times U(1)_{t_i}$ , their homology class and flux restrictions. For convenience, only the properties of 10, 5 are shown.  $\overline{10}, \overline{5}$  are characterized by opposite values of  $t_i \rightarrow -t_i$  etc. Note that the fluxes satisfy  $N = N_5 + N_7 + N_9$  and  $\sum_i M_{10_i} + \sum_j M_{5_j} = 0$  while  $\chi = \chi_5 + \chi_7 + \chi_9$ .

We know about the origins of the singlets also

Singlet	$Q_\chi$	$Q_\psi$	Representations
$\theta_{12}$	0	0	SO(10) singlet in 78
$\theta_{13}$	0	0	$45 \subset 78$
$\theta_{14}$	0	4	SO(10) singlet in $27_{t_{1,2}}$
$\theta_{15}$	-5	1	$16_{t_{1,2}} \subset 27_{t_{1,2}}$
$\theta_{34}$	0	4	SO(10) singlet in $27_{t_3}$
$\theta_{35}$	-5	-1	$16_{t_3} \subset 27_{t_3}$
$\theta_{45}$	-5	-3	$16_{t_4} \subset 78$

# A realistic model based on $E_6$

# A realistic model based on $E_6$

The only  $E_6$  allowed trilinear term in the superpotential is  $27_{t_1} 27_{t_1} 27_{t_3}$

# A realistic model based on $E_6$

The only  $E_6$  allowed trilinear term in the superpotential is  $27_{t_1} 27_{t_1} 27_{t_3}$

Anomalies can be cancelled by having complete 27 representations of  $E_6$

# A realistic model based on $E_6$

The only  $E_6$  allowed trilinear term in the superpotential is  $27_{t_1} 27_{t_1} 27_{t_3}$

Anomalies can be cancelled by having complete 27 representations of  $E_6$

$E_6, SO(10), SU(5)$	Charge	$N_Y$	$M_{U(1)}$	SM particle content
$27_{t_1}, 16_{t_1}, \bar{5}_3$	$t_1 + t_5$	$\tilde{N}$	$-M_{5_3}$	$-M_{5_3} d^c + (-M_{5_3} + \tilde{N})L$
$27_{t_1}, 16_{t_1}, 10_M$	$t_1$	$-\tilde{N}$	$-M_{5_3}$	$-M_{5_3} Q + (-M_{5_3} + \tilde{N})u^c + (-M_{5_3} - \tilde{N})e^c$
$27_{t_1}, 16_{t_1}, \theta_{15}$	$t_1 - t_5$	0	$-M_{5_3}$	$-M_{5_3} \nu^c$
$27_{t_1}, 10_{t_1}, 5_1$	$-t_1 - t_3$	$-\tilde{N}$	$-M_{5_3}$	$-M_{5_3} D + (-M_{5_3} - \tilde{N})H_u$
$27_{t_1}, 10_{t_1}, \bar{5}_2$	$t_1 + t_4$	$\tilde{N}$	$-M_{5_3}$	$-M_{5_3} \bar{D} + (-M_{5_3} + \tilde{N})H_d$
$27_{t_1}, 1_{t_1}, \theta_{14}$	$t_1 - t_4$	0	$-M_{5_3}$	$-M_{5_3} S$
$27_{t_3}, 16_{t_3}, \bar{5}_5$	$t_3 + t_5$	$-\tilde{N}$	$M_{5_{H_u}}$	$M_{5_{H_u}} d^c + (M_{5_{H_u}} - \tilde{N})L$
$27_{t_3}, 16_{t_3}, 10_2$	$t_3$	$\tilde{N}$	$M_{5_{H_u}}$	$M_{5_{H_u}} Q + (M_{5_{H_u}} - \tilde{N})u^c + (M_{5_{H_u}} + \tilde{N})e^c$
$27_{t_3}, 16_{t_3}, \theta_{35}$	$t_3 - t_5$	0	$M_{5_{H_u}}$	$M_{5_{H_u}} \nu^c$
$27_{t_3}, 10_{t_3}, 5_{H_u}$	$-2t_1$	$\tilde{N}$	$M_{5_{H_u}}$	$M_{5_{H_u}} D + (M_{5_{H_u}} + \tilde{N})H_u$
$27_{t_3}, 10_{t_3}, \bar{5}_4$	$t_3 + t_4$	$-\tilde{N}$	$M_{5_{H_u}}$	$M_{5_{H_u}} \bar{D} + (M_{5_{H_u}} - \tilde{N})H_d$
$27_{t_3}, 1_{t_3}, \theta_{34}$	$t_3 - t_4$	0	$M_{5_{H_u}}$	$M_{5_{H_u}} S$

# Spectrum



# Spectrum

- Choosing the case  $M_{5_3} = -3$ ,  $M_{5_{H_u}} = 0$  and  $\tilde{N} = 1$ , the spectrum is
- $3[Q, u^c, d^c, L, e^c, \nu^c]_{16}$
- $3[H_u, D, H_d, \overline{D}]_{10}$
- $3[S]_1$
- $L + \overline{L}, e^c + \overline{e^c}, u^c + \overline{u^c}, H_d + \overline{H_d}$

# Spectrum

- Choosing the case  $M_{5_3} = -3$ ,  $M_{5_{H_u}} = 0$  and  $\tilde{N} = 1$ , the spectrum is
- $3[Q, u^c, d^c, L, e^c, \nu^c]_{16}$
- $3[H_u, D, H_d, \bar{D}]_{10}$
- $3[S]_1$
- $L + \bar{L}, e^c + \bar{e}^c, u^c + \bar{u}^c, H_d + \bar{H}_d$
- With this choice, however,  $H_d$  comes from  $27_{t_1}$  and so down quark masses are forbidden at tree level!

# Spectrum

- Choosing the case  $M_{5_3} = -3$ ,  $M_{5_{H_u}} = 0$  and  $\tilde{N} = 1$ , the spectrum is
- $3[Q, u^c, d^c, L, e^c, \nu^c]_{16}$
- $3[H_u, D, H_d, \bar{D}]_{10}$
- $3[S]_1$
- $L + \bar{L}, e^c + \bar{e}^c, u^c + \bar{u}^c, H_d + \bar{H}_d$
- With this choice, however,  $H_d$  comes from  $27_{t_1}$  and so down quark masses are forbidden at tree level!
- $\Rightarrow$  Slightly modified structure:

$$M_{10_M} = -M_{5_3} = 4,$$

$$M_{5_1} = -M_{5_2} = 3$$

$$M_{10_2} = -M_{5_5} = -1,$$

$$M_{5_4} = M_{H_u} = 0,$$

$$M_{\theta_{15}} = 2,$$

$$\tilde{N} = 1$$

$E_6$	$SO(10)$	$SU(5)$	$N_Y$	$M_{U(1)}$	SM particle content	Low energy
$27_{t'_1}$	16	$\bar{5}_3$	1	4	$4d^c + 5L$	$3d^c + 3L$
$27_{t'_1}$	16	$10_M$	-1	4	$4Q + 5u^c + 3e^c$	$3Q + 3u^c + 3e^c$
$27_{t'_1}$	16	$\theta_{15}$	0	3	$3\nu^c$	-
$27_{t'_1}$	10	$5_1$	-1	3	$3D + 2H_u$	-
$27_{t'_1}$	10	$\bar{5}_2$	1	3	$3\bar{D} + 4H_d$	$H_d$
$27_{t'_3}$	16	$\bar{5}_5$	-1	-1	$\bar{d}^c + 2\bar{L}$	-
$27_{t'_3}$	16	$10_2$	1	-1	$\bar{Q} + 2\bar{u}^c$	-
$27_{t'_3}$	16	$\theta_{35}$	0	0	-	-
$27_{t'_3}$	10	$5_{H_u}$	1	0	$H_u$	$H_u$
$27_{t'_3}$	10	$\bar{5}_4$	-1	0	$\bar{H}_d$	-
$27_{t'_3}$	1	$\theta_{34}$	0	1	$\theta_{34}$	-
-	1	$\theta_{31}$	0	4	$\theta_{31}$	-
-	1	$\theta_{53}$	0	1	$\theta_{53}$	-
-	1	$\theta_{14}$	0	3	$\theta_{14}$	-
-	1	$\theta_{45}$	0	2	$\theta_{45}$	-



- The vector pairs with components in the  $27_{t_1}$  and  $27_{t_3}$  multiplets are removed by introducing  $\theta_{31}$  which is a singlet of  $E_6$  and has couplings:

$$\theta_{31} 27_{t'_1} \overline{27_{t'_3}} = \theta_{31} Q \overline{Q} + \theta_{31} (2u^c)(2\overline{u^c}) + \theta_{31} d^c \overline{d^c} + \theta_{31} (2L)(2\overline{L}) + \theta_{31} H_d \overline{H_d}$$

- The vector pairs with components in the  $27_{t_1}$  and  $27_{t_3}$  multiplets are removed by introducing  $\theta_{31}$  which is a singlet of  $E_6$  and has couplings:

$$\theta_{31} 27_{t_1} \overline{27_{t_3}} = \theta_{31} Q \overline{Q} + \theta_{31} (2u^c)(2\overline{u^c}) + \theta_{31} d^c \overline{d^c} + \theta_{31} (2L)(2\overline{L}) + \theta_{31} H_d \overline{H_d}$$

- If  $\theta_{31}$  gets a large VEV these vector states get large masses as required.

- The vector pairs with components in the  $27_{t_1}$  and  $27_{t_3}$  multiplets are removed by introducing  $\theta_{31}$  which is a singlet of  $E_6$  and has couplings:

$$\theta_{31} 27_{t_1} \overline{27}_{t_3} = \theta_{31} Q \overline{Q} + \theta_{31} (2u^c)(2\overline{u}^c) + \theta_{31} d^c \overline{d}^c + \theta_{31} (2L)(2\overline{L}) + \theta_{31} H_d \overline{H}_d$$

- If  $\theta_{31}$  gets a large VEV these vector states get large masses as required.
- To remove the remaining exotics we introduce  $\theta_{34}$  which has the couplings :

$$\theta_{34} 5_1 \overline{5}_2 = \theta_{34} [3D + 2H_u][3\overline{D} + 3H_d] = \theta_{34} [3(D\overline{D})] + \theta_{34} [2(H_u H_d)]$$

If it too acquires a large VEV it generates large mass to the three copies of  $D + \overline{D}$  (solving the doublet-triplet splitting problem) and two families of Higgs  $H_u, H_d$ , leaving just the MSSM spectrum



# Singlet VEVs

# Singlet VEVs

- To determine the large singlet VEVs we consider the  $F$ - and  $D$ -flatness conditions, which leads to

$$\begin{aligned}|\theta_{53}|^2 &= X \\ |\theta_{34}|^2 &= 2X \\ |\theta_{31}|^2 &= 37X\end{aligned}$$

where  $X = \frac{g_s^2 M_S^2}{192\pi^2}$

# Singlet VEVs

- To determine the large singlet VEVs we consider the  $F$ - and  $D$ -flatness conditions, which leads to

$$\begin{aligned} |\theta_{53}|^2 &= X \\ |\theta_{34}|^2 &= 2X \\ |\theta_{31}|^2 &= 37X \end{aligned}$$

where  $X = \frac{g_s^2 M_S^2}{192\pi^2}$

- The singlet  $\theta_{14}$  gets a TeV VEV in order to generate the  $\mu$  term  $\theta_{14} H_u H_d$ .

# Baryon- and lepton-number violating terms

# Baryon- and lepton-number violating terms

- The R-parity violating superpotential couplings  $u^c d^c d^c$ ,  $Q d^c L$ ,  $Le^c L$ ,  $\kappa L H_u$  are not allowed because of the underlying  $27_{t'_1} \rightarrow -27_{t'_1}$  symmetry. Under this symmetry the matter fields and  $H_d$  are odd but  $H_u$  is even.

# Baryon- and lepton-number violating terms

- The R-parity violating superpotential couplings  $u^c d^c d^c$ ,  $Qd^c L$ ,  $Le^c L$ ,  $\kappa L H_u$  are not allowed because of the underlying  $27_{t'_1} \rightarrow -27_{t'_1}$  symmetry. Under this symmetry the matter fields and  $H_d$  are odd but  $H_u$  is even.
- Dimension 5 terms correspond to the superpotential terms  $QQQL$  and  $u^c u^c d^c e^c$ . However they are forbidden by the  $U(1)$  symmetries that originate in the underlying  $E_6$ .

# Baryon- and lepton-number violating terms

- The R-parity violating superpotential couplings  $u^c d^c d^c$ ,  $Qd^c L$ ,  $Le^c L$ ,  $\kappa LH_u$  are not allowed because of the underlying  $27_{t'_1} \rightarrow -27_{t'_1}$  symmetry. Under this symmetry the matter fields and  $H_d$  are odd but  $H_u$  is even.
- Dimension 5 terms correspond to the superpotential terms  $QQQL$  and  $u^c u^c d^c e^c$ . However they are forbidden by the  $U(1)$  symmetries that originate in the underlying  $E_6$ .
- Allowing for arbitrary singlet fields to acquire VEVs the dangerous the baryon- and lepton-number violating operators arise through the terms  $\theta_{15} LH_u$ ,  $(\theta_{31}\theta_{45} + \theta_{41}\theta_{35})10_M \overline{5}_3^2$  and  $\theta_{31}\theta_{41}10_M^3 \overline{5}_3$ .

# Baryon- and lepton-number violating terms

- The R-parity violating superpotential couplings  $u^c d^c d^c$ ,  $Qd^c L$ ,  $Le^c L$ ,  $\kappa LH_u$  are not allowed because of the underlying  $27_{t'_1} \rightarrow -27_{t'_1}$  symmetry. Under this symmetry the matter fields and  $H_d$  are odd but  $H_u$  is even.
- Dimension 5 terms correspond to the superpotential terms  $QQQL$  and  $u^c u^c d^c e^c$ . However they are forbidden by the  $U(1)$  symmetries that originate in the underlying  $E_6$ .
- Allowing for arbitrary singlet fields to acquire VEVs the dangerous the baryon- and lepton-number violating operators arise through the terms  $\theta_{15} LH_u$ ,  $(\theta_{31}\theta_{45} + \theta_{41}\theta_{35})10_M \overline{5}_3^2$  and  $\theta_{31}\theta_{41}10_M^3 \overline{5}_3$ .
- $\Rightarrow$  provided  $\theta_{15}$ ,  $\theta_{41}$  and  $\theta_{45}$  do *not* acquire VEVs these dangerous terms will not arise.



# Summary

# Summary

- F-theory Guts can provide realistic models of flavour

# Summary

- F-theory Guts can provide realistic models of flavour
- A useful framework for model building can be built by studying the relationships between different GUT theories

# Summary

- F-theory Guts can provide realistic models of flavour
- A useful framework for model building can be built by studying the relationships between different GUT theories
- This framework has been used to develop a model which stabilises the proton and forbids R-parity violating operators