

$SO(10)$ inspired extended GMSB models

Tomasz Jeliński

Department of Field Theory and Particle Physics

University of Silesia, Katowice, Poland


CORFU2014, 3-14 September 2014

- GUT model: $SO(10)$ + GMSB + messenger couplings
- messenger-matter mixing & $y_{t,b,\tau}$ running
- $y_t - y_b - y_\tau$ unification in minimal $SO(10)$ model



1. LHC vs. MSSM

What do the LHC searches tell us about MSSM?

- no SUSY signal so far
- relevant exclusions only for 1st and 2nd family
- still \tilde{Q}_3, \dots can be as light as 500 GeV



2. Limits on stop mass



3. LHC vs. MSSM


What do the LHC searches tell us about MSSM?

- no SUSY signal so far
- relevant exclusions only for 1st and 2nd family
- still \tilde{Q}_3, \dots can be as light as 500 GeV

BUT important information comes from Higgs mass measurement:

- $m \sim 125$ GeV \rightarrow need for large loop corrections

ASSUME other MSSM Higgses are much heavier and masses of $\tilde{Q}_{1,2}$ and \tilde{g} are bigger than 1.8 TeV.



4. 1-loop corrections to m_{h^0}

- dominant contribution from top quarks and stops (due to $y_t \sim 1$):

$$\Delta(m_{h^0}^2) = \frac{h^0}{-} - \text{[solid loop with } t \text{]} - \text{[dashed loop with } \tilde{t} \text{]} + \frac{h^0}{-} - \text{[dashed loop with } \bar{\tilde{t}} \text{]} -$$

$$m_{h^0}^2 = m_Z^2 \cos^2 2\beta + \frac{3m_t^4}{4\pi^2 v^2} \left[\ln \frac{M_S^2}{m_t^2} + \frac{X_t^2}{M_S^2} \left(1 - \frac{X_t^2}{12M_S^2} \right) \right] \approx (125 \text{ GeV})^2,$$


$$M_S = \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$$

$$X_t = A_t - \mu \cot \beta$$

Large A -terms or heavy stops!

A-terms:


$$V_{\text{soft}} \supset y_t \textcolor{blue}{A}_t H_u \tilde{Q}_3 \tilde{\bar{U}}_3 \longrightarrow y_t \textcolor{blue}{A}_t h_0 \tilde{t}_1 \tilde{t}_2$$



Draper et al. 1112.3068

5. A -terms in GMSB

- in GMSB models A -terms = 0 at messenger scale



Draper et al. 1112.3068


$$\mu \frac{dA_t}{d\mu} \sim y_t^2 A_t + g_3^2 M_3$$

- hard to reconcile
 - $m_{h^0} \gtrsim 123 \text{ GeV}$
 - pure GMSB mechanism
 - light stops
 - $M_{\tilde{g}} \lesssim 2.5 \text{ GeV}$

- large A -terms at M ?

6. How to generate large A -terms?

- value of A -term gives initial condition for RGE evolution



Draper et al. 1112.3068


$$\mu \frac{dA_t}{d\mu} \sim y_t^2 A_t + g_3^2 M_3$$

- heavy \tilde{g} and RGE evolution from $M \gtrsim 10^{14}$ GeV
- or large A -terms at M

- how to get A -terms in GUT model?

Extended GMSB models (EGMSB)

7. SUSY breaking mediation



- singlet $\langle X \rangle = M + \theta^2 F \rightarrow$ spontaneous **SUSY breaking**

$$\xi = \frac{F}{M} \sim 10^5 \text{ GeV}$$

- messengers have large masses e.g. $M \sim 10^8 - 10^{14} \text{ GeV}$
- mediation = interactions between Y, \bar{Y} and other fields
- assumption: all messengers couple to the spurion X in the same way

$$XY_a \bar{Y}_a$$

and $M \gtrsim 10^8 \text{ GeV} \rightarrow$ 1-loop soft masses negligible

8. SO(10) GUT model

- at $M_{GUT} \sim 10^{16} \text{GeV}$: $SO(10) \rightarrow SU(5) \times U(1)_\chi \rightarrow \dots$
- chiral matter Φ

$$H_{10} : \quad 10 \rightarrow 5_2 + \bar{5}_{-2}, \quad \phi_{16} : \quad 16 \rightarrow 10_{-1} + \bar{5}_3 + 1_{-5}$$

- **messengers** $Y = (Y_{16}, Y_{\bar{16}})$

$$W = \textcolor{red}{y} H_{10} \phi_{16} \phi_{16} + \textcolor{blue}{h} H_{10} \phi_{16} Y_{16} + H_{10} Y_{16} Y_{16} + H_{10} Y_{\bar{16}} Y_{\bar{16}} + \underbrace{\frac{1}{2} M Y_{16} Y_{\bar{16}}}_{\text{mass term}}$$


- $\textcolor{red}{y} = y_t(t_{GUT}) = y_b(t_{GUT}) = y_\tau(t_{GUT})$
- $\phi_1 = N_R$, Y_1 and Higgs triplets masses $\sim M_{GUT}$
- only couplings to 3rd generation

$$W_\lambda = \frac{1}{6} \lambda_{ijk} \Phi_i \Phi_j \Phi_k = \frac{1}{6} y_{abc} \Phi_a \Phi_b \Phi_c + \frac{1}{2} h_{ab} \Phi_a \Phi_b Y + h_a \Phi_a Y Y + \eta Y Y Y$$

What are the soft terms?

9. Trilinear terms in EGMSSB models

$$W = \frac{1}{6} y_{abc} \Phi_a \Phi_b \Phi_c + \frac{1}{2} h_{ab} \Phi_a \Phi_b Y + h_a \Phi_a Y Y + \eta Y Y Y$$



$$V \supset T_{abc} \tilde{\Phi}_a \tilde{\Phi}_b \tilde{\Phi}_c, \quad T_{abc} = -\frac{\xi}{16\pi^2} [C_a h_{ad} h_{de}^* y_{ebc} + \dots] + (a \leftrightarrow b) + (a \leftrightarrow c)$$

- T_{abc} are ‘partially aligned’ to MSSM Yukawas y_{abc}

10. A -terms in EGMSB models

$$V \supset H_u \tilde{Q}(T_u) \tilde{\bar{U}} + H_d \tilde{Q}(T_d) \tilde{\bar{D}} + H_d \tilde{L}(T_e) \tilde{\bar{E}}$$

- $(T_{u,d,e})_{33} =: y_{t,b,\tau} A_{t,b,\tau}$

$$A_{t,b,\tau} \approx -\frac{\xi}{16\pi^2} C^{(t,b,\tau)} |h|^2 \quad C^{(t,b,\tau)} = 10, 12, 11$$

- A -terms
 - relevant to the m_{h^0}
 - may also lead to CCB when

$$A_f^2 > 3(m_{\tilde{f}_L}^2 + m_{\tilde{f}_R}^2 + \mu^2 + m_{H_u}^2)$$

- affect sfermion masses $m_{\tilde{f}_{1,2}}$


$$(\tilde{f}_L^* \ \tilde{f}_R^*) \begin{pmatrix} m_{\tilde{f}_{LL}}^2 & m_f(A_f - \mu \tan \beta^{\pm 1}) \\ m_f(A_f - \mu \tan \beta^{\pm 1}) & m_{\tilde{f}_{RR}}^2 \end{pmatrix} \begin{pmatrix} \tilde{f}_L \\ \tilde{f}_R \end{pmatrix}$$

$\rightarrow \tilde{f}_1$ may be tachyonic


11. Soft masses in EGMSSB models

- 2-loop contributions to soft masses

$$W_Y = h^{(I)} \Phi Y Y + h^{(II)} \Phi \Phi Y$$



$$\sim h^4, h^3 y, h^2 y^2$$




$$\sim h^2 g^2$$

$$m_{\tilde{\Phi}, h}^2 \sim \frac{\xi^2}{(4\pi)^4} (h^4 + h^3 y - h^2 y^2 - h^2 g^2)$$

12. 2-loop soft masses induced by YYY

$$W_Y = h_i^{(I)} \Phi_i Y Y + h_{ij}^{(II)} \Phi_i \Phi_j Y + \eta Y Y Y$$



$$m_{\tilde{\Phi}, \eta}^2 \sim \frac{\xi^2}{(4\pi)^4} (\eta^2 h^2 + \eta h^3 + \eta h^2 y)$$

Remark: η are relevant only if a model contains both $5 + \bar{5}$ and $10 + \bar{10}$ messengers

13. Kinetic mixing

- fields Y, ϕ with the same charges can mix: $\phi_{16} \leftrightarrow Y_{16}$

$$Q \leftrightarrow Y_Q, \bar{U} \leftrightarrow Y_{\bar{U}}, \dots \quad (\text{in some models: } H_d \leftrightarrow L \leftrightarrow Y_L)$$

- superpotential and Kähler potential K at scale $t = \log \mu$

$$W = \frac{1}{6} \lambda_{ijk} \Phi_i \Phi_j \Phi_k + \frac{1}{2} M_{ij} \Phi_i \Phi_j, \quad K = \Phi_i^\dagger Z_{ij}(t) \Phi_j, \quad Z = Z^\dagger, \quad Z > 0$$

- couplings $\tilde{\lambda}(t)$ and **masses** of canonically normalized fields $\tilde{\Phi}_i = Z_{ij}^{-1/2} \Phi_j$

$$\tilde{\lambda}_{ijk}(t) = \lambda_{i'j'k'} Z_{i'i}^{-1/2} Z_{j'j}^{-1/2} Z_{k'k}^{-1/2}, \quad \tilde{M}_{ij}(t) = M_{i'j'} Z_{i'i}^{-1/2} Z_{j'j}^{-1/2}$$

- RGE evolution of Z (re)introduces mixing mass terms!

e.g.

$$W = \tilde{M}_1 \tilde{Y}_{16} \tilde{Y}_{\overline{16}} + \tilde{M}_2 \tilde{\phi}_{16} \tilde{Y}_{\overline{16}} + \dots$$

- important for decouplings and running Yukawa (couplings between light states)!

14. Decoupling and running

$$W = \frac{1}{6}\lambda_{ijk}\Phi_i\Phi_j\Phi_k + \frac{1}{2}M_{ij}\Phi_i\Phi_j, \quad K = \Phi_i^\dagger Z_{ij}(t)\Phi_j, \quad Z = Z^\dagger, \quad Z > 0$$

- method 1 - rotate $\tilde{\Phi} = Z^{-1/2}\Phi$ such that at least one field in representation 16 of $SO(10)$ is massless
- **method 2** - instead of computing $Z^{-1/2}$ and then rotating $\tilde{\Phi}$ use Cholesky decomposition of Z :


$$Z = V^\dagger V, \quad \tilde{\Phi} = \begin{pmatrix} \tilde{\phi} \\ \tilde{Y} \end{pmatrix} = \underbrace{\begin{pmatrix} * & * \\ 0 & * \end{pmatrix}}_V \begin{pmatrix} \phi \\ Y \end{pmatrix}$$

$$\tilde{\lambda}_{ijk}(t) = \lambda_{i'j'k'} V_{i'i}^{-1} V_{j'j}^{-1} V_{k'k}^{-1}$$

- one can check that

$$\tilde{\lambda}_{abc}(t) = \frac{\lambda_{abc}}{\sqrt{\color{blue}Z_{aa}\color{blue}Z_{bb}\color{blue}Z_{cc}}}, \quad \phi_a - \text{light fields}$$

15. Standard RGE vs. Z



$$W = y_t H_u Q \bar{U} + h_t H_u Q Y_{\bar{U}} + M Y_U Y_{\bar{U}}, \quad y_t = 0.7, h_t = 0.4, M' = 10^{10} \text{ GeV}$$

16. Evolution of Z from GUT scale t_{GUT}

- RGE for $Z(t)$ with boundary condition $Z(t_{GUT}) = 1$

$$\frac{d}{dt} Z_{ij} = -\frac{1}{8\pi^2} \left(\frac{1}{2} d_{kl} \lambda_{ikl}^* Z_{km}^{-1*} Z_{ln}^{-1*} \lambda_{jmn} - 2C_{ij}^{(r)} Z_{ij} g_r^2 \right)$$

d_{kl} and C_{ij}^r - group theory factors

- solve numerically or use [approximate solution](#):

$$Z_{ij}(t) = 1 + Z_{ij}^{(1)}(t - t_{GUT}) + \frac{1}{2!} Z_{ij}^{(2)}(t - t_{GUT})^2 + \dots$$

- to compute $Z^{(n)}$ one needs all $Z^{(k)}$, $k < n$
- $Z^{(n)}$ are expressed in terms of $\epsilon = \ln 10/16\pi^2$, λ_{ijk} , d_{kl} , g_{GUT} and $\beta_{g_r}(t_{GUT})$

17. Z for $SO(10)$ model

- RGE for $Z(t)$ with boundary condition $Z(t_{GUT}) = 1$


$$\frac{d}{dt}Z_{ij} = -\frac{1}{8\pi^2} \left(\frac{1}{2}d_{kl}\lambda_{ikl}^*Z_{km}^{-1*}Z_{ln}^{-1*}\lambda_{jmn} - 2C_{ij}^{(r)}Z_{ij}g_r^2 \right)$$

- e.g for $SO(10)$ minimal model with superpotential

$$W = \textcolor{red}{y}H_{10}\phi_{16}\phi_{16} + \textcolor{blue}{h}H_{10}\phi_{16}Y_{16} + MY_{16}Y_{\overline{16}}$$

$$\begin{aligned} Z_{H_u H_u} &= 1 + \frac{6}{5}\epsilon[3g_{GUT}^2 - 5(2h^2 - y^2)](t - t_{GUT}) \\ &+ \frac{24}{25}\epsilon^2[29g_{GUT}^2 + 35(2h^2 + y^2) - 25(2h^4 + 4h^2y^2 + y^4)](t - t_{GUT})^2 \\ &+ \dots \end{aligned}$$

18. $t - b - \tau$ unification



$$W = y H_{10} \phi_{16} \phi_{16} + h H_{10} \phi_{16} Y_{16} + M Y_{16} Y_{\overline{16}}, \quad y = 0.7, \quad h = 0.4, \quad M' = 10^{10} \text{ GeV}$$

19. Phenomenology of minimal $SO(10)$ model

- scan over parameters

$$8 < t_M < 14, \quad 0.6 < y < 0.9, \quad 0 < h < 1.2$$

- check low-energy constraints

$$m_{h^0} \approx 125 \text{ GeV}, \quad M_{\tilde{g}, \tilde{q}_{1,2}} > 1.8 \text{ TeV}, \quad \text{UFB/CCB}, \quad a_\mu, \quad \dots$$

- for moderate $\tan \beta \sim 20$: no tachyons, $\tilde{\tau}$ is NLSP, but threshold corrections to $y_{b,\tau} \sim 200\%$ or more are needed
- to get $\sim 20\%$ threshold correction for y_b one has to fix $\tan \beta \sim 45 \rightarrow$ **tachyonic** $\tilde{\tau}$
- to avoid instabilities of the potential one could extend spectrum or allow additional messenger couplings

20. Conclusions

- messenger couplings λ not only generate soft terms but can also lead to kinetic mixing
- wave-function renormalization Z is a handy tool to analyze RG flow of Yukawas; this method can be implemented in a similar way at 2-loop level
- phenomenology of the simplest $SO(10)$ model is spoiled by tachionic $\tilde{\tau}$
→ extend spectrum or allow additional couplings