
F-theory models and their predictions for new physics phenomena

Symmetries for Neutrino Physics

Exotic states and Resonances

George K. Leontaris

The University of Ioannina

Ιωάννινα

GREECE

Outline of the Talk

- ▲ \mathcal{F} -Theory: A few basic notions...
- ▲ Model building with F-theory
- ▲ $SU(5) \times PSL_2(p)$ and Neutrinos ...
- ▲ New Physics Implications of ~~750~~GeV ... or rather \geq *few TeV resonances*
- ▲ Concluding Remarks

PART – I

F-Theory

why ?

★ Advantages

Consistent framework for unification

Calculability

testable predictions

Basic features of F-theory:

- ★ Geometrization of Type II-B String Theory
- ★ Elliptically fibred 8-dimensional compact space
- ★ Fibration described by a simple well known model (*Weierstraß model*)

A

... a short geometric description of the fibration ...

Any cubic equation with a rational point can be written in:

★ Weierstraß form:

$$y^2 = x^3 + fx + g$$

▲ Two important quantities characterising elliptic curves:

1. The Discriminant:

$$\Delta = 4f^3 + 27g^2$$

... classifies the curves with respect to its singularities


2. The j -invariant function:

$$j = 4 \frac{(24f)^3}{4f^3 + 27g^2}$$

... takes the same value for equivalent elliptic curves

basic ingredients: the elliptic curve equ and its discriminant:


$$y^2 = x^3 + fx + g, \quad \Delta = 4f^3 + 27g^2$$




non-singular $\Delta \neq 0 \longleftrightarrow$ Elliptic Curves \longleftrightarrow singular $\Delta = 0$

$$y^2 = x^3 + fx + g$$

Real





Complex




$$\Delta \neq 0$$

non-singular elliptic curve



$$\Delta = 0$$

singular elliptic curve



A torus cut along the two circles is topologically equivalent to a parallelogram.

Described by **Complex Modulus**: $\tau = \alpha + \beta i$.

★ F-theory ★

(Vafa hep-th/9602022)



Geometrisation of Type II-B superstring

II-B: closed string spectrum obtained by combining left and right moving open strings with NS and R-boundary conditions:

$$(NS_+, NS_+), (R_-, R_-), (NS_+, R_-), (R_-, NS_+)$$

Bosonic spectrum:

(NS_+, NS_+) : graviton, dilaton and 2-form Kalb-Ramond-field:

$$g_{\mu\nu}, \phi, B_{\mu\nu} \rightarrow B_2$$

(R_-, R_-) : scalar, 2- and 4-index fields (p -form potentials)

$$C_0, C_{\mu\nu}, C_{\kappa\lambda\mu\nu} \rightarrow C_p, p = 0, 2, 4$$

Definitions (*F*-theory bosonic part)

1. String coupling: $g_s = e^{-\phi}$
2. Combining the two scalars C_0, ϕ to one *modulus*:

$$\tau = C_0 + i e^{\phi} \rightarrow C_0 + \frac{i}{g_s}$$

(recall that τ can describe a torus)




1. Theory can be described by consistent properly invariant action (see for example arXiv:0803.1194)
2. ... gives the correct EoM
3. Consistent with $N = 1$ Supersymmetry

FIBRATION

- ▲ \Rightarrow 6-d compact space described by 3-complex dim. manifold \mathcal{B}_3
- ▲ \Rightarrow At each point on \mathcal{B}_3 assign a torus with modulus:

$$\tau = C_0 + i/g_s$$



\Rightarrow F-theory defined on $\mathcal{R}^{3,1} \times \mathcal{X}$

\mathcal{X} , is called elliptically fibered **CY** 4-fold over \mathcal{B}_3

Elliptic Fibration

described by Weierstraß Equation

$$y^2 = x^3 + f(w)xz^4 + g(w)z^6$$

For each point of B_3 , the above equation describes a torus

1. x, y, z homogeneous coordinates
2. $f(w), g(w) \rightarrow 8^{th}$ and 12^{th} degree polynomials.
3. Discriminant

$$\Delta(w) = 4f^3 + 27g^2$$

Fiber singularities at zeros of Discriminant.

$$\Delta(w) = 0 \rightarrow 24 \text{ roots } w_i$$

⇓

Kodaira classification:

- Type of Manifold **singularity** is specified by the **vanishing order** of $f(w)$, $g(w)$ and $\Delta(w)$
- **Geometric Singularities** classified in terms of ADE Lie groups (Kodaira ~ 1960...).

Interpretation of geometric singularities



CY_4 -Singularities \iff gauge symmetries

$$\text{Groups} \rightarrow \begin{cases} SU(n) \\ SO(m) \\ \mathcal{E}_n \end{cases}$$

Example:

$$f = w^3(b_3 + b_4w + \dots), \quad g = w^4(c_4 + c_5w + \dots), \quad \Delta = w^8(d_8 + d_9w + \dots) \rightarrow \mathcal{E}_6$$

$\text{ord}(f)$	$\text{ord}(g)$	$\text{ord}(\Delta)$	fiber type	Singularity
0	0	n	I_n	A_{n-1}
≥ 1	1	2	II	none
1	≥ 2	3	III	A_1
≥ 2	2	4	IV	A_2
2	≥ 3	$n + 6$	I_n^*	D_{n+4}
≥ 2	3	$n + 6$	I_n^*	D_{n+4}
≥ 3	4	8	IV^*	E_6
3	≥ 5	9	III^*	E_7
≥ 4	5	10	II^*	E_8

Table 1: Vanishing order of the polynomials f, g and the discriminant Δ .
(The **Kodaira** classification)

Tate's Algorithm

$$y^2 + \alpha_1 x y z + \alpha_3 y z^3 = x^3 + \alpha_2 x^2 z^2 + \alpha_4 x z^4 + \alpha_6 z^6$$

Table: Classification of Elliptic Singularities w.r.t. vanishing order of Tate's form coefficients α_i :

Group	α_1	α_2	α_3	α_4	α_6	Δ
$SU(2n)$	0	1	n	n	$2n$	$2n$
$SU(2n + 1)$	0	1	n	$n + 1$	$2n + 1$	$2n + 1$
$SU(5)$	0	1	2	3	5	5
$SO(10)$	1	1	2	3	5	7
\mathcal{E}_6	1	2	3	3	5	8
\mathcal{E}_7	1	2	3	3	5	9
\mathcal{E}_8	1	2	3	4	5	10

Basic ingredient in F-theory:

$D7$ - brane

GUTs are associated with 7-branes wrapping certain classes of 'internal' 2-complex dim. surface:

$$\mathbf{S} \subset B_3$$

▲ Gauge symmetry embedded in maximal exceptional group:

$$\mathcal{E}_8 \rightarrow \mathbf{G}_{GUT} \times \mathcal{C}$$

▲ $G_{GUT} = SU(5), SO(10), \dots$

★ \mathcal{C} Group can be reduced by \Rightarrow monodromies or some symmetry breaking mechanism to:

$$U(1)^n, \text{ or some discrete symmetry } A_4, S_4, \dots$$

... these act as family or discrete symmetries

B

Models


An $SU(5)$ Model (for GUTs see this morning's lecture by G.G. Ross)

$$\mathcal{E}_8 \rightarrow SU(5) \times SU(5)_\perp \rightarrow \mathcal{C} = SU(5)_\perp.$$

Spectral Cover description: $SU(5)_\perp \rightarrow$ described by Cartan roots:

$$t_i = SU(5) - \text{roots} \rightarrow \sum_i t_i = 0$$

Matter resides in 10 and $\bar{5}$ along intersections with other 7-branes (*intersecting branes: detailed description in Andoniadis' lecture*)



$\lambda_{t,b}$ -Yukawas at intersections and gauge symmetry enhancements

-
- ▲▼ Fluxes: ▲▼
 - ▲▼ $SU(5)$ Chirality
 - ▲▼ $SU(5)$ Symmetry Breaking
 - ▲▼ **Splitting of $SU(5)$ -reps**

Two types of fluxes:

▲ M_{10}, M_5 :

associated with flux-restrictions on $U(1)$'s $\in SU(5)_\perp$:

determine the chirality of **complete** $10, 5 \in SU(5)$.

▲ N_Y :

related to Cartan generators of $SU(5)_{GUT}$.

They are taken along $U(1)_Y \in SU(5)_{GUT}$ and **split** $SU(5)$ -reps.

$U(1)_\perp$ –Flux on SM reps $\in \mathbf{10}$'s:

$$\#10 - \#\overline{10} = \begin{cases} n_{(3,2)_{\frac{1}{6}}} - n_{(\overline{3},2)_{-\frac{1}{6}}} & = M_{10} \\ n_{(\overline{3},1)_{-\frac{2}{3}}} - n_{(3,1)_{\frac{2}{3}}} & = M_{10} \\ n_{(1,1)_1} - n_{(1,1)_{-1}} & = M_{10} \end{cases}$$

$U(1)_\perp$ – Flux on SM reps $\in \mathbf{5}$'s:

$$\#5 - \#\overline{5} = \begin{cases} n_{(3,1)_{-\frac{1}{3}}} - n_{(\overline{3},1)_{\frac{1}{3}}} & = M_5 \\ n_{(1,2)_{\frac{1}{2}}} - n_{(1,2)_{-\frac{1}{2}}} & = M_5 \end{cases}$$

(...subject to: $\sum_i M_{10}^i + \sum_j M_5^j = 0$)

$U(1)_Y$ – **Flux**-splitting of **10**'s:

$$n_{(3,2)_{\frac{1}{6}}} - n_{(\bar{3},2)_{-\frac{1}{6}}} = M_{10}$$

$$n_{(\bar{3},1)_{-\frac{2}{3}}} - n_{(3,1)_{\frac{2}{3}}} = M_{10} - N_{Y_{10}}$$

$$n_{(1,1)_1} - n_{(1,1)_{-1}} = M_{10} + N_{Y_{10}}$$

$U(1)_Y$ – **Flux**-splitting of **5**'s:

$$n_{(3,1)_{-\frac{1}{3}}} - n_{(\bar{3},1)_{\frac{1}{3}}} = M_5$$

$$n_{(1,2)_{\frac{1}{2}}} - n_{(1,2)_{-\frac{1}{2}}} = M_5 + N_{Y_5}$$

(... subject to $\sum_i N_{Y_{10}}^i = \sum_j N_{Y_5}^j = 0, \dots$ etc)

▲ **Spectrum** (... in brief) ▼

- **MSSM** spectrum + **natural** doublet-triplet splitting
- **vector-like** fields $f + \bar{f}$ (always present for $G_S \geq SO(10)$)
- **singlets** + KK-modes ...

Two ways to obtain **Fermion Mass Hierarchy** in F-theory

▲▼ All families on the same curve(s) ($\Sigma_{10}, \Sigma_{\bar{5}}$)

non-commutative geometry, ... **Flux** corrections \Rightarrow **Hierarchy**...

▲▼ Families assigned on different matter curves ($\Sigma_{10}^{1,2,3}, \Sigma_{\bar{5}}^{1,2,3}$)

Monodromy \rightarrow Rank one mass matrices at tree level.

Hierarchy organised by $U(1)$'s (Froggatt Nielsen mechanism)

from underlying E_8 via **Singlet** vevs $\langle \theta_{ij} \rangle$

Choice: $\langle \theta_{14} \rangle \cdot \langle \theta_{43} \rangle \neq 0$

▲ Rank one Quark mass matrices (*GKL and GG Ross*) [JHEP02\(2011\)108](#)

$$M_d = \begin{pmatrix} \lambda_{11}^d \theta_{14}^2 \theta_{43}^2 & \lambda_{12}^d \theta_{14} \theta_{43}^2 & \lambda_{13}^d \theta_{14} \theta_{43} \\ \lambda_{21}^d \theta_{14}^2 \theta_{43} & \lambda_{22}^d \theta_{14} \theta_{43} & \lambda_{23}^d \theta_{14} \\ \lambda_{31}^d \theta_{14} \theta_{43} & \lambda_{32}^d \theta_{43} & 1 \times \lambda_{33}^d \end{pmatrix} v_b, \quad (1)$$

$$M_u = \begin{pmatrix} \lambda_{11}^u \theta_{14}^2 \theta_{43}^2 & \lambda_{12}^u \theta_{14}^2 \theta_{43} & \lambda_{13}^u \theta_{14} \theta_{43} \\ \lambda_{21}^u \theta_{14}^2 \theta_{43} & \lambda_{22}^u \theta_{14}^2 & \lambda_{23}^u \theta_{14} \\ \lambda_{31}^u \theta_{14} \theta_{43} & \lambda_{32}^u \theta_{14} & 1 \times \lambda_{33}^u \end{pmatrix} v_u \quad (2)$$

▲ Yukawa strengths λ_{ij} computed from overlapping integrals ... expected of $\mathcal{O}(1)$.

▲ Singlet vevs θ_{ij} fixed by F- and D-flatness.

Particles' Wavefunctions: solving **EoM** \rightarrow Gaussian profile: $\psi \sim f(z_i)e^{-M|z_i|^2}$




Figure 1: Overlapping of three wavefunctions at triple intersection (Yukawa coupling)

Strength of Yukawa coupling \propto integral of overlapping ψ 's at 3-intersection:

$$\lambda_{ij} \propto \int \psi_i(z_1, z_2) \psi_j(z_1, z_2) \psi_H(z_1, z_2) dz_1 \wedge dz_2 \approx 0.3 - 0.5$$

PART – II

Discrete Symmetries and Neutrinos

.

E.G.Floratos, GKL [arXiv:1511.01875](https://arxiv.org/abs/1511.01875)

Neutrino data: parametrization of mixing angles

$$U_\nu = \begin{pmatrix} c_{12}c_{13} & c_{13}s_{12} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - c_{12}s_{13}s_{23}e^{i\delta} & c_{12}c_{23} - s_{12}s_{13}s_{23}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{23}s_{12}s_{13} - c_{12}s_{23}e^{i\delta} & c_{13}c_{23} \end{pmatrix} \quad (3)$$

Experimental data (3σ range) of the angles ($c_{ij} \equiv \cos \theta_{ij}$)

$$\sin^2 \theta_{12} = [0.259 - 0.359]$$

$$\sin^2 \theta_{23} = [0.331 - 0.637]$$

$$\sin^2 \theta_{13} = [0.0169 - 0.0313]$$

$$\delta = 0.77\pi - 1.36\pi$$

(4)

... for theoretical analysis... see yesterday's Lectures by J.W.F. Valle

Working with $M = m_\nu m_\nu^\dagger$ (assuming Hermitian matrix):

A Hermitian can be written in terms a unitary U (...+ Cayley Hamilton theorem)

$$M = i \log(U) = c_1 I + c_2 U + c_3 U^2 \quad (\mathcal{R}_1)$$

Assuming *invariance* under group generator(s) A_i

$$[M, A_i] = 0 \rightarrow [U, A_i] = 0 \quad (\mathcal{R}_2)$$

▲ (\mathcal{R}_1) \rightarrow disentangles mixing from eigenvalues ... $m_{\nu_i} = m_{\nu_i}(c_{0,1,2})$

▲ (\mathcal{R}_2) $\rightarrow M, U, A_i$ **common** system of eigenvectors:

i) search for groups with $3 - d$ irreps A_i and the right eigenvectors $\rightarrow \nu$ -mixing or ...

ii)...try to express $M = \sum_i A_i$.

... a unified method to construct discrete group representations... **required**.

... this is **feasible** for a wide class of **Discrete Groups** $PSL_2(p)$, p prime

▲ **Requirements:** ▲

▲ Relevant only those with $3 - dim.$ representations

▲ GUT and “perpendicular”-group embedded in maximal symmetry E_8 :

$$E_8 \supset \begin{cases} E_6 \times SU(3)_\perp \\ SO(10) \times SU(4)_\perp \\ SU(5) \times SU(5)_\perp \end{cases} \quad (5)$$

→ In the context of F-theory, $PSL_2(p)$ must be subgroups of

$SU(5)_\perp, SU(4)_\perp, SU(3)_\perp$

$\dots \rightarrow p \leq 11$

Definition of $SL_2(p)$ $p \in \mathbb{Z}/p\mathbb{Z}$

$SL_2(p)$: group of 2×2 matrices with **integer** entries

$$\mathfrak{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad ad - bc = 1 \pmod{p}, \quad p \in \mathbb{Z}/p\mathbb{Z}$$

Group generated by two 2×2 matrices (**Artin's** representation):

$$\mathfrak{a} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \mathfrak{b} = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$$

$$\mathfrak{a}^2 = \mathfrak{b}^3 = -\mathcal{I} \equiv - \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

... additional conditions depend on p .

Definition of $PSL_2(p)$, $p \in \mathbb{Z}/p\mathbb{Z}$

Observing that $Z_2 = \{I, -I\}$ is normal subgroup $\in SL_2(p)$...

...Quotient defines the **projective linear group**

$$SL_2(p) / \{I, -I\} = PSL_2(p)$$

AIM: construction of 3-dim. representations of $PSL_2(p)$. (3×3 matrices)

Method: use of **Weil's Metaplectic Representation**

(based on work of *Balian & Itzykson Acad. Dc. Paris 303 (1986).*)

...this method provides the p -dimensional *reducible* representation of $SL_2(p)$...


... p -dim. splits to two lower dimensional **irreducible** representations:

$$p = \frac{p+1}{2} + \frac{p-1}{2}$$

of discrete groups $\in SU(\frac{p+1}{2})$ and $SU(\frac{p-1}{2})$

Cases of Physical Interest: $p = 3, 5, 7$

- $PSL_2(3) \sim A_4$, (and $SL_2(3)$ its double covering)
- $PSL_2(5) \sim A_5$ (*smallest non-abelian simple group*)
- $SL_2(7)$ and its projective $PSL_2(7) \subset SU(3)$ with 168 elements...
...isomorphic to the group preserving the discrete projective geometry of *Fano plane*.



BACKGROUND

Consider the *GF=Galois field* of discrete circle $GF[p]$ and position eigenfunctions $|q\rangle$

$$GF[p] = \{0, 1, 2, \dots, p-1\}, \quad |q\rangle_i = \delta_{ij}, \quad (i, j) = 1, 2, \dots, p-1$$

Define **Translation** and **Momentum** operators :

$$P|q\rangle = |q+1\rangle \quad (6)$$

$$Q|q\rangle = \omega|q\rangle \quad (7)$$

with

$$\omega = e^{2\pi i/p}, \quad P_{kl} = \delta_{k-1,l}, \quad Q_{kl} = \omega^k \delta_{kl}$$

Properties

Commutation Relation

$$QP = \omega PQ$$

Associated to each-other through the **Discrete Fourier Transform (DFT)** $F_{kl} = \frac{1}{\sqrt{p}}\omega^{kl}$:

$$P = F^{-1}QF$$

Heisenberg Group \mathcal{H}

P and Q generate \mathcal{H} with elements of the form:

$$J_{n_1, n_2, t} = \omega^t P^{n_1} Q^{n_2},$$

with $t \in \mathbb{Z}/p\mathbb{Z}$, $n_1, n_2 \in (\mathbb{Z}/p\mathbb{Z})^2$. Isomorphic to the group of matrices:

$$J_{n_1, n_2, t} \leftrightarrow \begin{pmatrix} 1 & 0 & 0 \\ n_1 & 1 & 0 \\ t & n_2 & 1 \end{pmatrix}$$

Working with a subset of it ($t \rightarrow \frac{n_1 n_2}{2}$):

$$J_{\vec{n}} \equiv J_{n_1, n_2} = \omega^{\frac{n_1 n_2}{2}} P^{n_1} Q^{n_2}, \quad \vec{n} = (n_1, n_2)$$

which obeys the 'multiplication' law

$$J_{\vec{m}} J_{\vec{n}} = \omega^{\frac{\vec{n} \times \vec{m}}{2}} J_{\vec{m} + \vec{n}}$$

Metaplectic Representation

... the action of an $SL_2(p)$ element $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ on coordinates (r, s) of periodic lattice $\mathbb{Z}_p \times \mathbb{Z}_p$ induces unitary automorphism $U(A)$:

$$U(A)J_{r,s}U^\dagger(A) = J_{r',s'}, \quad \text{where } (r',s') = (r,s) \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

Formula of $U(A)$ has been given by *Balian and Itzykson (1986)*:

$$U(A) = \frac{\sigma(1)\sigma(\delta)}{p} \sum_{r,s} \omega^{[br^2 + (d-a)rs - cs^2]/(2\delta)} J_{r,s}$$

for $\delta = 2 - a - d \neq 0$, and:

$$\delta = 0, \quad b \neq 0 : \quad U(A) = \frac{\sigma(-2b)}{\sqrt{p}} \sum_s \omega^{s^2/(2b)} J_{s(a-1)/b,s}$$

$$\delta = b = 0, \quad c \neq 0 : \quad U(A) = \frac{\sigma(2c)}{\sqrt{p}} \sum_r \omega^{-r^2/(2c)} P^r$$

$$\delta = b = 0 = c = 0 : \quad U(1) = I$$

(8)

A few clarifications on notation

$\sigma(a)$ is the **Quadratic Gauss Sum**,

$$\sigma(a) = \frac{1}{\sqrt{p}} \sum_{k=0}^{p-1} \omega^{ak^2} = (a|p) \times \begin{cases} 1 & \text{for } p = 4k + 1 \\ i & \text{for } p = 4k - 1 \end{cases}$$

and $(a|p)$ the **Legendre symbol**

$$\left(\frac{a}{p}\right) = \begin{cases} 0 & \text{if } a \text{ divides } p \\ +1 & \text{if } a = \mathcal{QR} \ p \\ -1 & \text{if } a \neq \mathcal{QR} \ p \end{cases}$$

(9)

(integer q is \mathcal{QR} \rightarrow Quadratic Residue **iff** $\exists x : x^2 = q \pmod{p}$.)

for $x = 0, 1, 2, 3, \dots$, $x^2 \pmod{5} = 0, 1, 4, 4, 1, 0, 1, 4, 4, 1, \dots$

The construction of the $SL_2(p)$ representations

... it suffices to construct only the two generators $\mathfrak{a} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $\mathfrak{b} = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$. Observe that

$$U(\mathfrak{a}) = (-1)^{k+1} i^n F, \quad \begin{cases} n = 0 & \text{for } p = 2k + 1 \\ n = 1 & \text{for } p = 2k - 1 \end{cases}$$

Observe also that **DFT** generates an *Abelian group* with four elements

$$F, \quad S = F^2, \quad F^3 = F^*, \quad S^2 \equiv F^4 = I$$

and... since $S^2 = I \Rightarrow S$: can be used to define projection operators

$$P_{\pm} = \frac{1 \pm S}{2}$$

... *split* $SL_2(p)$ *reducible* representations to $\frac{p+1}{2}$ & $\frac{p-1}{2}$ dim. **irreps**:

$$U(A)_{\pm} = U(A)P_{\pm}$$

... $U(A)_\pm$ block diagonal form achieved by orthogonal matrix \mathcal{O} of S eigenvectors:

$$(e_0)_k = \delta_{k0},$$

$$(e_j^+)_k = \frac{1}{\sqrt{2}}(\delta_{k,j} + \delta_{k,-j}), \quad j = 1, \dots, \frac{p-1}{2}$$

$$(e_j^-)_k = \frac{1}{\sqrt{2}}(\delta_{k,j} - \delta_{k,-j}), \quad j = \frac{p+1}{2}, \dots, p$$

Example. $SL_2(7)$ case:

$$\mathcal{O} = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}$$

Final block-diagonal form:

$$V_\pm(A) = \mathcal{O}U(A)_\pm\mathcal{O}$$

▲ $SL_2(7)$ has $(p^2 - 1)p = 336$ elements

▲ $PSL_2(7)$ has 168 elements ($= 7 \times 24$ hours in week!)

Construction of 3-d. irreducible representation of $PSL_2(7)$ satisfying:

$$\mathbf{a}^2 = \mathbf{b}^3 = (\mathbf{ab})^7 = ([\mathbf{a}, \mathbf{b}])^4 = I$$

from 7-d. reducible rep. of $SL_2(7)$.

Defining $\eta = e^{2\pi i/7}$, (7th root of unity)

$$\mathbf{a} \rightarrow A^{[3]} = \frac{i}{\sqrt{7}} \begin{pmatrix} \eta^2 - \eta^5 & \eta^6 - \eta & \eta^3 - \eta^4 \\ \eta^6 - \eta & \eta^4 - \eta^3 & \eta^2 - \eta^5 \\ \eta^3 - \eta^4 & \eta^2 - \eta^5 & \eta - \eta^6 \end{pmatrix}$$

and

$$\mathbf{b} \rightarrow B^{[3]} = \frac{i}{\sqrt{7}} \begin{pmatrix} \eta - \eta^4 & \eta^4 - \eta^6 & \eta^6 - 1 \\ \eta^5 - 1 & \eta^2 - \eta & \eta^5 - \eta \\ \eta^2 - \eta^3 & 1 - \eta^3 & \eta^4 - \eta^2 \end{pmatrix}$$

APPLICATION

... assuming invariance of neutrino mass matrix

$$[m_\nu, A^{[3]}] = 0 \rightarrow m_\nu = \begin{pmatrix} z_5 + x & z_4 - y & z_4 \\ z_4 - y & z_5 - y & z_6 \\ z_4 & z_6 & z_5 \end{pmatrix}, \begin{cases} x = \alpha (z_4 + z_6) \\ \alpha = 32 \sin^3\left(\frac{\pi}{14}\right) \sin^2\left(\frac{3\pi}{14}\right) \\ y = \beta (z_4 + z_6) \\ \alpha + \beta = 1 \end{cases}$$

The neutrino mixing matrix is found analytically.

$$\mathbf{U}_{PMNS} = V_\ell^T V_\nu = \begin{pmatrix} -0.814857 & 0.57735 & 0.051711 \\ 0.452212 & 0.57735 & 0.679832 \\ -0.362646 & -0.57735 & 0.731543 \end{pmatrix}$$

Comparison with experimental data:

- ▲ θ_{12}, θ_{23} in perfect agreement with experimental values.
- ▲ θ_{13} automatically non-zero, although smaller than experimental value.

A systematic study required ...

... work in progress with *N.D. Vlachos*

*C*oncluding *R*emarks

F-theory models :



Geometric interpretation of GUTs

Calculability, form handful of topological properties, natural Doublet-Triplet splitting...

Prediction of Vector-like pairs and singlets ...

hints for New Physics

such as ... resonances, diphoton events at a few TeV...

Discrete Symmetries interpreting the Neutrino data naturally incorporated in E_8 singularity

The analytic form of V_ν

The analytic form of the neutrino mixing matrix

$$V_\nu = \begin{pmatrix} \frac{1 + \frac{2 \cos\left(\frac{\pi}{14}\right)}{\sqrt{7}} - \frac{2 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}}{\sqrt{6 \left(1 - \frac{2 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}\right)}} & -\frac{1}{\sqrt{3}} \frac{1 - \frac{2 \cos\left(\frac{\pi}{14}\right)}{\sqrt{7}} + \frac{2 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}}{\sqrt{2 \left(1 + \frac{6 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}\right)}} \\ \frac{1 - \frac{2 \sin\left(\frac{\pi}{7}\right)}{\sqrt{7}} - \frac{2 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}}{\sqrt{6 \left(1 - \frac{2 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}\right)}} & \frac{1}{\sqrt{3}} \frac{1 + \frac{2 \sin\left(\frac{\pi}{7}\right)}{\sqrt{7}} + \frac{2 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}}{\sqrt{2 \left(1 + \frac{6 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}\right)}} \\ \frac{2 \sin\left(\frac{\pi}{7}\right)}{\sqrt{7}} + \frac{2 \cos\left(\frac{\pi}{14}\right)}{\sqrt{7}}}{\sqrt{6 \left(1 - \frac{2 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}\right)}} & \frac{1}{\sqrt{3}} \frac{-\frac{2 \sin\left(\frac{\pi}{7}\right)}{\sqrt{7}} - \frac{2 \cos\left(\frac{\pi}{14}\right)}{\sqrt{7}}}{\sqrt{2 \left(1 + \frac{6 \cos\left(\frac{3\pi}{14}\right)}{\sqrt{7}}\right)}} \end{pmatrix}$$