

Noncommutative Spherically Symmetric Spacetimes

Christopher Fritz¹

in collaboration with Shahn Majid²

¹Sussex University ²QMUL

September 21, 2017

Published in Class. Quantum Grav.
DOI: 10.1088/1361-6382/aa72a5
arXiv:1611.04971

Outline

- ▶ Algebra → Differential Algebra
- ▶ Differential Algebra → Metric, Connection and Geometry
- ▶ Compatibility with Semiclassical Data
- ▶ Uniqueness Theorem for Spherically Symmetric Spacetime

Differential Algebra

Algebra

- ▶ (\mathcal{A}, \cdot) unital, associative etc.
- ▶ Not necessarily commutative: $ab \neq ba$

Differential Algebra

- ▶ $\Omega^1 \equiv \Omega^1(\mathcal{A})$ is a bimodule of \mathcal{A} so $a((db)c) = (a(db))c$
- ▶ $d : \mathcal{A} \rightarrow \Omega^1$ so $d(ab) = (da)b + a(db)$
- ▶ $\{adb\}$ span Ω^1
- ▶ $\ker d = k$

Extends to Differential Graded Algebra $\Omega = \bigoplus_i \Omega^i$

- ▶ $d^2 = 0$
- ▶ Product $\wedge : \Omega^n(\mathcal{A}) \otimes \Omega^m(\mathcal{A}) \rightarrow \Omega^{n+m}$
- ▶ $d(\alpha \wedge \beta) = d(\alpha) \wedge \beta + (-1)^{|\alpha|} \alpha \wedge d(\beta)$

Quantum metric

$g \in \Omega^1 \otimes \Omega^1$ with quantum symmetry $\wedge(g) = 0$

Require

- ▶ Inverse $(\cdot, \cdot) : \Omega^1 \otimes \Omega^1 \rightarrow \mathcal{A}$
- ▶ Bimodule

Forces g to be central: $[g, f] = 0 \quad \forall f \in \mathcal{A}$

Also want a compatible bimodule connection

$$\nabla : \Omega^1 \rightarrow \Omega^1 \otimes \Omega^1, \quad \sigma : \Omega^1 \otimes \Omega^1 \rightarrow \Omega^1 \otimes \Omega^1$$

- ▶ Metric compatible $\nabla(g) = 0$
- ▶ Torsion free $\wedge\nabla = d$

Quantum Levi-Civita connection!

Example

S. Majid, W. Tao; Phys. Rev. D 91 (2015)

Take Majid-Ruegg model: $[x^i, t] = i\lambda_P x^i$ & $[t, dt] = i\lambda_P \alpha dt$

$$g = a_{ij}dx^i \otimes dx^j + b_i(dx^i \otimes dt + dt \otimes dx^i) + cdt \otimes dt$$

$$[g, f] = 0 \quad \forall f \in \mathcal{A}, \quad \wedge(g) = 0$$



$$g = \delta^{-1}d\Omega^2 + ar^{-2}dr \otimes dr + br^{\alpha-1}(dr \otimes dt + dt \otimes dr) + cr^{2\alpha}dt \otimes dt$$

$$\delta > 0 \quad a, b, c, \delta \in \mathbf{R}$$

This is the Bertotti-Robinson metric.

Algebra forces the metric!

Semi-Classical Quantum Gravity

E. Beggs, S.Majid; Class.Quant.Grav. 31 (2014)

Interested in quantization to $\mathcal{O}(\lambda) \rightarrow$ Semiquantization

- ▶ Controlled by Poisson bracket $(\{ , \}, C^\infty(\mathcal{M}))$
- ▶ Have $[a, b] = \lambda \{a, b\} + \mathcal{O}(\lambda^2)$
- ▶ $\{ , \} \leftrightarrow \omega^{ij}$

For differential structure have

- ▶ For $\eta \in \Omega^1$ have $[a, \eta] = \lambda \nabla_{\hat{a}} \eta + \mathcal{O}(\lambda^2)$
- ▶ $\nabla_{\hat{a}}$ is a Poisson (pre)connection defined along a Hamiltonian vector field $\hat{a} = \{a, \}$
- ▶ $d\{a, b\} = \nabla_{\hat{a}} b - \nabla_{\hat{b}} a$

Look at associativity \longrightarrow Jacobi identity

$$[a, [db, c]] + [c, [a, db]] + [db, [c, a]] \sim \nabla_{\hat{a}} \nabla_{\hat{b}} db - \nabla_{\hat{a}} \nabla_{\hat{c}} db - \nabla_{\widehat{\{a,c\}}} db$$

Nonflat connection \longrightarrow nonassociative calculus at $\mathcal{O}(\lambda^2)$ (but associative functions)

Semiquantization

E. Beggs, S.Majid; J.Geom.Phys. 114 (2017)

- ▶ Want quantum Levi-Civita connection and central metric
- ▶ Semiclassical data: ω^{ij} and $\nabla_{\hat{a}}$
- ▶ Compatibility?

$Q:$ (vector bundles and connections) \longrightarrow (bimodules over a deformed algebra)

From the standpoint of physics can view λ as the effective scale of the theory (e.g. Planck scale), so it is reasonable to work only to first order.

Semiquantization

What do we need? Have

- ▶ Metric g
- ▶ Poisson tensor $\omega^{ij} \Leftrightarrow \sum_{\text{cyclic}(i,j,k)} \omega^{il}\omega^{jk},_l = 0$
- ▶ Poisson connection ∇
- ▶ Levi-Civita connection: $\widehat{\nabla} = \nabla + S$
$$S^\alpha{}_{\mu\nu} = \frac{1}{2}g^{\alpha\beta} (T_{\beta\mu\nu} + T_{\mu\beta\nu} + T_{\nu\beta\mu})$$

And want

- ▶ Metric: $g \rightarrow g_1$
- ▶ Connection: $\widehat{\nabla} \rightarrow \nabla_1$
- ▶ Tensor product: $E \otimes F \rightarrow Q(E) \otimes_1 Q(F)$
- ▶ Wedge product: $\eta \wedge \xi \rightarrow Q(\eta) \wedge_1 Q(\xi)$

Semiquantization

Generalized Ricci two form

$$\mathcal{R} = g_{\alpha\beta}\omega^{\alpha\gamma}(\nabla_\gamma S^\beta{}_{\mu\nu} - R^\beta{}_{\mu\nu\gamma})dx^\nu \wedge dx^\mu$$

Then for $\hat{\nabla} = \nabla + S$ have the conditions

- ▶ Poisson Compatibility $\hat{\nabla}_\gamma \omega^{\alpha\beta} + S^\alpha{}_{\delta\gamma} \omega^{\delta\beta} + S^\beta{}_{\delta\gamma} \omega^{\alpha\delta} = 0$
- ▶ Metric compatibility $\nabla g = 0$
- ▶ Quantum Levi-Civita condition

$$\hat{\nabla}\mathcal{R} + \omega^{\alpha\beta} g_{\rho\sigma} S^\sigma{}_{\beta\nu} (R^\rho{}_{\mu\gamma\alpha} + \nabla_\alpha S^\rho{}_{\gamma\mu}) dx^\gamma \otimes dx^\mu \wedge dx^\nu = 0$$

Application

Take generic spherically symmetric metric

$$g = a^2(r, t)dt \otimes dt + b^2(r, t)dr \otimes dr + c^2(r, t)(d\theta \otimes d\theta + \sin^2(\theta)d\phi \otimes d\phi)$$

Poisson tensor? Ansatz: spherical symmetry

$$\omega^{23} = \frac{f(r, t)}{\sin(\theta)} \quad \omega^{01} = h(r, t)$$

So have Levi-Civita connection $\hat{\nabla}$ and need

- ▶ ω^{ij} Poisson: $h\partial_t f = h\partial_r f = 0$
- ▶ Poisson connection $\nabla = \hat{\nabla} - S$
- ▶ Quantum L.C. condition

Uniqueness Theorem

Find $f = k$ and $g = 0$ and

$$S_{022} = c\partial_t c, S_{122} = c\partial_r c, S_{033} = c\partial_t c \sin^2(\theta), S_{133} = c\partial_r c \sin^2(\theta)$$

$$S_{120} = S_{123} = S_{223} = S_{320} = S_{130} = S_{132} = S_{230} = S_{233} = 0$$

- ▶ Unique quantization up to $\mathcal{O}(\lambda^2)$ with algebra

$$[z^i, z^j] = \lambda \epsilon^{ij}_k z^k, \quad [z^i, dz^j] = \lambda z^i \epsilon^j_{mn} z^m dz^n$$

$$\sum (z^i)^2 = 1, \quad \sum z^i dz^i = 0$$

- ▶ Nonassociative fuzzy sphere.
- ▶ Central (classical) t, r, dt, dr

Semiquantization

Quantum metric

$$g_1 = g_{\mu\nu} dx^\mu \otimes_1 dx^\nu + \frac{\lambda}{2} \omega^{\alpha\beta} \Gamma_{\mu\alpha\kappa} \Gamma^\kappa{}_{\beta\nu} dx^\mu \otimes_1 dx^\nu + \frac{\lambda}{2} \mathcal{R}_{\mu\nu} dx^\mu \otimes_1 dx^\nu$$

Quantum Connection

$$\begin{aligned} \nabla_1 dx^\iota = - & \left[\widehat{\Gamma}^\iota{}_{\mu\nu} + \frac{\lambda}{2} \omega^{\alpha\beta} \left(\widehat{\Gamma}^\iota{}_{\mu\kappa,\alpha} \Gamma^\kappa{}_{\beta\nu} - \widehat{\Gamma}^\iota{}_{\kappa\tau} \Gamma^\kappa{}_{\alpha\mu} \Gamma^\tau{}_{\beta\nu} \right. \right. \\ & \left. \left. + \widehat{\Gamma}^\iota{}_{\alpha\kappa} (R^\kappa{}_{\nu\mu\beta} + \nabla_\beta S^\kappa{}_{\mu\nu}) \right) \right] dx^\mu \otimes_1 dx^\nu \end{aligned}$$

Quantum Laplace operator

$$\square_1 f := (\ , \)_1 \nabla_1 df = \square f + \frac{\lambda}{2} \omega^{\alpha\beta} (\text{Ric}^\gamma{}_\alpha - S^\gamma{}_{;\alpha}) (\widehat{\nabla}_\beta df)_\gamma$$

Metric and Connection

Quantum Metric

$$g_1 = g_{\mu\nu} dx^\mu \otimes_1 dx^\nu + \frac{\lambda c^2}{2(z^3)^2} \epsilon_{3ij} (z^3 dz^i \otimes_1 dz^j - z^i dz^3 \otimes_1 dz^j)$$

Quantum Connection

$$\nabla_1(dt) = -\widehat{\Gamma}_\mu{}^\nu dx^\mu \otimes_1 dx^\nu - \frac{\lambda}{2(z^3)^2} \frac{c \partial_t c}{a^2} \epsilon_{3ij} (z^3 dz^i \otimes_1 dz^j - z^i dz^3 \otimes_1 dz^j)$$

$$\nabla_1(dr) = -\widehat{\Gamma}_\mu{}^\nu dx^\mu \otimes_1 dx^\nu + \frac{\lambda}{2(z^3)^2} \frac{c \partial_r c}{b^2} \epsilon_{3ij} (z^3 dz^i \otimes_1 dz^j - z^i dz^3 \otimes_1 dz^j)$$

$$\nabla_1(dz^a) = -\widehat{\Gamma}_\mu{}^\nu dx^\mu \otimes_1 dx^\nu + \frac{\lambda}{2} \left(\epsilon_{ijk} z^k z^a dz^i \otimes_1 dz^j - \frac{1}{(z^3)^2} \epsilon^a{}_{i3} dz^3 \otimes_1 dz^i \right)$$

Quantum Laplace operator

$$\square_1 f = g^{\alpha\beta} \left(f_{,\alpha\beta} + f_{,\gamma} \widehat{\Gamma}^\gamma{}_{\alpha\beta} \right)$$