

Workshop on the standard model and beyond
Corfu Summer Institute
27 August - 7 September 2023

A grandunified RHINO miracle
in IceCube

Pasquale Di Bari
(University of Southampton)

Why new physics?

Even ignoring:

- (more or less) compelling theoretical motivations (quantum gravity theory, flavour problem, hierarchy and naturalness problems,...) and
- Experimental anomalies (e.g., $(g-2)_\mu$, R_K , R_K^* , ...)

The SM+GR cannot explain:

- Cosmological Puzzles :

1. Dark matter
2. Matter - antimatter asymmetry
3. Inflation
4. Accelerating Universe

- Neutrino masses and mixing


problem of the origin of matter in the universe

Dark Matter


At the present time dark matter acts as a cosmic glue keeping together

stars in galaxies and

galaxies in clusters of galaxies




Coma cluster




bullet cluster

...but it also needs to be primordial to understand structure formation and CMB anisotropies



(Hu, Dodelson, astro-ph/0110414)




CMB +
BAO

(Planck 2018, 1807.06209)

$$\Omega_{CDM,0} h^2 = 0.11933 \pm 0.0009 \sim 5 \Omega_{B,0} h^2$$

WIMP miracle




$$\Omega_X h^2 \simeq \frac{4 \times 10^{-10}}{\langle \sigma_{\text{ann}} \beta_{\text{rel}} \rangle} \left(\frac{hc}{\text{GeV}} \right)^2 \simeq \frac{1.6 \times 10^{-37} \text{ cm}^2}{\langle \sigma_{\text{ann}} \beta_{\text{rel}} \rangle t},$$

$$\langle \sigma_{\text{ann}}^{\text{weak}} \beta_{\text{rel}} \rangle \simeq 0.1 \frac{\alpha_{\text{weak}}^2}{m_X^2} \simeq 4 \times 10^{-36} \text{ cm}^2 \left(\frac{100 \text{ GeV}}{m_X} \right)^2 \Rightarrow \Omega_{\text{DM}} h^2 \sim \mathcal{O}(0.1) \left(\frac{m_X}{100 \text{ GeV}} \right)^2$$

- ❑ embeddable in models addressing naturalness+hierarchy problems
- ❑ \Rightarrow **new physics at the 100 GeV - TeV scale**
- ❑ The **WIMP miracle** has been for long time regarded as a strong argument in favour of WIMPs as dark matter particles.
- ❑ The lack of evidence of new physics at the TeV scale makes the WIMP miracle, if not completely ruled out, certainly less compelling.

Beyond the WIMP paradigm: the DM particle zoo

(from Baer et al.1407.0017)



(PDB, Anisimov '08)

From an old to a new miracle?

?



EW




Searching for a new miracle



Examples of DM beyond the standard WIMPs:

- Dark matter could decay after freeze-out
example: gravitino dark matter with R parity breaking
(Buchmuller, Covi, Hamaguchi Ibarra, Yanagida hep--ph/0702184)
- Freeze-in solution (FIMPs)



$$\Omega_{DM0} h^2 \propto \langle \sigma_{\text{ann}} \beta_{\text{rel}} \rangle$$

- Or both: freeze-in + decaying DM!
(example: keV seesaw neutrino solution)

Minimal seesaw mechanism (type I)

• Dirac + (right-right) Majorana mass term

(Minkowski '77; Gell-mann, Ramond, Slansky; Yanagida; Mohapatra, Senjanovic '79)

$$-\mathcal{L}_{mass}^{\nu} = \bar{\nu}_L m_D \nu_R + \frac{1}{2} \bar{\nu}_R^c M \nu_R + h.c. = -\frac{1}{2} (\bar{\nu}_L \quad \bar{\nu}_R^c) \begin{pmatrix} 0 & m_D^T \\ m_D & M \end{pmatrix} \begin{pmatrix} \nu_L \\ \nu_R^c \end{pmatrix} + h.c.$$

In the *see-saw limit* ($M \gg m_D$) the mass spectrum splits into 2 sets:


- 3 light **Majorana neutrinos** with masses (seesaw formula): $m_{\nu} = -m_D M^{-1} m_D^T \Rightarrow \text{diag}(m_1, m_2, m_3) = -U^\dagger m_{\nu} U^*$
- 3(?) very heavy Majorana neutrinos N_1, N_2, N_3 with $M_3 > M_2 > M_1 \gg m_D$

1 generation toy model :

$$m_D \sim m_{\text{top}},$$

$$m \sim m_{\text{atm}} \sim 50 \text{ meV}$$

$$\Rightarrow M \sim 10^{15} \text{ GeV}$$



Dark matter from active-sterile neutrino mixing

(Dodelson Widrow '94; Shi, Fuller '99; Dolgov and Hansen '00; Asaka, Blanchet, Shaposhnikov

- LH-RH
(active-sterile)
neutrino mixing

$$\nu_{1L} \simeq U_{1\alpha}^\dagger \left(\nu_{L\alpha} - \frac{m_{D\alpha 1}}{M_1} \nu_{R1}^c \right)$$

$$N_{1R} \simeq \nu_{1R} + \frac{m_{D\alpha 1}}{M_1} \nu_{L\alpha}^c \longrightarrow \text{lightest RH neutrino}$$

- Solving Boltzmann equations an abundance is produced at $T \sim 100$ MeV:

$$\Omega_{N_1} h^2 \sim 0.1 \frac{\theta^2}{10^{-8}} \left(\frac{M_1}{\text{keV}} \right)^2 \sim \Omega_{DM,0} h^2 \quad \theta^2 \equiv \frac{\sum_\alpha |m_{D\alpha 1}|^2}{M_1^2}$$

- For $M_1 \ll m_e \Rightarrow \tau_1 = 5 \times 10^{26} \text{ s} \left(\frac{M_1}{\text{keV}} \right)^{-5} \left(\frac{10^{-8}}{\theta^2} \right) \gg t_0$

- The lightest neutrino mass $m_1 \lesssim 10^{-5} \text{ eV} \Rightarrow$ **hierarchical neutrino masses**
- The N_1 's also radiatively decay and this produces constraints from X-rays (or opportunities to observe it).
- Considering also structure formation constraints, one is forced to consider a resonant production induced by a large lepton asymmetry
- $L \sim 10^{-4}$: 3.5 keV line? (Horiuchi et al. '14; Bulbul et al. '14; Abazajian '14)

The XRISM satellite should give a final answer

Dark matter from active-sterile neutrino mixing

(Dodelson Widrow '94; Shi, Fuller '99; Dolgov and Hansen '00; Asaka, Blanchet, Shaposhnikov

$$\nu_{1L} \simeq U_{1\alpha}^\dagger \left(\nu_{L\alpha} - \frac{m_{D\alpha 1}}{M_1} \nu_{R1}^c \right)$$

- LH-RH

X-Ray Imaging and Spectroscopy Mission (XRISM)

Announcement

The XRISM launch scheduled for Sunday, Aug. 27 (Monday, Aug. 28, in Japan) has been **postponed**. The new launch date and time will be announced once confirmed. [updated on Aug 27, 2023]

Launch [edit]

JAXA planned to launch XRISM on August 27 using [H-IIA](#) rocket from [Tanegashima Space Center](#), but scrapped because of bad weather. Launch window is reserved till September 15. ^[25]

- $L \sim 10^{-4}$: 3.5 keV line? (Horiuchi et al. '14; Bulbul et al. '14; Abazajian '14)

The XRISM satellite should give a final answer

Heavy RH neutrino as dark matter ?

(Anisimov, PDB '08)

What production mechanism? For high masses just a tiny abundance is needed:

$$N_{DM} \simeq 10^{-9} (\Omega_{DM,0} h^2) N_\gamma(t_{prod}) \frac{\text{TeV}}{M_{DM}}$$

Suppose there is a RH neutrino with tiny Yukawa couplings (e.g., proportional to a small symmetry breaking parameter) referred to as **dark neutrino** N_D :

$$m_D \simeq \begin{pmatrix} \varepsilon_{e1} & m_{De2} & m_{De3} \\ \varepsilon_{\mu1} & m_{D\mu2} & m_{D\mu3} \\ \varepsilon_{\tau1} & m_{D\tau2} & m_{D\tau3} \end{pmatrix} \text{ or } m_D \simeq \begin{pmatrix} m_{De1} & \varepsilon_{e2} & m_{De3} \\ m_{D\mu1} & \varepsilon_{\mu2} & m_{D\mu3} \\ m_{D\tau1} & \varepsilon_{\tau2} & m_{D\tau3} \end{pmatrix} \text{ or } m_D \simeq \begin{pmatrix} m_{De1} & m_{De2} & \varepsilon_{e3} \\ m_{D\mu1} & m_{D\mu2} & \varepsilon_{\mu3} \\ m_{D\tau1} & m_{D\tau2} & \varepsilon_{\tau3} \end{pmatrix}$$

$$m_D = V_L^\dagger D_{m_D} U_R \quad D_{m_D} \equiv v \text{diag}(h_A, h_B, h_C) \text{ with } h_A \leq h_B \leq h_C$$

$$\tau_{DM} = \frac{4\pi}{h_A^2 M_{DM}} = 0.87 h_A^2 10^{-26} \frac{\text{TeV}}{M_{DM}} s$$

\Rightarrow

$$\tau_{DM} > \tau_{DM}^{\min} \simeq 10^{28} s \Rightarrow h_A < 10^{-27} \sqrt{\frac{\text{TeV}}{M_{DM}} \times \frac{10^{28} s}{\tau_{DM}^{\min}}}$$

Too small to reproduce the correct abundance with any production mechanism within a minimal type-I seesaw extension

5-dimensional Higgs portal-like operators as a way out

(Anisimov hep-ph/0612024, Anisimov,PDB 0812.5085)

$$\mathcal{L} = \mathcal{L}_{SM} + \mathcal{L}_{Y+M}^{\nu} + \mathcal{L}_A$$

Type-I
Seesaw
Lagrangian

$$-\mathcal{L}_{Y+M}^{\nu} = \bar{L}_{\alpha} h_{\alpha I} N_I \tilde{\phi} + \frac{1}{2} \overline{N_I^c} M_I N_I + h.c.$$

Anisimov
operator(s)

$$\mathcal{L}_A = \sum_{I,J} \frac{\lambda_{IJ}}{\Lambda} \phi^{\dagger} \phi \overline{N_I^c} N_J + h.c.$$

$$= \frac{\lambda_{DS}}{\Lambda} \phi^{\dagger} \phi \overline{N_D^c} N_S + \frac{\lambda_{SS}}{\Lambda} \phi^{\dagger} \phi \overline{N_S^c} N_S + \frac{\lambda_{DD}}{\Lambda} \phi^{\dagger} \phi \overline{N_D^c} N_D + h.c. \quad (N_D = N_3; N_S = N_2)$$

Remarks: \longrightarrow RH-RH (sterile-sterile) Higgs-induced neutrino mixing (RHINO)

- from SMEFT to vSMEFT
- They are Weinberg-like operators but a further step up
- They extend Higgs portal renormalizable operator (Patt, Wilczek hep-ph/0605188)

RHINO dark matter


(Anisimov '06, Anisimov, PDB '08)

Focus on the RH-RH Higgs-induced neutrino mixing (RHINO) operator:

$$\mathcal{L}_A = \frac{\lambda_{DS}}{\Lambda} \phi^\dagger \phi \overline{N_D^c} N_S \quad \tilde{\Lambda}_{DS} = \Lambda / \lambda_{DS}$$


In general, $\lambda_{DS} \neq 0$ generates a dark-source RH neutrino mixing. The Yukawa and Anisimov interactions both generate effective potentials from self-energies:

From Yukawa interactions



$$\Rightarrow V_S^Y = \frac{T^2}{8p} h_S^2$$

From mixing



$$\Rightarrow V_{DS}^\Lambda = \frac{T^2}{12\Lambda} \lambda_{DS}$$

Effective mixing Hamiltonian :

$$\Delta H \approx \begin{pmatrix} -\frac{\Delta M^2}{4p} - \frac{T^2}{16p} h_S^2 & \frac{T^2}{12\tilde{\Lambda}_{DS}} \\ \frac{T^2}{12\tilde{\Lambda}_{DS}} & \frac{\Delta M^2}{4p} + \frac{T^2}{16p} h_S^2 \end{pmatrix}$$

mixing term

$$\Delta M^2 \equiv M_S^2 - M_D^2$$


Density matrix calculation of the relic abundance

(P.Di Bari, K. Farrag, R. Samanta, Y. Zhou, 1908.00521)

Density matrix equation for the dark-source RH neutrino system (using a monochromatic approximation $p \sim 3T$)

$$\frac{dN_{IJ}}{dt} = -i[\Delta H, N]_{IJ} - \begin{pmatrix} 0 & \frac{1}{2}(\Gamma_D + \Gamma_S)N_{DS} \\ \frac{1}{2}(\Gamma_D + \Gamma_S)N_{SD} & (\Gamma_D + \Gamma_S)(N_{N_S} - N_{N_S}^{eq}) \end{pmatrix}$$

Assuming an **initial thermal N_S -abundance**




Dark neutrinos are necessarily unstable

(Anisimov,PDB '08; Anisimov,PDB'10; P.Ludl.PDB,S.Palomarez-Ruiz'16)

2 body decays ($M_S > M_W$)

Dark neutrinos unavoidably decay today into $A + \text{leptons}$ ($A = H, Z, W$) through the same mixing that produced them in the very early Universe




$$\theta_{\Lambda 0} = \frac{2 v^2 / \tilde{\Lambda}_{DS}}{M_D (1 - M_S / M_D)}$$

mixing angle today
(for $\theta_{\Lambda 0} \ll 1$)

$$\Gamma_{D \rightarrow A + l_S} = \frac{h_S^2}{\pi} \left(\frac{v^2}{\tilde{\Lambda}} \right)^2 \frac{M_D}{(M_D - M_S)^2}$$

\Rightarrow Lower bound on M_D

4 body decays



$$N_{DM} \rightarrow 2A + N_S \rightarrow 3A + l_S \quad (A = W^\pm, Z, H).$$


$$\Gamma_{D \rightarrow 3A + l_S} = \frac{\Gamma_S}{15 \cdot 2^{11} \cdot \pi^4} \frac{M_D}{M_S} \left(\frac{M_D}{\tilde{\Lambda}_{DS}} \right)^2$$

\Rightarrow Upper bound on M_D

3 body decays and annihilations can also occur but yield weaker constraints

DM lifetime vs. mass plane: allowed regions

(P. Di Bari, K. Farrag, R. Samanta, Y. Zhou, 1908.00521)



It works only for **initial thermal N_S abundance**, unless $M_S \sim 1$ GeV and $M_D \gtrsim 10^7$ GeV


Can one think of processes able to thermalize the N_S abundance prior to the oscillations?

Two good motivations

Unifying Leptogenesis and Dark Matter

(PDB, K. Farrag, R. Samanta, Y. Zhou, 1908.00521)

A solution for initial thermal N_S abundance:



Lower bound on the lifetime of decaying DM





FIG. 2: *Constraints on the lifetime of dark matter decaying to neutrinos $\chi \rightarrow \bar{\nu}\nu$.* Solid lines bordering shaded regions represent limits from existing neutrino telescope data, solid lines without shading correspond to limits from existing gamma-ray observatories (as shown in Fig. 3), and dashed lines show the reach of future experiments. Labels with a heart symbol (♡) correspond to limits derived for this work.

(Argüelles et al., 2210.01303)

$$\chi \rightarrow \nu \bar{\nu}$$


IceCube



- Neutrinos are perfect astronomical messengers (from the edge of the universe)
- In the range 10 TeV - 10 EeV only neutrinos are unabsorbed and undeflected
- 2013: IceCube discovered cosmic VHE neutrinos (30 TeV - 1 PeV range)
- Some observed in coincidence with blazar γ -ray flare: **extragalactic origin**
- **High Energy Starting Events (HESE)** veto to reduce overwhelming atmospheric background at energies $\lesssim 300$ TeV \Rightarrow first evidence of cosmic neutrinos
- **Up-going muon data set** has confirmed the existence of cosmic neutrinos but ...

IceCube up-going muon neutrinos

IceCube 8 years data



Standard single power-law spectrum for an astrophysical flux


$$\frac{d\Phi}{dE} = \Phi_0 \cdot \left(\frac{E_\nu}{100\text{TeV}} \right)^{-\gamma_{astro}}$$

Best fit


$$\frac{d\Phi_{\nu+\bar{\nu}}}{dE} = (1.01 \pm_{0.23}^{0.26}) \left(\frac{E}{100\text{TeV}} \right)^{-2.19 \pm 0.10} \cdot 10^{-18} \text{GeV}^{-1} \text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}.$$

An extra component at ~100 TeV ?

IceCube 6 year HESE data (1710.01191)



A multimessenger analysis confirms an 100 TeV excess




IceCube 6 year HESE data (1710.01191)

Very high energy neutrinos from N_D decays

(Anisimov,PDB,0812.5085;PDB, P.Ludl,S. Palomarez-Ruiz 1606.06238)

- Dark neutrinos unavoidably decay today into $A+\text{leptons}$ ($A=H,Z,W$) through the same mixing that produced them in the very early Universe
- The produced neutrinos can be responsible for the excess at ~ 100 TeV in IceCube

Example: $M_{DM}=300\text{TeV}$




(from 1606.06238)

Searches for Connections between Dark Matter and High-Energy Neutrinos with IceCube

IceCube Collaboration


(2205.12950)



2.5 σ significance when compared to the null hypothesis
best fit point: $m_D=386$ TeV, $\tau_D=2.8\times 10^{27}$ s

Absence of strong anisotropies

Arrival directions of most energetic neutrino events (HESE 6yr (magenta) & $\nu_\mu + \bar{\nu}_\mu$ 8yr (red))



This disfavors scenarios with strong Galactic emissions, the dominant component is of extra-galactic origin

Observation of high energy neutrinos from the Galactic plane

(IceCube 10 years data 2011-2021 2307.04427)





Figure 4: **All-sky point source search.** The best-fitting pre-trial significance for the all-sky search is shown as a function of direction in an Aitoff projection of the celestial sphere, in equatorial coordinates (J2000 equinox). The Galactic plane is indicated by a grey curve, and the Galactic Center as a dot. Although some locations appear to have significant emission, the trial factor for the number of points searched means these points are all individually statistically consistent with background fluctuations.

DM lifetime vs. mass plane: allowed regions

(P.Di Bari, K. Farrag, R. Samanta, Y. Zhou, 1908.00521)



What processes can thermalize the N_S -abundance prior to the oscillations?

Including Higgs portal interactions for N_S

(PDB, A. Murphy, arXiv 2210.10801)

$$\mathcal{L}_A = \frac{\lambda_{DS}}{\Lambda} \phi^\dagger \phi \overline{N_{DM}^c} N_S + \frac{\lambda_{SS}}{\Lambda} \phi^\dagger \phi \overline{N_S^c} N_S$$

effective scales

$$\tilde{\Lambda}_{DS} \equiv \Lambda / \lambda_{DS}$$

$$\tilde{\Lambda}_{SS} \equiv \Lambda / \lambda_{SS}$$

Can these interactions thermalise the source neutrinos prior to oscillations?
Let us modify the kinetic equations including these processes:

$$\frac{dN_{IJ}}{dt} = -i[\Delta H, N]_{IJ} - \begin{pmatrix} 0 & \frac{1}{2}(\Gamma_D + \Gamma_S)N_{DS} \\ \frac{1}{2}(\Gamma_D + \Gamma_S)N_{SD} & (\Gamma_D + \Gamma_S)(N_{N_S} - N_{N_S}^{eq}) + \frac{\langle \sigma_{\phi\phi \rightarrow N_S N_S^V} \rangle}{R^3} (N_{N_S}^2 - N_{N_S}^{eq2}) \end{pmatrix}$$


$$A(z) \equiv \frac{\langle \sigma_{\phi\phi \rightarrow N_S N_S^V} \rangle}{R^3 H z} = \frac{A(z=1)}{z^2}; \quad \langle \sigma_{\phi\phi \rightarrow N_S N_S^V} \rangle_{T \gg M_S} \simeq \frac{1}{\sim 2 \cdot 4\pi \Lambda_{SS}} \quad (\text{Kolb, Long, 1708.04293})$$

$$\Rightarrow A(z=1) \simeq g_N \frac{3}{16} \frac{\xi(3)}{\pi^3} \sqrt{\frac{90}{8\pi^3 g_R}} \frac{M_D M_{Pl}}{\sim 2 \Lambda_{SS}}$$

Condition for the thermalisation of the N_S abundance

(PDB, A. Murphy, arXiv 2210.10801)

$$\Rightarrow N_{N_S}(z_{in} \ll z \ll 1) - N_{N_S}(z_{in}) \simeq \frac{A_1}{z_{in}} \simeq 1.0 \times \left(\frac{T_{in}}{10^{16} \text{ GeV}} \right) \left(\frac{10^{16} \text{ GeV}}{\tilde{\Lambda}_{SS}} \right)^2 \simeq 1$$




$$\tilde{\Lambda}_{DS} \sim 10^{23} \text{ GeV}$$

$$\tilde{\Lambda}_{SS} \sim 10^{16} \text{ GeV}$$


↓
GUT
scale !!

The scale 10^{16} GeV maximises the production of DM



DM lifetime vs. mass plane: allowed regions


(PDB, A. Murphy, 2210.10801)



Decaying DM best fit (2.5σ) from IceCube 7.5 year data (2205.12950)

DM lifetime vs. mass plane: allowed regions


(PDB, A. Murphy, 2210.10801)



The scale of new physics cannot be made too much lower the GUT scale in order to explain the IceCube excess (respecting the LHAASO lower bound)



DM lifetime vs. mass plane: allowed regions

(PDB, A. Murphy, 2210.10801)



Upper bound on the seesaw (=leptogenesis) scale

(PDB, A. Murphy, 2210.10801)




The mechanism is compatible with (resonant) leptogenesis at a scale between 10 and 100 TeV

A possible GUT origin ? Heavy scalar H as mediator

(Anisimov,PDB, 2008; P.Ludl.PDB,S.Palomarez-Ruiz'16; Kolb and Long 1708.04293; PDB, A. Murphy, arXiv 2210.10801)

$$\mathcal{L}_H = \frac{1}{2} \partial_\mu H \partial^\mu H - \frac{1}{2} M_H^2 H^2 - \sum_{I,J} \lambda_{IJ} H \overline{N}_I^c N_J - \mu H \phi^\dagger \phi.$$




$$\mathcal{L}_H^{\text{eff}} = \frac{1}{2} \sum_{I,J,K,L} \frac{\lambda_{IJ} \lambda_{KL}}{M_H^2} (\overline{N}_I^c N_J) (\overline{N}_K^c N_L) + \frac{1}{2} \frac{\mu^2}{M_H^2} (\phi^\dagger \phi)^2 + \sum_{I,J} \frac{\mu \lambda_{IJ}}{M_H^2} \Phi^\dagger \Phi \overline{N}_I^c N_J. \quad \Rightarrow \quad \tilde{\Lambda}_{IJ} = \Lambda / \lambda_{IJ}, \text{ and } \Lambda = M_H^2 / \mu.$$

For $\mu \sim 10^9 \text{ GeV}$ one can have $\tilde{\Lambda}_{DS} \sim 10^{23} \text{ GeV}$ and $\lambda_{DS} \sim O(1)$ but one cannot reproduce simultaneously $\tilde{\Lambda}_{SS} \sim 10^{16} \text{ GeV}$ with the same scale Λ


A possible GUT origin? Heavy fermion F as mediator

(Anisimov, PDB, 2008; PDM, A. Murphy 2210.10801)

$$\mathcal{L}_F = \bar{F} (i \not{\partial} - M_F) F - \sum_I y_I (\bar{F} \phi N_I + \bar{N}_I \phi^\dagger F)$$



$$-\mathcal{L}_F^{\text{eff}} = \sum_{I,J} \frac{y_I y_J}{M_F} \bar{N}_I N_J \phi^\dagger \phi, \implies \Lambda = M_F \text{ and } \lambda'_{IJ} = y_I y_J.$$



This time one can have one scale $\Lambda = M_F \sim M_{\text{GUT}}$ and for $y_S \sim 1$ and $y_D \sim 10^{-7}$:

$$\tilde{\Lambda}_{DS} = \frac{\Lambda}{y_D y_S} \sim 10^{23} \text{ GeV} \quad \tilde{\Lambda}_{SS} = \frac{\Lambda}{y_S y_S} \sim \Lambda \sim 10^{16} \text{ GeV} \quad \tilde{\Lambda}_{DD} = \frac{\Lambda}{y_D y_D} \sim 10^{30} \text{ GeV}$$


$y_D \sim 10^{-7}$ could be understood as a small symmetry (e.g. Z_2) breaking parameter

Summary

- The DM puzzle might have a solution at higher scales than those traditionally explored so far and...
- ...heavy RH neutrinos provide an interesting option. An heavy RH neutrino playing the role of DM requires an extension of the usual type-I seesaw Lagrangian
- Higgs induced sterile-sterile neutrino mixing provides not only a way to produce dark neutrinos with the right abundance but also **it makes them detectable at neutrino telescopes.**
- Higgs portal interactions for the seesaw (source) neutrino enhance the dark neutrino production and allow to lift the scale of leptogenesis up to 100 TeV.
- Interestingly, the IceCube collaboration find an excess in the neutrino flux at energies well explained by RHINO DM decays (with $M_D \sim 100$ TeV) and further support comes from multimessenger astronomy
- Soon (?) new analysis of anisotropies in the IceCube high energy neutrino flux might provide a crucial test for heavy decaying DM
- **The emerging scale of new physics that can accommodate all constraints and also address the IceCube excess at ~ 100 TeV is $M_{GUT} \sim 10^{15} - 10^{16}$ GeV**

New frontiers

(SHIP proposal, 1504.04855)



Minimal seesaw mechanism (type I)

• Dirac + (right-right) Majorana mass term

(Minkowski '77; Gell-mann, Ramond, Slansky; Yanagida; Mohapatra, Senjanovic '79)

$$-\mathcal{L}_{mass}^{\nu} = \bar{\nu}_L m_D \nu_R + \frac{1}{2} \bar{\nu}_R^c M \nu_R + h.c. = -\frac{1}{2} (\bar{\nu}_L \quad \bar{\nu}_R^c) \begin{pmatrix} 0 & m_D^T \\ m_D & M \end{pmatrix} \begin{pmatrix} \nu_L \\ \nu_R^c \end{pmatrix} + h.c.$$

In the *see-saw limit* ($M \gg m_D$) the mass spectrum splits into 2 sets:


- 3 light **Majorana neutrinos** with masses (seesaw formula): $m_\nu = -m_D M^{-1} m_D^T \Rightarrow \text{diag}(m_1, m_2, m_3) = -U^\dagger m_\nu U^*$
- 3(?) very heavy Majorana neutrinos N_1, N_2, N_3 with $M_3 > M_2 > M_1 \gg m_D$

1 generation toy model :

$$m_D \sim m_{\text{top}},$$

$$m \sim m_{\text{atm}} \sim 50 \text{ meV}$$

$$\Rightarrow M \sim 10^{15} \text{ GeV}$$




Matter-antimatter asymmetry with leptogenesis

(Fukugita, Yanagida '86)

- Type I seesaw mechanism

- Thermal production of RH neutrinos: $T_{RH} \gtrsim T_{lep} \simeq M_i / (2 \div 10)$



total CP asymmetries $\varepsilon_I \equiv -\frac{\Gamma - \bar{\Gamma}}{\Gamma + \bar{\Gamma}}$

$\Rightarrow N_{B-L}^{fin} = \sum_{I=1,2,3} \varepsilon_I \times \underbrace{K_I^{fin}}_{\text{efficiency factors}}$


- Sphaleron processes in equilibrium

$\Rightarrow T_{RH} \gtrsim T_{sphalerons}^{off} \simeq 132 \text{ GeV}$

(Kuzmin, Rubakov, Shaposhnikov '85
D'Onofrio, Rummukainen, Tranberg 1404.3565)

$\Rightarrow \eta_{B0}^{lep} = \frac{a_{sph} N_{B-L}^{fin}}{N_\gamma^{rec}} \simeq 0.01 N_{B-L}^{fin}$

successful leptogenesis $\eta_{B0}^{lep} = \eta_{B0}^{obs} \sim 6 \cdot 10^{-10}$



Many proposed production mechanisms


Many production mechanisms have been proposed especially to address **IceCube** initially seemingly anomalous **PeV neutrino** events:

- from $SU(2)_R$ extra-gauge interactions (LRSM) (Fornengo, Niro, Fiorentin);
- from inflaton decays (Anisimov, PDB'08; Higaki, Kitano, Sato '14);
- from resonant annihilations through $SU(2)'$ extra-gauge interactions (Dev, Kazanas, Mohapatra, Teplitz, Zhang '16);
- From new $U(1)_Y$ interactions connecting DM to SM (Dev, Mohapatra, Zhang '16);
- From $U(1)_{B-L}$ interactions (Okada, Orikasa '12);

•

In all these models IceCube data are fitted through fine tuning of parameters responsible for decays (they are post-dictive)

Freeze-in solution for annihilating particles (FIMPs)



$$\Omega_{DM0} h^2 \propto \langle \sigma_{\text{ann}} \beta_{\text{rel}} \rangle$$

FIMPs evade all constraints, even too much: they are typically untestable!


Very high energy neutrinos from N_D decays

(Anisimov,PDB,0812.5085;PDB, P.Ludl,S. Palomarez-Ruiz 1606.06238)


- DM neutrinos unavoidably decay today into A +leptons ($A=H,Z,W$) through the same mixing that produced them in the very early Universe
- Potentially testable high energy neutrino contribution

Energy neutrino flux

Flavour composition at the detector




Hard component



Neutrino events at IceCube: 2 examples

$M_{DM} = 300 \text{ TeV}$



$M_{DM} = 8 \text{ PeV}$

