

# Possible late-time transition of $M_B$ inferred via neural networks

Kostas Dialektopoulos



Transilvania  
University  
of Brasov



Dark Side of the Universe 2024  
Corfu, Greece

Possible Late-time Transition of the absolute magnitude  
of Type Ia galaxies using Artificial Neural Networks

## Take home message

- ANNs are a model independent tool that can help us reconstruct cosmological parameters.
- We can use them to distinguish between the plethora of theories in the literature, based solely on the data without any physical or statistical assumption.


# Take home message

- ANNs are a model independent tool that can help us reconstruct cosmological parameters.
- We can use them to distinguish between the plethora of theories in the literature, based solely on the data without any physical or statistical assumption.


In this work,

- We use ANNs to agnostically constrain the value of  $M_B$  and assess the impact and statistical significance of a possible variation with redshift from the Pantheon+ compilation.
- We find an indication for a possible transition redshift at the  $z \approx 1$  region.

# What are Gaussian processes?



# What are Gaussian processes?





Definition: A GP is a stochastic (random) process where any finite subset is a **multivariate Gaussian distribution** with mean  $\mu(x)$  and covariance  $k(x, x')$ .

Setting each  $\mu(x)$  to zero, the **covariance function** can be used to learn the behavior that produced the data points.

# Gaussian Process Regression

- The covariance function contains **non-physical hyperparameters**  $\theta$  which define the distribution  $k(\theta, x, x')$ .
- Iterating over these values using Bayesian inference (or others) can produce better hyperparameters.
- The result is a **model independent reconstruction** (in physics) of the behavior of some parameter.
- This is superior to regular fitting because it is nonparametric and so **assumes no physical model** whatsoever.

# Squared Exponential $H_0$ GP (GaPP code: Seikel et al. 2012)



$$H_0 = 67.539 \pm 4.772 \text{ km/s/Mpc}$$

$$H_0 = 67.001 \pm 1.653 \text{ km/s/Mpc}$$

$$H_0 = 66.197 \pm 1.464 \text{ km/s/Mpc}$$


$$H_0 = 73.782 \pm 1.374 \text{ km/s/Mpc}$$

$$H_0 = 72.022 \pm 1.076 \text{ km/s/Mpc}$$


$$H_0 = 71.180 \pm 1.025 \text{ km/s/Mpc}$$

# Open problems with GP reconstructions


- **Overfitting:** GP is very prone to overfitting for small data sets, which is especially pronounced at the origin, i.e. Hubble constant



- **Kernel Selection Problem:** There is no natural kernel for cosmology





# Artificial Neural Networks (ANN)



ReFANN code from Wang et al. (2020)

# Training data for the ANN



$$P(z, \alpha, \lambda) = \frac{\lambda^\alpha}{\Gamma(\alpha)} z^{\alpha-1} e^{-\lambda z}$$

**Mean:**  $\sigma_H = 14.25 + 3.42z$

**Upper error:**  $\sigma_H = 21.37 + 10.79z$

**Lower error:**  $\sigma_H = 7.14 - 3.95z$

# Designing the ANN

- **Risk**: Optimizes the **number of hidden layers and neurons** in an ANN

$$\text{risk} = \sum_{i=1}^N (\text{Bias}_i^2 + \text{Variance}_i) = \sum_{i=1}^N \left( [H_{\text{obs}}(z_i) - H_{\text{pred}}(z_i)]^2 + \sigma_H^2(z_i) \right)$$

- **Loss**: Balances the **number of iterations** a system needs to predict the observational data

- 1 Least absolute deviation (**L1**)

$$L1 = \sum_{i=1}^N |H_{\text{obs}}(z_i) - H_{\text{pred}}(z_i)|$$

- 2 Smoothed L1 (**SL1**)
- 3 Mean Square Error (**MSE**)

$$\text{MSE} = \frac{1}{N} \sum_{i=1}^N (H_{\text{obs}}(z_i) - H_{\text{pred}}(z_i))^2$$



# Designing the ANN

What we use here

$$L_{\chi^2} = \sum_{i,j} [m_{\text{obs}}(z_i) - m_{\text{pred}}(z_i)]^T \mathcal{C}_{ij}^{-1} [m_{\text{obs}}(z_j) - m_{\text{pred}}(z_j)] ,$$

where  $\mathcal{C}_{ij}$  is the total noise covariance matrix of the data, which includes the statistical noise and systematics.


# Building the ANN



**Figure:** **Left:** Risk function for **one layer** (number of neurons  $2^n$ ,  $n \in 7, \dots, 14$ ), **Right:** Evolution of L1, SL1 and MSE loss functions

Reconstructing  $H(z)$  and  $H'(z)$ 

## Using the ANN (KD, Levi Said et al. '21) (KD, Mukherjee et al. '23)



**Figure:** Reconstructed reduced Hubble parameter from the (i) Pantheon SN compilation (left) and (ii) combined CC+BAO Hubble data set (right), using ANNs.

# Observational Datasets

- **Pantheon+**: SNIa observations from 1701 light curves that represent 1550 distinct SNIa spanning the redshift range  $z < 2.3$ .
- **CC**: 32  $H(z)$  measurements, along with the full covariance matrix that includes systematic and calibration errors, as reported in *Moresco, et al, Astrop. J. 2020*.

# ANN Training and Validation

- 1 We split the Pantheon+ dataset into training (70%) and validation (30%) sets and we train the network.
- 2 To incorporate the covariance matrix of the dataset, we minimize the  $\chi^2$  loss function.
- 3 The optimal network is one with two hidden layers and 128 neurons each.
- 4 The optimal network architecture is iterated over 500 times for random initialization of the hyperparameters along with the dropout effect. Out of these 500 samples, we compute the mean function and the respective uncertainties.

# Theoretical Framework

In a spatially flat Friedmann-Lemaître-Robertson-Walker universe, the luminosity distance is related to the Hubble parameter  $H(z)$  at some redshift  $z$ , as,

$$d_L(z) = c(1+z) \int_0^z \frac{d\tilde{z}}{H(\tilde{z})}.$$

# Theoretical Framework

In a spatially flat Friedmann-Lemaître-Robertson-Walker universe, the luminosity distance is related to the Hubble parameter  $H(z)$  at some redshift  $z$ , as,

$$d_L(z) = c(1+z) \int_0^z \frac{d\tilde{z}}{H(\tilde{z})}.$$

The observed luminosity of SNIa, from a specific redshift, is related to the apparent peak magnitude  $m$  via the following relation, independent of any physical model as,

$$m(z) - M_B = 5 \log_{10} \left[ \frac{d_L(z)}{1 \text{ Mpc}} \right] + 25.$$

# Theoretical Framework

We can rewrite the luminosity distance as,

$$d_L(z) = 10^{\frac{1}{5}[m(z) - M_B - 25]}.$$

and we can compute  $d'_L$ , the first order derivative of  $d_L$  with respect to the redshift  $z$  as,


$$d'_L(z) = \frac{\log(10)}{5} 10^{-\frac{M_B}{5}} m'(z).$$

Combining them, we can express the Hubble parameter as,

$$H(z) = \frac{c(1+z)^2}{(1+z)d'_L(z) - d_L(z)}.$$

In this way, we can derive the Hubble parameter  $H(z)$ , from the Pantheon+ apparent magnitudes  $m$  and its corresponding derivatives  $m'$  employing specific values of  $M_B$ .

# Constraints on $M_B$




**Figure:** ANN reconstruction of the Pantheon+ apparent magnitudes  $m(z)$  (left panel) and its corresponding derivatives  $m'(z)$  [right panel] at the CC redshifts ( $z_{CC}$ ).


# Constraints on $M_B$ [Mukherjee, KD, Levi-Said, Mifsud, 2402.10502 (JCAP)]

Reference	Methodology	Datasets	$M_B$
Camarena & Marra[144]	Cosmography	$\alpha_{\text{BAO}} + r_d^{\text{CMB}} + \text{Pantheon}$	$-19.401 \pm 0.027$
		$\theta_{\text{BAO}} + r_d^{\text{CMB}} + \text{Pantheon}$	$-19.262 \pm 0.030$
Mukherjee & Mukherjee[118]	<b>Gaussian Process</b>	<b>CC + Pantheon</b>	<b><math>-19.387 \pm 0.060</math></b>
Mukherjee & Banerjee[61]	<b>Gaussian Process</b>	<b>CC + Pantheon</b>	<b><math>-19.360 \pm 0.059</math></b>
		CC + $r_{\text{BAO}}$ + Pantheon	$-19.398 \pm 0.041$
Dinda & Banerjee[115]	<b>Gaussian Process</b>	<b>CC + Pantheon</b>	<b><math>-19.384 \pm 0.052</math></b>
		BAO + Pantheon	$-19.396 \pm 0.016$
		CC + BAO + Pantheon	$-19.395 \pm 0.015$
Benisty <i>et al.</i> [68]	Neural Networks	$\alpha_{\text{BAO}} + r_d^{\text{CMB}} + \text{Pantheon}$	$-19.38 \pm 0.20$
		$\alpha_{\text{BAO}} + r_d^{\text{SH0ES}} + \text{Pantheon}$	$-19.22 \pm 0.20$
Gómez-Valent[145]	Index of Inconsistency	CC + BAO + Pantheon	$-19.374 \pm 0.080$
		CC + BAO + Pantheon ( $\Omega_k \neq 0$ )	$-19.362^{+0.078}_{-0.067}$
Favale <i>et al.</i> [122]	Gaussian Process	CC + Pantheon+	$-19.344^{+0.116}_{-0.090}$
		CC + BAO + Pantheon+	$-19.314^{+0.086}_{-0.108}$
		CC + SH0ES + Pantheon+	$-19.252^{+0.024}_{-0.036}$
		CC + BAO + SH0ES + Pantheon+	$-19.252^{+0.024}_{-0.036}$
Banerjee <i>et al.</i> [119]	Gaussian Process	CC + $r_{\text{BAO}}$ + Pantheon+ ( $\Omega_k \neq 0$ )	$-19.404^{+0.099}_{-0.104}$
Shah <i>et al.</i> [111]	Neural Networks	Pantheon + $\alpha_{\text{BAO}} + r_d^{\text{CMB}}$	$-19.394^{+0.018}_{-0.017}$
		Pantheon + $\theta_{\text{BAO}} + r_d^{\text{CMB}}$	$-19.257^{+0.028}_{-0.027}$
<b>This work*</b>	<b>Neural Networks</b>	<b>CC &amp; Pantheon+</b>	<b><math>-19.353^{+0.073}_{-0.078}</math></b>

**Figure:** Comparison between the model-independent constraints on  $M_B$  obtained in this work vs those present in the literature.

# Constraints on $M_B$ [Mukherjee, KD, Levi-Said, Mifsud, 2402.10502 (JCAP)]



Cumulative binning method/Redshift layer binning with  $\bar{z}$ 

**Figure:** Predictions of the supernovae absolute magnitudes:  $M_B(z_{\max})$  by adopting cumulative binning, where  $M_B(z_{\max})$  is the derived value of  $M_B$  by considering CC  $H(z)$  data up to  $z_{\max}$  (left panel), and  $M_B(\bar{z})$  by adopting the redshift layer binning, where  $M_B(\bar{z})$  is the derived value of  $M_B$  by considering CC  $H(z)$  data within that redshift layer with a mean redshift  $\bar{z}$  (right panel). The purple region corresponds to the  $1 - \sigma$  model independent constraint  $M_B = -19.214 \pm 0.037$ , as inferred from the Pantheon and SH0ES data sets.

# Conclusions

- Given the observed apparent magnitudes  $m$  of the SNIa, we reconstruct  $m(z)$  using ANNs.
- We express the Hubble parameter inferred from SN-Ia as a function of its peak absolute magnitude.
- We obtain constraints on  $M_B$  by minimizing its negative log-likelihood. The result we get is  $M_B = -19.353^{+0.073}_{-0.078}$ .
- We test the evolution of  $M_B$  as a function of redshift and we find a possible transition at  $z \simeq 1$ .

# Conclusions


- Given the observed apparent magnitudes  $m$  of the SNIa, we reconstruct  $m(z)$  using ANNs.
- We express the Hubble parameter inferred from SN-Ia as a function of its peak absolute magnitude.
- We obtain constraints on  $M_B$  by minimizing its negative log-likelihood. The result we get is  $M_B = -19.353^{+0.073}_{-0.078}$ .
- We test the evolution of  $M_B$  as a function of redshift and we find a possible transition at  $z \simeq 1$ .

**Thank you!**

# $Om$ diagnostics (Sahni, Shafieloo, Starobinsky '08) (Shafieloo, Clarkson '10)

Distinguish  $\Lambda$ CDM from alternative dark energy and modified gravity models:

$$Om(z) = \frac{E^2(z) - 1}{(1+z)^3 - 1}.$$



**Figure:** Reconstructed  $Om$  diagnostics using (i) ANNs (left) and (ii) GPs (right) from the Pantheon SN data for three different priors.

# H<sub>0</sub> diagnostics (Krishnan, Colgáin, Sheikh-Jabbari, Yang '20)

It is defined as

$$H_0 = \frac{H(z)}{\sqrt{\Omega_{m0}(1+z)^3 + 1 - \Omega_{m0}}},$$

and its non-constancy suggests evidence for new physics beyond  $\Lambda$ CDM.




Figure: Reconstructed  $H_0$  diagnostics using (i) ANNs (left) and (ii) GPs (right)

# Constraining theories Arjona, Cardona, Nesseris '19

**Example:** Horndeski mapping:

$$G_2 = K(X), \quad G_3 = G(X), \quad G_4 = 1/2, \quad \text{and} \quad G_5 = 0,$$

The action is given by:


$$S = \int d^4x \sqrt{-g} \left( \frac{R}{2} - K(X) - G(X) \square \phi \right) + S_{\text{mat}}(\psi, g_{\mu\nu}).$$

Cosmological equations (flat FLRW):

$$K(X) = -3H_0^2 (1 - \Omega_{m0}) + \frac{\mathcal{J} \sqrt{2X} H^2(X)}{H_0^2 \Omega_{m0}} - \frac{\mathcal{J} \sqrt{2X} (1 - \Omega_{m0})}{\Omega_{m0}},$$

and


$$G_X(X) = -\frac{2\mathcal{J}H'(X)}{3H_0^2\Omega_{m0}}.$$




(KFD, Mukherjee, Levi Said, Mifsud '23)

We can also compute the DE EoS as

$$w_\phi = \frac{-K + \sqrt{2X\dot{X}}G_X}{K - 2X(K_X + 3\sqrt{2X}HG_X)}$$



**Figure:** Plots for dark energy EoS  $w_\phi(z)$  (left) and its compactified form  $\arctan(1 + w_\phi(z))$  (right) considering R21, TRGB, and P18  $H_0$  priors. The shaded regions with ‘-’, ‘|’ and ‘x’ hatches represent the  $1\sigma$  confidence levels for the above priors respectively.



**Figure:** Plots showing the posteriors of probability distribution of the compactified dark energy EoS for the theory at some sample redshifts for the R21, TRGB, and P18  $H_0$  priors, respectively.