

The QCD Axion Sum Rule


Workshop on Standard Model and Beyond

Corfu August 27- September 7 2023

Belén Gavela

Univ. Autónoma de Madrid and IFT

H2020



**Why axions
or ALPs ?**

Axions and ALPs a

are the tell-tale of hidden

symmetries

awaiting discovery

as they are (pseudo)Goldstone bosons

Many small unexplained SM parameters

Hidden symmetries
can explain small parameters




If spontaneously broken:
Goldstone bosons *a*

—> derivative couplings to SM particles

(Pseudo)Goldstone Bosons appear in many BSM theories

* e.g. Extra-dim Kaluza-Klein: 5d gauge field compactified to 4d

The Wilson line around the circle is a GB, which behaves as an axion in 4d



* Majorons, for dynamical neutrino masses

* From string models

* The Higgs itself may be a pGB ! (“composite Higgs” models)

* Axions **a** that solve the strong CP problem, and ALPs (axion-like particles)

.....

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?


$$\mathcal{L}_{\text{QCD}} = G_{\mu\nu} G^{\mu\nu} + \theta G_{\mu\nu} \tilde{G}^{\mu\nu}$$

where $\tilde{G}_{\mu\nu} = \epsilon_{\mu\nu\rho\sigma} G^{\rho\sigma}$

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?

$$\mathcal{L}_{\text{QCD}} = G_{\mu\nu} G^{\mu\nu} + \theta G_{\mu\nu} \tilde{G}^{\mu\nu}$$



$$\vec{E}^2 - \vec{B}^2 \qquad \theta \vec{E} \cdot \vec{B}$$

(CP even) (CP odd)

experimentally (neutron EDM): $\bar{\theta} \leq 10^{-10}$?

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?

$$\bar{\theta} \leq 10^{-10}$$




$$\mathcal{L}_{\text{QCD}} \supset \theta G_{\mu\nu} \tilde{G}^{\mu\nu}$$

$$\tilde{G}_{\mu\nu} = \epsilon_{\mu\nu\rho\sigma} G^{\rho\sigma}$$

A dynamical $U(1)_A$ solution ?

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?




$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

A dynamical $U(1)_A$ solution
→ **the axion a**

[Peccei+Quinn 77]
[Weinberg, 78]
[Wilczek, 78]

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?




$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

A dynamical $U(1)_A$ solution

→ the axion \mathbf{a}

It is a **pGB**:


$$V\left(\frac{\mathbf{a}}{f_a}\right)$$



$\mathbf{m}_a \neq 0$

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?




$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

A dynamical $U(1)_A$ solution

→ the axion \mathbf{a}

It is a **pGB**:

Excellent DM candidate !!




$\mathbf{m_a} \neq 0$

[Abbot+Sikivie, 83]
[Dine and W. Fischler, 83]
[Preskil et al, 91]

If axions or ALPs are the dark matter of the universe


e.g. for $m_a = 10^{-6}$ eV, inside each cm^{-3} there must be



about one thousand million axions per cm^{-3} !


In “true axion” models (= which solve the strong CP problem):

$$m_a f_a = \text{cte.}$$




In “true axion” models (= which solve the strong CP problem):

$$m_a f_a = \text{cte.}$$




In “true axion” models (= which solve the strong CP problem):

$$m_a f_a = \text{cte.}$$



In “true axion” models (= which solve the strong CP problem):

$$m_a f_a = \text{cte.}$$




The value of the constant is determined by the strong gauge group

In “true axion” models (= which solve the strong CP problem)

$$m_a f_a = \text{cte.}$$

* If the confining group is QCD:




$$V_{SM}\left(\frac{a}{f_a}\right) = -m_\pi^2 f_\pi^2 \sqrt{1 - \frac{4m_u m_d}{(m_u + m_d)^2} \sin^2\left(\frac{a}{2f_a}\right)}$$

In “true axion” models (= which solve the strong CP problem)

$$m_a f_a = \text{cte.}$$

* If the confining group is QCD:



$$\leftarrow \frac{a}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$


$$m_a^2 f_a^2 = m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$$

canonical QCD axion

In “true axion” models (= which solve the strong CP problem)

$$m_a f_a = \text{cte.}$$

* If the confining group is QCD:



$$\leftarrow \frac{a}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

$$m_a^2 f_a^2 = \underbrace{m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}}_{\chi_{\text{QCD}}}$$


canonical QCD axion

QCD topological susceptibility = χ_{QCD}

In “true axion” models (= which solve the strong CP problem)

$$m_a f_a = \text{cte.}$$

* If the confining group is QCD:



$$\leftarrow \frac{a}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

$$m_a^2 f_a^2 = \underbrace{m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}}_{\chi_{\text{QCD}}}$$


standard QCD axion

QCD topological susceptibility = χ_{QCD}

In “true axion” models (= which solve the strong CP problem)

$$m_a f_a = \text{cte.}$$

* If the confining group is QCD:



$$\leftarrow \frac{a}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

$$m_a^2 f_a^2 = \underbrace{m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}}_{\chi_{\text{QCD}}}$$

invisible axion

QCD topological susceptibility = χ_{QCD}

How come the QCD axion mass is NOT $\sim \Lambda_{\text{QCD}}$

Because two pseudo scalars couple to the QCD anomalous current :

$N_{\text{ps}} :$

η'_{QCD}

a

With only QCD:


one combination
must be (almost) massless

\rightarrow “Invisible axion”

$N_{\text{inst}} :$

$G \quad \tilde{G}$

QCD \uparrow



How come the QCD axion mass is NOT $\sim \Lambda_{\text{QCD}}$

Because two pseudo scalars couple to the QCD anomalous current :

N_{ps} :

η'_{QCD}

a

With only QCD:

**one combination
must be (almost) massless**

\rightarrow “Invisible axion”

N_{inst} :

$G \quad \tilde{G}$

QCD


The tiny axion mass is due to mixing
with η' and pion:

$$m_a^2 f_a^2 \sim m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$$

relation independent of the UV axion model

QCD: $m_a^2 f_a^2 = m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$

$10^{-5} < m_a < 10^{-2} \text{ eV} \quad , \quad 10^9 < f_a < 10^{12} \text{ GeV}$



Because of SN and hadronic data,
if axions light enough to be emitted

“Invisible axion”

Intensely looked for experimentally...

$\{m_a, 1/f_a\}$: direct **a** - gluon coupling





figure from cajohare.github.io

... and theoretically

e.g. Casper electric

$\{m_a, 1/f_a\}$: direct **a** - gluon coupling

$$\mathcal{L} \supset \frac{a}{f_a} \frac{\alpha_s}{8\pi} G\tilde{G}$$

$$\delta\mathcal{L} \equiv -\frac{i}{2} \frac{0.011 e}{m_n} \frac{a}{f_a} \bar{n} \sigma_{\mu\nu} \gamma_5 n F^{\mu\nu}$$
$$\equiv g_a \gamma n$$

Coupling to the
nEDM

$$m_a^2 f_a^2 \simeq m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$$

Axion mass

Intensely looked for experimentally...

$$g_{a\gamma} \sim C \frac{\alpha}{8\pi f_a} \quad (\text{GeV}^{-1})$$







figure from cajohare.github.io

... and theoretically

m_a (eV)

Intensely looked for experimentally...

$$g_{a\gamma} \sim C \frac{\alpha}{8\pi f_a} \quad (\text{GeV}^{-1})$$





[Farina et al, 17]
[Craig et al, 18]
[Di Luzio+Nardi et al, 17]


figure from cajohare.github.io

... and theoretically m_a (eV)

ALPs (axion-like particles) territory




ALPs (axion-like particles) territory




Difference between and ALP and a true axion:

an ALP does not intend to solve the strong CP problem

otherwise, the phenomenology is alike




ALPs (axion-like particles) territory




Difference between ALP and a true axion:

$$\{ m_a, f_a \}$$

are independent parameters



Axions and ALPs can explain Dark Matter




within the blueish bands
axions/ALPs would
account for all the DM

The field is BLOOMING


in Experiment ... and Theory



My task today: **can ALPs be true axions ?** (i.e. solve strong CP)



ALPs territory: can they be true axions ?




constraints in plot from
 Jaeckel+ Spannowsky 2015

“True” QCD axion


**“True” axion region
 amplifies?**

ALPs territory: can they be true axions ?



constraints in plot from
 Jaeckel+ Spannowsky 2015

ALPs territory: can they be true axions ?



constraints in plot from
 Jaeckel+ Spannowsky 2015

“True” QCD axion

“True” axion region amplifies?

but.....

**Let me revisit, and challenge,
the standard QCD wisdom**

``The QCD axion sum rule''

with Pablo Quilez and Maria Ramos, arXiv2305.15465

In “true axion” models (= which solve the strong CP problem):

$$m_a f_a = \text{cte.}$$

* If the confining group is QCD:

$$m_a^2 f_a^2 = m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$$

I am going to
challenge
this !


QCD topological susceptibility = χ_{QCD}

The Peccei-Quinn symmetry

PQ symmetry = a global $U(1)_A$ symmetry,

exact at classical level

but explicitly broken only by QCD instantons



QCD axion a

The Peccei-Quinn symmetry

PQ symmetry = a global $U(1)_A$ symmetry,

exact at classical level

but explicitly broken only by QCD instantons

→ QCD axion a

$$\mathcal{L}_{\text{QCD}} \supset \frac{a}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

$$m_a^2 f_a^2 = m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$$

The Peccei-Quinn symmetry

PQ symmetry = a global $U(1)_A$ symmetry,

exact at classical level

but explicitly broken only by QCD instantons

➔ QCD axion a

$$\mathcal{L}_{\text{QCD}} \supset \frac{a}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

$$m_a^2 f_a^2 = m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$$

This connection
assumes that
 a is a mass
eigenstate

ungranted !

The Peccei-Quinn symmetry

PQ symmetry = a global $U(1)_A$ symmetry,

exact at classical level

but explicitly broken only by QCD instantons

➔ QCD axion a

$$\mathcal{L}_{\text{QCD}} \supset \frac{a}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

$$m_a^2 f_a^2 = m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$$

to be more precise:
that it only mixes
with the η' , i.e.:

QCD eigenstate
= mass eigenstate


ungranted !

Remember:

In SM electroweak interactions families mix because

The weak interaction basis \neq the mass basis

(they are not simultaneously diagonal, unlike for QCD or QED)



In QCD-axion interactions, axions may mix because

The gluon interaction basis \neq the mass basis

(they are not necessarily simultaneously diagonal)

In QCD-axion interactions, axions may mix because

The gluon interaction basis \neq the mass basis

(they are not necessarily simultaneously diagonal)

**The axion field may not be the only singlet scalar in Nature.
It may mix with other singlet scalars**

**As long as the total scalar potential has a PQ symmetry,
the strong CP problem is solved**

coupling to gluons

Standard QCD axion: $\mathcal{L} = \frac{\alpha_s}{8\pi} \left(\frac{a_{G\tilde{G}}}{f_a} - \bar{\theta} \right) G\tilde{G}$

$$\Rightarrow m_a^2 f_a^2 = \chi_{\text{QCD}} \simeq m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$$

Instead, we can have:

$$\mathcal{L} = \frac{\alpha_s}{8\pi} \left(\frac{\hat{a}_{G\tilde{G}}}{F} - \bar{\theta} \right) G\tilde{G} - V'(\hat{a}_{G\tilde{G}}, \dots, \hat{a}_N)$$
$$\Rightarrow m_i^2 f_i^2 = g_i \chi_{\text{QCD}} \quad \text{within QCD}$$

distance to standard case ($g=1$)

Axion-exotic scalars mixing has appeared before in other constructions (clockwork, GUT, multiHiggs...)

Kim, Niles, Peloso 2005

Choi, Kim, Yun 2014

Kaplan, Ratazzi 2016

Giudice, McCullough 2017

Di Luzio et al. 2018

Fraser, Reece 2020

Darme et al. 2021

Chen et al. 2022

Agrawal Nee, Reig 2022

but, either by choice or by construction,
they took the limit where all but one decouple

Plan

- 1) A toy model with $N=2$ scalars
- 2) N fields and the most general PQ-invariant potential

N=2 toy example

$$\mathcal{L}_{N=2} = \left(\frac{a_{G\tilde{G}}}{F} + \theta \right) G\tilde{G} - V(a_{G\tilde{G}}, a_{\perp})$$

or equivalently:

$$\frac{a_{\tilde{G}}}{F} \equiv \frac{\hat{a}_1}{\hat{f}_1} + \frac{\hat{a}_2}{\hat{f}_2}$$

$$\mathcal{L}_{N=2} = \left(\frac{\hat{a}_1}{\hat{f}_1} + \frac{\hat{a}_2}{\hat{f}_2} + \theta \right) G\tilde{G} - V(\hat{a}_1, \hat{a}_2)$$

N=2 toy example

$$\mathcal{L}_{N=2} = \left(\frac{a_{G\tilde{G}}}{F} + \theta \right) G\tilde{G} - V(a_{G\tilde{G}}, a_{\perp})$$

$$\frac{a_{\tilde{G}}}{F} \equiv \frac{\hat{a}_1}{\hat{f}_1} + \frac{\hat{a}_2}{\hat{f}_2}$$

or equivalently:

$$\mathcal{L}_{N=2} = \left(\frac{\hat{a}_1}{\hat{f}_1} + \frac{\hat{a}_2}{\hat{f}_2} + \theta \right) G\tilde{G} - \frac{1}{2} \hat{m}_2^2 \hat{a}_2^2 \square$$

PQ symmetry : $\hat{a}_1 \rightarrow \hat{a}_1 - \theta \hat{f}_1$

N=2 toy example

$$\mathcal{L}_{N=2} = \left(\frac{a_{G\tilde{G}}}{F} + \theta \right) G\tilde{G} - V(a_{G\tilde{G}}, a_{\perp})$$

or equivalently:

$$\frac{a_{\tilde{G}}}{F} \equiv \frac{\hat{a}_1}{\hat{f}_1} + \frac{\hat{a}_2}{\hat{f}_2}$$

$$\mathcal{L}_{N=2} = \left(\frac{\hat{a}_1}{\hat{f}_1} + \frac{\hat{a}_2}{\hat{f}_2} + \theta \right) G\tilde{G} - \frac{1}{2} \hat{m}_2^2 \hat{a}_2^2$$

PQ symmetry : $\hat{a}_1 \rightarrow \hat{a}_1 - \theta \hat{f}_1$

After confinement (and for $\hat{f}_1 = \hat{f}_2 = \hat{f}$ and $r \equiv \frac{\hat{m}_2^2 \hat{f}^2}{\chi_{\text{QCD}}}$):

$$\mathbf{M}^2 = \frac{\chi_{\text{QCD}}}{\hat{f}^2} \begin{pmatrix} 1 & 1 \\ 1 & 1+r \end{pmatrix} \rightarrow \begin{matrix} \{m_1, f_1\} \\ \{m_2, f_2\} \end{matrix}$$

Both eigenstates couple to gluons: $\mathcal{L} \supset \frac{\alpha_s}{8\pi} \left[\frac{a_1}{f_1} + \frac{a_2}{f_2} \right] G\tilde{G}$

N=2 toy example

$$\mathcal{L}_{N=2} = \left(\frac{a_{G\tilde{G}}}{F} + \theta \right) G\tilde{G} - V(a_{G\tilde{G}}, a_{\perp})$$

or equivalently:

$$\frac{a_{\tilde{G}}}{F} \equiv \frac{\hat{a}_1}{\hat{f}_1} + \frac{\hat{a}_2}{\hat{f}_2}$$

$$\mathcal{L}_{N=2} = \left(\frac{\hat{a}_1}{\hat{f}_1} + \frac{\hat{a}_2}{\hat{f}_2} + \theta \right) G\tilde{G} - \frac{1}{2} \hat{m}_2^2 \hat{a}_2^2$$

PQ symmetry : $\hat{a}_1 \rightarrow \hat{a}_1 - \theta \hat{f}_1$

After confinement (and for $\hat{f}_1 = \hat{f}_2 = \hat{f}$ and $r \equiv \frac{\hat{m}_2^2 \hat{f}^2}{\chi_{\text{QCD}}}$):

$$\mathbf{M}^2 = \frac{\chi_{\text{QCD}}}{\hat{f}^2} \begin{pmatrix} 1 & 1 \\ 1 & 1+r \end{pmatrix} \rightarrow \begin{matrix} \{m_1, f_1\} \\ \{m_2, f_2\} \end{matrix}$$


Both eigenstates couple to gluons: $\mathcal{L} \supset \frac{\alpha_s}{8\pi} \left[\frac{a_1}{f_1} + \frac{a_2}{f_2} \right] G\tilde{G}$

Distance to standard QCD band

$$\{m_a, 1/f_a\}$$

$$g_i \equiv \frac{m_i^2 f_i^2}{m_a^2 f_a^2} \Big|_{\text{single QCD axion}}$$

in the N=2 toy model: $g_{1(2)} = \frac{2\sqrt{4+r^2}}{\sqrt{4+r^2} \pm (r-2)}$




Distance to standard QCD band

$$\{m_a, 1/f_a\}$$

$$g_i \equiv \frac{m_i^2 f_i^2}{m_a^2 f_a^2} \Big|_{\text{single QCD axion}}$$

in the N=2 toy model: $g_{1(2)} = \frac{2\sqrt{4+r^2}}{\sqrt{4+r^2} \pm (r-2)}$




Distance to standard QCD band

$$\{m_a, 1/f_a\}$$

$$g_i \equiv \frac{m_i^2 f_i^2}{m_a^2 f_a^2} \Big|_{\text{single QCD axion}}$$

$$\beta_i \equiv \frac{1}{g_i}$$

in the N=2 toy model: $g_{1(2)} = \frac{2\sqrt{4+r^2}}{\sqrt{4+r^2} \pm (r-2)}$



$$\frac{1}{g_1} + \frac{1}{g_2} \equiv \frac{\chi_{\text{QCD}}}{m_1^2 f_1^2} + \frac{\chi_{\text{QCD}}}{m_2^2 f_2^2} = 1.$$


Distance to standard QCD band

$$\{m_a, 1/f_a\}$$

$$g_i \equiv \frac{m_i^2 f_i^2}{m_a^2 f_a^2} \Big|_{\text{single QCD axion}}$$

$$\beta_i \equiv \frac{1}{g_i}$$

in the N=2 toy model: $g_{1(2)} = \frac{2\sqrt{4+r^2}}{\sqrt{4+r^2} \pm (r-2)}$




$$\beta_i \equiv \frac{1}{g_i}$$

$$\beta_1 + \beta_2 = 1$$


$\{m_a, 1/f_a\}$

**coupling
to gluons**



$\{m_a, 1/f_a\}$


**coupling
to gluons**



$\{m_a, 1/f_a\}$

Distance to standard QCD band


**coupling
to gluons**



Multiple QCD axion for N=2

$\{m_a, 1/f_a\}$


coupling
to gluons



Multiple QCD axion for N=2

$\{m_a, 1/f_a\}$

coupling
to gluons




Distance to standard QCD band

$$\{m_a, 1/f_a\}$$

$$g_i \equiv \frac{m_i^2 f_i^2}{m_a^2 f_a^2} \Big|_{\text{single QCD axion}}$$

$$\beta_i \equiv \frac{1}{g_i}$$

in the N=2 toy model: $g_{1(2)} = \frac{2\sqrt{4+r^2}}{\sqrt{4+r^2} \pm (r-2)}$



$$\frac{1}{g_1} + \frac{1}{g_2} \equiv \frac{\chi_{\text{QCD}}}{m_1^2 f_1^2} + \frac{\chi_{\text{QCD}}}{m_2^2 f_2^2} = 1.$$


Distance to standard QCD band

$$\{m_a, 1/f_a\}$$

$$g_i \equiv \frac{m_i^2 f_i^2}{m_a^2 f_a^2} \Big|_{\text{single QCD axion}}$$

$$\beta_i \equiv \frac{1}{g_i}$$

in the N=2 toy model: $g_{1(2)} = \frac{2\sqrt{4+r^2}}{\sqrt{4+r^2} \pm (r-2)}$



$$\frac{1}{g_1} + \frac{1}{g_2} \equiv \frac{\chi_{\text{QCD}}}{m_1^2 f_1^2} + \frac{\chi_{\text{QCD}}}{m_2^2 f_2^2} = 1.$$

Distance to standard QCD band


$$\{m_a, 1/f_a\}$$

$$g_i \equiv \frac{m_i^2 f_i^2}{m_a^2 f_a^2} \Big|_{\text{single QCD axion}}$$

$$\beta_i \equiv \frac{1}{g_i}$$


in the N=2 toy model:

$$g_{1(2)} = \frac{2\sqrt{4+r^2}}{\sqrt{4+r^2} \pm (r-2)}$$




Maxions (maximally deviated QCD axions): the maximal distance possible for the closest axion eigenstate is... **2**, and $g_1 = g_2 = 2$

N=2 QCD MAXION

 $\{m_a, 1/f_a\}$ 



N=2 QCD MAXION

 $\{m_a, 1/f_a\}$ 

General MAXION condition for N=2

In general, $N(N+1)/2$ maxion families

$$\mathbf{M}_{N=2}^2 = \frac{\chi_{\text{QCD}}}{\hat{f}^2} \begin{pmatrix} 2-p & 1 + \sqrt{p(2-p)} \\ 1 + \sqrt{p(2-p)} & 1+p \end{pmatrix}$$



Limiting case: Massless state has no mixing with gluons, the heavy one with mass $\sim 4 \frac{\chi_{\text{QCD}}}{\hat{f}^2}$

General potential for arbitrary N scalars

Exact results and sum rules

Multiple QCD axion for any N

$$\mathcal{L} = \frac{\alpha_s}{8\pi} \frac{a_{G\tilde{G}}}{F} G\tilde{G} - V_B^R(\underbrace{a_{G\tilde{G}}, \dots}_N)$$

Multiple QCD axion for any N and arbitrary potential

$$\mathcal{L} = \frac{\alpha_s}{8\pi} \frac{a_{G\tilde{G}}}{F} G\tilde{G} - V_B^R(\underbrace{a_{G\tilde{G}}, \dots}_N)$$

$$\mathbf{M}^2 = \mathbf{M}_{\text{QCD}}^2 + \mathbf{M}_{\text{ext}}^2 = \frac{\chi_{\text{QCD}}}{F^2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} b_{11} & -\frac{\chi_{\text{QCD}}}{F^2} \mathbf{X}^\dagger \\ \mathbf{X} & \mathbf{M}_1^2 \end{pmatrix}$$

eigenvalues: $m_i^2 = \beta_i m_i^2 + \langle a_i | \mathbf{M}_{\text{ext}}^2 | a_i \rangle$

Multiple QCD axion for any N and arbitrary potential

$$\mathcal{L} = \frac{\alpha_s}{8\pi} \frac{a_{G\tilde{G}}}{F} G\tilde{G} - V_B^R(\underbrace{a_{G\tilde{G}}, \dots}_N)$$

$$\mathbf{M}^2 = \mathbf{M}_{\text{QCD}}^2 + \mathbf{M}_{\text{ext}}^2 = \frac{\chi_{\text{QCD}}}{F^2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} b_{11} & -\frac{\chi_{\text{QCD}}}{F^2} \mathbf{X}^\dagger \\ \mathbf{X} & \mathbf{M}_1^2 \end{pmatrix}$$

eigenvalues: $m_i^2 = \beta_i m_i^2 + \langle a_i | \mathbf{M}_{\text{ext}}^2 | a_i \rangle$

β_i is the fraction of the total m_i due to QCD: the *QCD-axionness*

$$\beta_i \equiv \frac{1}{g_i}$$

Multiple QCD axion for any N and arbitrary potential

$$\mathcal{L} = \frac{\alpha_s}{8\pi} \frac{a_{G\tilde{G}}}{F} G\tilde{G} - V_B^R(\underbrace{a_{G\tilde{G}}, \dots}_N)$$

$$\mathbf{M}^2 = \mathbf{M}_{\text{QCD}}^2 + \mathbf{M}_{\text{ext}}^2 = \frac{\chi_{\text{QCD}}}{F^2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} b_{11} & -\frac{\chi_{\text{QCD}}}{F^2} \mathbf{X}^\dagger \\ \mathbf{X} & \mathbf{M}_1^2 \end{pmatrix}$$

eigenvalues: $m_i^2 = \beta_i m_i^2 + \langle a_i | \mathbf{M}_{\text{ext}}^2 | a_i \rangle$

β_i is the fraction of the total m_i due to QCD: the *QCD-axionness*

$$\beta_i \equiv \frac{1}{g_i}$$

$$g_i \equiv \frac{m_i f_i}{m_a f_a} \Big|_{\text{single QCD axion}} \geq 1$$

Multiple QCD axion for any N and arbitrary potential

$$\mathcal{L} = \frac{\alpha_s}{8\pi} \frac{a_{G\tilde{G}}}{F} G\tilde{G} - V_B^R(\underbrace{a_{G\tilde{G}}, \dots}_N)$$

$$\mathbf{M}^2 = \mathbf{M}_{\text{QCD}}^2 + \mathbf{M}_{\text{ext}}^2 = \frac{\chi_{\text{QCD}}}{F^2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} b_{11} & -\frac{\chi_{\text{QCD}}}{F^2} \mathbf{X}^\dagger \\ \mathbf{X} & \mathbf{M}_1^2 \end{pmatrix}$$

eigenvalues: $m_i^2 = \beta_i m_i^2 + \langle a_i | \mathbf{M}_{\text{ext}}^2 | a_i \rangle$

β_i is the fraction of the total m_i due to QCD: the *QCD-axionness*

$$\beta_i \equiv \frac{1}{g_i}$$

$$g_i \equiv \frac{m_i f_i}{m_a f_a} \Big|_{\text{single QCD axion}} \geq 1$$

1 PQ field \rightarrow N eigenvectors a_i coupled to $G\tilde{G}$

$$\mathcal{L} \supset \frac{\alpha_s}{8\pi} \frac{a_i}{f_i} G\tilde{G} \quad \frac{1}{F^2} = \sum_{i=1}^N \frac{1}{f_i^2}$$

Located to the right of the standard band

Several exact results follow from the eigenvalue-eigenvector theorem

Jacobi.....

PETER B. DENTON, STEPHEN J. PARKE, TERENCE TAO, AND XINING ZHANG

ABSTRACT. If A is an $n \times n$ Hermitian matrix with eigenvalues $\lambda_1(A), \dots, \lambda_n(A)$ and $i, j = 1, \dots, n$, then the j^{th} component $v_{i,j}$ of a unit eigenvector v_i associated to the eigenvalue $\lambda_i(A)$ is related to the eigenvalues $\lambda_1(M_j), \dots, \lambda_{n-1}(M_j)$ of the minor M_j of A formed by removing the j^{th} row and column by the formula

$$|v_{i,j}|^2 \prod_{k=1; k \neq i}^n (\lambda_i(A) - \lambda_k(A)) = \prod_{k=1}^{n-1} (\lambda_i(A) - \lambda_k(M_j)) .$$

We refer to this identity as the *eigenvector-eigenvalue identity*

<https://arxiv.org/pdf/1908.03795.pdf>

$$\mathcal{L} \supset \frac{\alpha_s}{8\pi} \frac{a_i}{f_i} G \tilde{G} \quad \text{with} \quad \frac{1}{f_i} = \frac{\langle a_{G\tilde{G}} | a_i \rangle}{F} \equiv \frac{v_{i1}}{F} \implies \sum_{i=1}^N \frac{1}{f_i^2} = \frac{1}{F^2}$$

Peccei-Quinn condition for arbitrary M

$$\lim_{\chi_{\text{QCD}} \rightarrow 0} \det \mathbf{M}^2 = 0 \implies \det \mathbf{M}_{\text{ext}} = 0$$

$$\frac{1}{F^2} = \sum_{i=1}^N \frac{1}{f_i^2}$$

$$\frac{\det \mathbf{M}^2}{\det \mathbf{M}_1^2} = \frac{f_\pi^2 m_\pi^2}{F^2} \frac{m_u m_d}{(m_u + m_d)^2}$$

PQ-invariance
condition
for arbitrary
potential


Peccei-Quinn condition for arbitrary M

$$\lim_{\chi_{\text{QCD}} \rightarrow 0} \det \mathbf{M}^2 = 0 \implies \det \mathbf{M}_{\text{ext}} = 0$$

$$\frac{1}{F^2} = \sum_{i=1}^N \frac{1}{f_i^2}$$

$$\frac{\det \mathbf{M}^2}{\det \mathbf{M}_1^2} = \frac{f_\pi^2 m_\pi^2}{F^2} \frac{m_u m_d}{(m_u + m_d)^2}$$

PQ-invariance
condition
for arbitrary
potential



$$\exists U(1)_{PQ} \implies \sum_{i=1}^N \frac{1}{g_i} = 1$$

PQ sum-rule

or equivalently

$$\exists U(1)_{PQ} \implies \sum_{i=1}^N \beta_i = 1; \quad \beta_i = \frac{1}{g_i}$$

QCD-axionness is shared

An intuitive view of the *QCD-axionness*

$$\beta_i \equiv \frac{1}{g_i}$$

field (or combination of) that has shift symmetry

eigenstate

field (or combination of) that couples to $G\tilde{G}$

$$\beta_i = \frac{1}{g_i} = \frac{\langle a_{\text{PQ}} | a_i \rangle \langle a_i | a_{G\tilde{G}} \rangle}{\langle a_{\text{PQ}} | \hat{a}_{G\tilde{G}} \rangle}$$

An intuitive view of the *QCD-axionness*

$$\beta_i \equiv \frac{1}{g_i}$$

field (or combination of) that has shift symmetry

eigenstate

field (or combination of) that couples to $G\tilde{G}$

$$\beta_i = \frac{1}{g_i} = \frac{\langle a_{PQ} | a_i \rangle \langle a_i | a_{G\tilde{G}} \rangle}{\langle a_{PQ} | \hat{a}_{G\tilde{G}} \rangle}$$

* e.g. in the N=2 toy model:

$$\mathcal{L}_{N=2} = \left(\frac{\hat{a}_1}{\hat{f}_1} + \frac{\hat{a}_2}{\hat{f}_2} + \theta \right) G\tilde{G} - \frac{1}{2} \hat{m}_2^2 \hat{a}_2^2$$

a_{PQ}

$a_{G\tilde{G}}$

An intuitive view of the QCD-axionness

$$\beta_i \equiv \frac{1}{g_i}$$

$|a_i\rangle$: eigenstates

$|a_{G\tilde{G}}\rangle$: field(s) that couple to $G\tilde{G}$

$|a_{PQ}\rangle$: field(s) that maintain shift invariance

then
$$\beta_i = \frac{1}{g_i} = \frac{\langle a_{PQ} | a_i \rangle \langle a_i | a_{G\tilde{G}} \rangle}{\langle a_{PQ} | a_{G\tilde{G}} \rangle}$$

and it can be proven that:

$$1 = \frac{\langle a_{PQ} | a_{G\tilde{G}} \rangle}{\langle a_{PQ} | a_{G\tilde{G}} \rangle} = \sum_i^N \frac{\langle a_{PQ} | a_i \rangle \langle a_i | a_{G\tilde{G}} \rangle}{\langle a_{PQ} | a_{G\tilde{G}} \rangle} = \sum_i^N \frac{\chi_{\text{QCD}}}{m_i^2 f_i^2} = \sum_i^N \frac{1}{g_i}$$

Maxions (maximally deviated QCD axions):

$$\max \left\{ \min_i \{g_i\} \right\} = N \quad \Longrightarrow \quad g_i = N, \quad \forall i$$


$$\text{Tr}[\mathbf{M}^2] = \sum_i m_i^2 = N \frac{\chi_{\text{QCD}}}{F^2}$$

N relations

Examples of $N > 2$ axions and Maxions


a multiple QCD axion for N=3

$\{m_a, 1/f_a\}$




a N=3 Maxion

$\{m_a, 1/f_a\}$




Potential=
Laguerre
matrices

Maxions




a N=8 Maxion


$\{m_a, 1/f_a\}$



$\{m_a, 1/f_a\}$: **coupling to gluons**



$\{m_a, 1/f_a\}$: **coupling to gluons**



Coupling to photons

Coupling to photons for the multiple QCD axion

Standard single QCD axion:

$$\mathcal{L} \supset \frac{\alpha_{em}}{2\pi} \left[\frac{E}{\mathcal{N}} - 1.92 \right] \frac{a}{f_a} F \tilde{F}$$

↑
model-dependent

Coupling to photons for the multiple QCD axion

Standard single QCD axion:

$$\mathcal{L} \supset \frac{\alpha_{em}}{2\pi} \left[\frac{E}{\mathcal{N}} - 1.92 \right] \frac{a}{f_a} F \tilde{F}$$

Multiple QCD axion:

$$\mathcal{L} \supset \frac{\alpha_{em}}{2\pi} \sum_i \left[\frac{E_i}{\mathcal{N}_i} - 1.92 \right] \frac{a_i}{f_i} F \tilde{F}$$

↑
model-dependent

Coupling to photons for the multiple QCD axion

Standard single QCD axion:

$$\mathcal{L} \supset \frac{\alpha_{em}}{2\pi} \left[\frac{E}{\mathcal{N}} - 1.92 \right] \frac{a}{f_a} F \tilde{F}$$

Multiple QCD axion:

$$\mathcal{L} \supset \frac{\alpha_{em}}{2\pi} \sum_i \left[\frac{E_i}{\mathcal{N}_i} - 1.92 \right] \frac{a_i}{f_i} F \tilde{F}$$

↑
model-dependent

if E_i / \mathcal{N}_i universal:


$$\mathcal{L} \supset \frac{\alpha_{em}}{2\pi} \left[\frac{E}{\mathcal{N}} - 1.92 \right] \sum_i \frac{a_i}{f_i} F \tilde{F}$$




$$\frac{m_i^2}{g_{a_i \gamma \gamma}^2} = \frac{m_a^2}{g_{a \gamma \gamma}^2} \Big|_{\text{single QCD axion}} \times g_i$$

$$\frac{(2\pi)^2 \chi_{\text{QCD}}}{\alpha_{em}^2} \left[\frac{E}{\mathcal{N}} - 1.92 \right]^{-2} \sum_{i=1}^N \frac{g_{a_i \gamma \gamma}}{m_i^2} = 1 \quad \text{sum-rule}$$

Coupling to photons for Maxions



Coupling to photons for Maxions



UV completions: one example

a simple KSVZ with 2 true QCD axions

$$\Psi_{1,2} \sim (3, 1, 0) \quad S_{1,2} \sim (1, 1, 0)$$

$$\mathcal{L}_{UV} = |\partial_\mu S_1|^2 + |\partial_\mu S_2|^2 + \bar{\Psi}_1 i \not{D} \Psi_1 + \bar{\Psi}_2 i \not{D} \Psi_2 - [y_1 \bar{\Psi}_1 \Psi_1 S_1 + y_2 \bar{\Psi}_2 \Psi_2 S_2 + \text{h.c.}] - V(S_{1,2})$$

$$S_i = \frac{1}{\sqrt{2}} \left(\hat{f}_i + \rho_i \right) e^{i\hat{a}_i/\hat{f}_i}$$

for instance $V(S_{1,2}) \sim S_2^4$ reduces the system to just one PQ

and gives precisely the first N=2 mass matrix I showed you !

a simple KSVZ with 2 true QCD axions

$$\Psi_{1,2} \sim (3, 1, 0) \quad S_{1,2} \sim (1, 1, 0)$$

$$\mathcal{L}_{UV} = |\partial_\mu S_1|^2 + |\partial_\mu S_2|^2 + \bar{\Psi}_1 i \not{D} \Psi_1 + \bar{\Psi}_2 i \not{D} \Psi_2 - [y_1 \bar{\Psi}_1 \Psi_1 S_1 + y_2 \bar{\Psi}_2 \Psi_2 S_2 + \text{h.c.}] - V(S_{1,2})$$

$$S_i = \frac{1}{\sqrt{2}} \left(\hat{f}_i + \rho_i \right) e^{i \hat{a}_i / \hat{f}_i}$$

for instance $V(S_{1,2}) = \lambda S_1^3 S_2 + \text{h.c.}$ reduces the system to just one PQ


$$V_{\text{eff}} = \frac{1}{2} \chi_{\text{QCD}} \left(\frac{\hat{a}_1 + \hat{a}_2}{\hat{f}} - \bar{\theta} \right)^2 + \frac{\lambda}{4} \hat{f}^4 \left(\frac{3\hat{a}_1 + \hat{a}_2}{\hat{f}} \right)^2$$

$$(\hat{f}_1 = \hat{f}_2 = \hat{f})$$



$$\mathbf{M}^2 = \frac{\chi_{\text{QCD}}}{F^2} \begin{pmatrix} 2 + 8r & -4r \\ -4r & 2r \end{pmatrix}$$

$$1/F^2 = 2/\hat{f}^2$$




Maxion solution for $r=1/5$

$$\text{Tr}[\mathbf{M}^2] = \sum_i m_i^2 = N \frac{\chi_{\text{QCD}}}{F^2}$$

How far from Λ_{QCD} must the new scales be to impact experiment?

Consider $N=3$ and an extra potential of the form:


$$\hat{\mathbf{M}}^2 = \frac{\chi_{\text{QCD}}}{\hat{f}^2} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 + \tilde{\lambda}_1 & 1 \\ 1 & 1 & 1 + \tilde{\lambda}_2 \end{pmatrix}$$


and ratios of the other two scales : $\tilde{\lambda}_1 = 10^{-3} \tilde{\lambda}_2 = 0.5$

This would lead to: $(g_1, g_2, g_3) \approx (1.2, 7.3, 497)$

—> Measuring g_1 and g_2 with enough precision would allow to infer the existence of a third axion even if $1/g_3 \ll 1$

right **ALP** territory: they can be pure QCD axions



constraints in plot from
 Jaeckel+ Spannowsky 2015


“True” QCD axion

Conclusions


- * **The PQ solution to the strong CP problem leads in all generality to multiple QCD axion signals**
 - > displaced to the right of the canonical QCD band
 - > the usual single QCD axion is just one limit
- * **The smoking gun is the multiplicity of signals.**
- * **Exact PQ invariance condition and exact PQ sum rule**
- * The main experimental impact is from scales not far from the QCD contribution
- * Beautiful synergy between different experiments.

$$g_{a\gamma} \sim \frac{\alpha}{8\pi f_a}$$

Log₁₀(GeV⁻¹)



Strong case for looking everywhere for a spin 0 particle with derivative couplings



cons. Jaeckel+ ...
wsky 2015

“True” QCD axion

“True” axion region has amplified

Conclusions / Outlook

It is a deep pleasure to be here today

Thank you very very much for the invitation!



Backup

Clockwork axions

$$\hat{\mathbf{M}}^2 = \frac{\chi_{\text{QCD}}}{\hat{f}^2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} + r \frac{\chi_{\text{QCD}}}{\hat{f}^2} \begin{pmatrix} 1 & -q & 0 \\ -q & 1 + q^2 & -q \\ 0 & -q & q^2 \end{pmatrix}$$

Certainly comply with the PQ condition: $\det \mathbf{M}^2 / \det \mathbf{M}_1^2 = \chi_{\text{QCD}} / F^2$

but do not allow Maxion solutions:

e.g. $q=1/3$

$$\text{Tr } \mathbf{M}^2 = N \frac{\chi_{\text{QCD}}}{F^2} \Leftrightarrow r = \frac{1}{10},$$

$$\text{Tr}^2 \mathbf{M}^2 - \text{Tr } \mathbf{M}^2 \cdot \text{Tr } \mathbf{M}^2 = N \frac{\chi_{\text{QCD}}}{F^2} \text{Tr } \mathbf{M}_1^2 \Leftrightarrow r = 0 \vee r = \frac{11}{182}$$

Not
compatible

$\{m_a, 1/f_a\}$: **coupling to gluons**

The single QCD axion line


$$\mathcal{L} \supset \frac{a}{f_a} \frac{\alpha_s}{8\pi} G\tilde{G}$$

$$\delta\mathcal{L} \equiv -\frac{i}{2} \frac{0.011 e a}{m_n f_a} \bar{n} \sigma_{\mu\nu} \gamma_5 n F^{\mu\nu} \equiv g_{a\gamma n}$$

**Coupling to the
nEDM**

$$m_a^2 f_a^2 \simeq m_\pi^2 f_\pi^2 \frac{m_u m_d}{(m_u + m_d)^2}$$

Axion mass



Adapted from AxionLimits
[Ciaran O'hare, 20]




Figure 1. Existing and projected (dashed lines) constraints on the parameter space of ALP-photon g_{aBB} and g_{aAB} couplings versus ALP mass and decay constant together with the lines corresponding to g_{aBB} (solid), $|g_{aAB}|$ (dashed) and $\sqrt{|g_{aAB}|g_{aBB}}$ (dash-dotted) in different hadronic axion models with one heavy PQ-charged fermion ψ with the parameters given in a box and $N_{\text{DW}} \equiv$

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?

$$\bar{\theta} \leq 10^{-10}$$



$$\mathcal{L}_{\text{QCD}} \supset \theta G_{\mu\nu} \tilde{G}^{\mu\nu}$$


$$\tilde{G}_{\mu\nu} = \epsilon_{\mu\nu\rho\sigma} G^{\rho\sigma}$$

A dynamical $U(1)_A$ solution ?

It substitutes θ by a spin 0 particle \mathbf{a} , i.e. a field $\mathbf{a}(x)$, which has a small potential with minimum at zero

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?



$$\mathcal{L}_{\text{QCD}} \supset \left(\frac{a}{f_a} - \theta \right) G_{\mu\nu} \tilde{G}^{\mu\nu}$$


scale f_a

A dynamical $U(1)_A$ solution

[Peccei+Quinn 77]
[Weinberg, 78]
[Wilczek, 78]

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?



$$\mathcal{L}_{\text{QCD}} \supset \left(\frac{\mathbf{a}}{f_a} - \boldsymbol{\theta} \right) G_{\mu\nu} \tilde{G}^{\mu\nu}$$

scale f_a


A dynamical $U(1)_A$ solution

[Peccei+Quinn 77]
[Weinberg, 78]
[Wilczek, 78]

with minimum at $\boldsymbol{\theta} f_a$: $\mathbf{a} = \boldsymbol{\theta} f_a + \mathbf{a}'$

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?



$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}'}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

← scale f_a


A dynamical $U(1)_A$ solution

$$\mathbf{a} = \theta f_a + \mathbf{a}'$$

[Peccei+Quinn 77]
[Weinberg, 78]
[Wilczek, 78]

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?



$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}'}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$


A dynamical $U(1)_A$ solution

→ the axion \mathbf{a}'

[Peccei+Quinn 77]
[Weinberg, 78]
[Wilczek, 78]

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?




$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

A dynamical $U(1)_A$ solution
→ the axion \mathbf{a}

[Peccei+Quinn 77]
[Weinberg, 78]
[Wilczek, 78]

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?



$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$


A dynamical $U(1)_A$ solution

→ **the axion a**

[Peccei+Quinn 77]
[Weinberg, 78]
[Wilczek, 78]

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?



$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$


A dynamical $U(1)_A$ solution

→ the axion \mathbf{a}

[Peccei+Quinn 77]
[Weinberg, 78]
[Wilczek, 78]

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?




$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

A dynamical $U(1)_A$ solution

→ the axion \mathbf{a}

It is a **pGB**:


$$V\left(\frac{a}{f_a}\right)$$



$\mathbf{m}_a \neq 0$

Strong motivation for singlet (pseudo)scalars from fundamental SM problems

The strong CP problem: Why is the QCD θ parameter so small?



$$\mathcal{L}_{\text{QCD}} \supset \frac{\mathbf{a}}{f_a} G_{\mu\nu} \tilde{G}^{\mu\nu}$$

A dynamical $U(1)_A$ solution
→ **the axion a**

[Peccei+Quinn 77]
[Weinberg, 78]
[Wilczek, 78]

It is a pGB: ~mainly derivative couplings

$$\partial_\mu a$$

Also excellent DM candidate

$$\mathbf{m}_a \neq 0$$

[Abbot+Sikivie, 83]
[Dine and W. Fischler, 83]
[Preskil et al, 91]

Maxions (maximally deviated QCD axions): N relations

$$p_{\mathbf{M}^2}(\lambda) \equiv \sum_{k=0}^N c_k^{\mathbf{M}} \lambda^k$$

$$c_k^{\mathbf{M}} = -N \frac{\chi_{\text{QCD}}}{F^2(N-k)} c_k^{\mathbf{M}_1}$$

Maxion
conditions