

Neutrino electromagnetic interactions: A window to new physics



24th Hellenic School and Workshops
on Elementary Particle Physics and
Gravity

Workshop on the
Standard Model and Beyond
01/09/2024

Alexander Studenikin
Moscow State University

a-studenik@yandex.ru

Within the Programme # 8 of National Centre for Physics and Mathematics "Effects of neutrino electromagnetic properties in neutrino propagation and interaction in matter"

Supported by Russian Science Foundation under grant No. 24-12-00084

Outline

- ① reminder of ν electromagnetic properties
- ② constraints on μ_ν , d_ν , g_ν and $\langle r_\nu^2 \rangle$
from laboratory experiments
- ③ effects of electromagnetic ν interactions in
astrophysics
- ④ astrophysical probes of electromagnetic ν
- ⑤ new effects in ν oscillations related to
electromagnetic ν interactions
... new phenomena in ν spin (flavor) oscillations in **moving** and
polarized mater and **magnetic field** of interest for astrophysical applications ...
- ⑥ conclusions – most recent constraints on ν emp
 - **SATURNE** tritium neutrino experiment -
first observation of *coherent elastic neutrino-atom scattering* (CEvAS) and search for ν **magnetic moment**



**Neutrino electromagnetic interactions:
A window to new physics**

+ upgrade:

Neutrino magnetic moment: a window to new physics

A. Studenikin*

Department of Theoretical Physics, Moscow State University, 119991 Moscow, Russia

A short review on a neutrino magnetic moment is presented.

Introduction. Experimental and theoretical studies of flavour conversion in solar, atmospheric, reactor and accelerator neutrino fluxes give strong evidence of non-zero neutrino mass. A massive neutrino can have non-trivial electromagnetic properties [1]. For a recent review on neutrino electromagnetic properties see [2].

The neutrino dipole moment (along with the electric dipole moment) is the most well studied among neutrino electromagnetic properties. The effective Lagrangian, that is in charge of a neutrino coupling to the electromagnetic field, can be written in the form

$$L_{int} = \frac{1}{2} \bar{\nu}_i \sigma_{\alpha\beta} (\mu_{ij} + \epsilon_{ij\gamma\delta} \nu_j) F^{\alpha\beta} + h.c. \quad (1)$$

where the magnetic moments μ_{ij} , in the presence of mixing between different neutrino states, are associated with the neutrino mass eigenstates ν_i . The interplay between magnetic moment and neutrino mixing effects is important. Note that electric (transition) moments ϵ_{ij} do also contribute to the coupling.

A Dirac neutrino may have non-zero diagonal electric moments in models where CP invariance is violated. For a Majorana neutrino the diagonal magnetic and electric moments are zero. Therefore, neutrino magnetic moments can be used to distinguish Dirac and Majorana neutrinos (see [3] and also [2] for a detailed discussion).

Neutrino magnetic moment in a minimal extension of Standard Model. The explicit evaluation of the one-loop contributions to the Dirac neutrino magnetic moment in the leading approximation over small parameters $b_i = \frac{m_i^2}{M_W^2}$ (m_i are the neutrino masses, $i = 1, 2, 3$), that however exceeds $\sim 10^{-10}$ μ_B , is

actly accounts for $a_l = \frac{g_l^2}{4\pi^2} (l = e, \mu, \tau)$, leads the following result [4],

$$\mu_l^D = \frac{e G_F m_l}{8\sqrt{2}\pi^2} \left(1 + \frac{m_l}{m_i}\right) \sum_{i=e,\mu,\tau} f(a_i) U_{ij} U_{il}^*, \quad (2)$$

where U_{ij} is the neutrino mixing matrix. The correspondent result in the absence of mixing was confirmed in [5,6]. A Majorana neutrino may also have transition moment of the value $\mu_{ij}^M = 2\mu_{ij}^D$ (see [2] for a detailed discussion and references).

For the diagonal magnetic moment of the Dirac neutrino, from (2) in the limit $a_i \ll 1$ the result [1] can be obtained

$$\mu_l^D = \frac{3e G_F m_l}{8\sqrt{2}\pi^2} \left(1 - \frac{1}{2} \sum_{i=e,\mu,\tau} a_i |U_{il}|^2\right). \quad (3)$$

The magnetic moment for hypothetical heavy neutrino was studied in [6]. In particular, it was obtained

$$\mu_{\nu} = \frac{e G_F m_{\nu}}{8\sqrt{2}\pi^2} \begin{cases} 3 + \frac{3}{2}b, & m_l \ll m_{\nu} \ll M_W, \\ 1, & m_l \ll M_W \ll m_{\nu}. \end{cases} \quad (4)$$

Note that the *LEP* data set a limit on number of light neutrinos coupled to Z boson. The numerical value of the Dirac neutrino magnetic moment within a minimal extension of the Standard Model, as it follows from (3), is

$$\mu_l^D \approx 3.2 \times 10^{-10} \left(\frac{m_i}{1 \text{ eV}}\right) \mu_B. \quad (5)$$

This is several orders of magnitude smaller than the present experimental limits if to account for the existed constraints on neutrino masses.

0920-5633/\$ - see front matter © 2009 Elsevier B.V. All rights reserved.

Detailed review on emp of

... > 500 citations

Electromagnetic properties of neutrino 2023, Physics of Particles and Nuclei 55 (2024) 1444

Studenikin,

Overview of neutrino electromagnetic properties (the theory, laboratory experiments and astrophysical probes), PoS (NuFact2021) 402(2022)052

Carlo Giunti*
INFN, Torino Section, Via P. Giuria 1, I-10125 Torino, Italy

Alexander Studenikin*
Department of Theoretical Physics, Faculty of Physics, Moscow State University and Joint Institute for Nuclear Research, Dubna, Russia

(published 16 June 2015)

A review is given of the theory and phenomenology of neutrino electromagnetic interactions, which provide powerful tools to probe the physics beyond the standard model. After a derivation of the general structure of the electromagnetic interactions of Dirac and Majorana neutrinos in the one-photon approximation, the effects of neutrino electromagnetic interactions in terrestrial experiments and in astrophysical environments are discussed. The experimental bounds on neutrino electromagnetic properties are presented and the predictions of theories beyond the standard model are confronted.

DOI: 10.1103/RevModPhys.87.531

PACS numbers: 14.60.St, 13.15.+g, 13.35.Hb, 14.60.Lm

CONTENTS

I. Introduction	531
II. Neutrino Masses and Mixing	532
A. Dirac neutrinos	533
B. Majorana neutrinos	533
C. Three-neutrino mixing	534
D. Neutrino oscillations	535
E. Status of three-neutrino mixing	538
F. Sterile neutrinos	540
III. Electromagnetic Form Factors	540
A. Dirac neutrinos	541
B. Majorana neutrinos	545
C. Massless Weyl neutrinos	546
IV. Magnetic and Electric Dipole Moments	547
A. Theoretical predictions for Dirac neutrinos	547
B. Theoretical predictions for Majorana neutrinos	549
C. Neutrino-electron elastic scattering	550
D. Effective magnetic moment	551
E. Experimental limits	553
F. Theoretical considerations	554

V. Radiative Decay and Related Processes	556
A. Radiative decay	556
B. Radiative decay in matter	559
C. Cherenkov radiation	560
D. Plasmon decay into a neutrino-antineutrino pair	561
E. Spin light	562
VI. Interactions with Electromagnetic Fields	563
A. Effective potential	564
B. Spin-flavor precession	565
C. Magnetic moment in a strong magnetic field	571
D. Beta decay of the neutron in a magnetic field	573
E. Neutrino pair production by an electron	574
F. Neutrino pair production by a strong magnetic field	575
G. Energy quantization in rotating media	576
VII. Charge and Anapole Form Factors	578
A. Neutrino electric charge	578
B. Neutrino charge radius	580
C. Neutrino anapole moment	583
VIII. Summary and Perspectives	585
Acknowledgments	585
References	585

... problem and puzzle ...

✓ electromagnetic properties
up to now nothing has been seen

... in spite of reasonable efforts ...

- results of terrestrial lab experiments on μ_0 (and ✓ EM properties in general)
- as well as data from astrophysics and cosmology

are in agreement with “ZERO”
✓ EM properties

... However, in course of recent development of knowledge on ✓ mixing and oscillations,


... Why ν electromagnetic properties are important ?

... Why ν em properties

to new physics ?




... How does it all relate to ν oscillations ?



$m_\nu \neq 0$




magnetic moment $\mu_\nu \neq 0$




Arthur McDonald

The Nobel Prize in Physics 2015

Takaaki Kajita



«for the discovery of neutrino oscillations, which shows that neutrinos have mass»



in Standard Model
 $m_\nu = 0 !!!$

In the easiest generalization of SM

$$\mu_{ii}^D = \frac{3eG_F m_i}{8\sqrt{2}\pi^2} \approx 3.2 \times 10^{-19} \left(\frac{m_i}{1 \text{ eV}} \right) \mu_B$$

Fujikawa, Shrock,
Phys.Rev.Lett. 45
(1980) 963

if $m_\nu \leq 0.45 \text{ eV}$ \ll new KATRIN limit arXiv: 2406.13516, 19 Jun 2024

then $\mu_{ii}^D \sim 1.8 \times 10^{-19} \mu_B$!

many orders of magnitude smaller than present experimental limits:

- $\mu_\nu \sim 10^{-11} \mu_B$ reactor \checkmark limits GEMMA 2012

- $\mu_\nu \sim 10^{-11} \div 10^{-12} \mu_B$ astrophysical ($\checkmark_{\text{solar}}$, \checkmark_{SN} , DM) limits
Borexino 2017 - XENONnT 2023, LUX-ZEPLIN 2023


- μ_ν is no less extravagant than possibility of $q_\nu \neq 0$

- limitations imposed by general principles of any theory are very strict

- $q_\nu \leq 3 \times 10^{-21} e$ from neutrality of hydrogen atom

- slightly weaker constraints are imposed by astrophysics

Studenikin, Tokarev, NPB, 2014 $q_\nu \leq 1.3 \times 10^{-19} e_0$



Direct neutrino-mass measurement based on 259 days of KATRIN data

Alexey Lokhov
on behalf of the
KATRIN collaboration

Karlsruhe Institute
of Technology,
Germany



XXXI International Conference on
Neutrino Physics and Astrophysics

Milano (Italy) - June 16-22, 2024



Alexey Lokhov
Univ. of Münster
& INR RAS


... PhD 2013
at Department of
Theoretical Physics,
MSU

Meeting of the MSU  group
Faculty of Physics, Moscow State University
January 2019

... a bit of  electromagnetic
properties theory ...

✓ electromagnetic vertex function

$$\langle \psi(p') | J_\mu^{EM} | \psi(p) \rangle = \bar{u}(p') \Lambda_\mu(q, l) u(p)$$



Matrix element of electromagnetic current is a Lorentz vector

$\Lambda_\mu(q, l)$ should be constructed using

matrices $\hat{1}, \gamma_5, \gamma_\mu, \gamma_5 \gamma_\mu, \sigma_{\mu\nu}$

tensors $g_{\mu\nu}, \epsilon_{\mu\nu\sigma\gamma}$

vectors q_μ and l_μ

$$q_\mu = p'_\mu - p_\mu, \quad l_\mu = p'_\mu + p_\mu$$

Lorentz covariance (1)
and electromagnetic gauge invariance (2)



Matrix element of electromagnetic current between neutrino states

$$\langle \nu(p') | J_\mu^{EM} | \nu(p) \rangle = \bar{u}(p') \Lambda_\mu(q) u(p)$$

where vertex function generally contains 4 form factors

$$\Lambda_\mu(q) = f_Q(q^2) \gamma_\mu + f_M(q^2) i \sigma_{\mu\nu} q^\nu - f_E(q^2) \sigma_{\mu\nu} q^\nu \gamma_5$$

1. electric

dipole

2. magnetic

3. electric

$$+ f_A(q^2) (q^2 \gamma_\mu - q_\mu \not{q}) \gamma_5$$

4. anapole

Hermiticity and discrete symmetries of EM current J_μ^{EM} put constraints on form factors

Dirac ✓

- 1) CP invariance + Hermiticity $\Rightarrow f_E = 0$,
- 2) at zero momentum transfer only electric Charge $f_Q(0)$ and magnetic moment $f_M(0)$ contribute to $H_{int} \sim J_\mu^{EM} A^\mu$

- 3) Hermiticity itself \Rightarrow three form factors are real: $Im f_Q = Im f_M = Im f_A = 0$

Majorana ✓

- 1) from CPT invariance (regardless CP or ~~CP~~).

$$f_Q = f_M = f_E = 0$$

...as early as 1939, W. Pauli...

EM properties \Rightarrow a way to distinguish Dirac and Majorana ✓

In general case matrix element of J_μ^{EM} can be considered between different initial $\psi_i(p)$ and final $\psi_j(p')$ states of different masses

$$\langle \psi_j(p') | J_\mu^{EM} | \psi_i(p) \rangle = \bar{u}_j(p') \Lambda_\mu(q) u_i(p)$$

$$p^2 = m_i^2, p'^2 = m_j^2:$$

... beyond SM...

and

$$\Lambda_\mu(q) = \left(f_Q(q^2)_{ij} + f_A(q^2)_{ij} \gamma_5 \right) (q^2 \gamma_\mu - q_\mu \not{q}) + f_M(q^2)_{ij} i \sigma_{\mu\nu} q^\nu + f_E(q^2)_{ij} \sigma_{\mu\nu} q^\nu \gamma_5$$



form factors are matrices in mass eigenstates space.



Dirac



(off-diagonal case $i \neq j$)

Majorana



1) Hermiticity itself does not apply restrictions on form factors,

1) CP invariance + hermiticity

2) CP invariance + Hermiticity

$$\mu_{ij}^M = 2\mu_{ij}^D \text{ and } \epsilon_{ij}^M = 0 \text{ or}$$

$$\mu_{ij}^M = 0 \text{ and } \epsilon_{ij}^M = 2\epsilon_{ij}^D$$

$f_Q(q^2), f_M(q^2), f_E(q^2), f_A(q^2)$ are relatively real (no relative phases).

... quite different EM properties ...

✓ form factors in gauge models

$$\langle \psi_j(p') | J_\mu^{EM} | \psi_i(p) \rangle = \bar{u}_j(p') \Lambda_\mu(q) u_i(p)$$

Form Factors at zero momentum transfer $q^2 = 0$ are elements of scattering matrix in any consistent theoretical model **FF** in matrix element

gauge independent and finite

...therefore...

- FF at $q^2 = 0$ determine static properties of that can be probed (measured) in direct interaction with external em fields


This is the case for $f_Q(q^2)$, $f_M(q^2)$, $f_E(q^2)$ in minimally extended SM ($f_A(q^2)$ is an exceptional case)

- In non-Abelian gauge models, FF at $q^2 \neq 0$ can be not invariant under gauge transformation because (in general) off-shell photon propagator is gauge dependent!

... One-photon approximation is not enough to get physical quantity...

... FF in matrix element cannot be directly measured in experiment with em field...

... FF can contribute to higher order processes accessible for experimental observation



... a bit more on \checkmark electromagnetic
properties theory

(*em* properties in gauge models)

ν em vertex function

The most general study of the
massive neutrino vertex function
(including electric and magnetic
form factors) in arbitrary R_ξ gauge
in the context of the SM + $SU(2)$ -singlet
 ν_R accounting for masses of particles
in polarization loops




M. Dvornikov, A. Studenikin

* Phys. Rev. D 63, 073001, 2004,

"Electric charge and magnetic moment of massive neutrino";

JETP 126 (2004), N 8, 1

* "Electromagnetic form factors of a massive neutrino."



charge

magnetic moment

$$\Delta_{\mu}(q) = \underbrace{f_Q(q^2)}_{\text{charge}} \gamma_{\mu} + \underbrace{f_M(q^2)}_{\text{magnetic moment}} i \sigma_{\mu\nu} q^{\nu} -$$

$$- \underbrace{f_E(q^2)}_{\text{electric moment}} i \sigma_{\mu\nu} q^{\nu} \gamma_5 - \underbrace{f_A(q^2)}_{\text{anapole moment}} (q^2 \gamma_{\mu} - q_{\mu} \not{q}) \gamma_5$$

electric moment

anapole moment

Direct calculations of complete set of one-loop contributions to ν vertex function in **minimally extended SM** (for a massive Dirac neutrino)

M.Dvornikov, A.Studenikin
PRD, 2004


... in case **CP** conservation

• $\Lambda_\mu(q) \Rightarrow f_Q(q^2), f_M(q^2), f_E(q^2), f_A(q^2)$

• Electric charge $f_Q(0) = 0$ and is **gauge-independent**

• Magnetic moment $f_M(0)$ **finite and gauge-independent**

• **Gauge and $q \times q$ dependence ...**



Magnetic moment dependence

$$\mu_\nu = \mu_\nu(m_\nu)$$

on neutrino mass 

3.2

Calculation of ν magnetic moment (massive ν , arbitrary R_ξ -gauge)



$$\Lambda_\mu(q) = f_Q(q^2) \gamma_\mu + f_M(q^2) i \sigma_{\mu\nu} q^\nu - f_E(q^2) \sigma_{\mu\nu} q^\nu \gamma_5 + f_A(q^2) (q^2 \gamma_\mu - q_\mu \not{q}) \gamma_5$$

magnetic moment

Dvornikov, Studenikin PRD 2004

$$\mu(a, b, \alpha) = f_M(q^2 = 0)$$

two mass parameters

$$a = \left(\frac{m_\ell}{M_W} \right)^2$$

$$b = \left(\frac{m_\nu}{M_W} \right)^2$$


$$\mu(a, b, \alpha) = \sum_{i=1}^6 \mu^{(i)}(a, b, \alpha)$$

and gauge-fixing parameter


$$\alpha = \frac{1}{\xi}$$

$\xi = 0$ - unitary gauge, $\xi = 1$ - 't Hooft-Feynman gauge


Proper vertices




(a)




(b)




(c)



(d)



(e)



(f)

Gauge and $q \times q$ dependence ...

Dvornikov,
Studenikin,
PRD 2004

✓ magnetic moment

● $\mu_e = \frac{3eG_F}{8\sqrt{2}\pi^2} m_{\nu_e}$

$\bar{f}_M(t)$

$$\bar{f}_M(t) = \sum_{i=1}^6 \bar{f}_M^{(i)}(t)$$



$\alpha = 100$
 $\alpha = 1$ ('t Hooft-Feynman)
 $\alpha = 0.1$

$$\alpha = \frac{1}{\xi}$$

$t \times 10^{-4} M_W^2$

$$f_M(q^2) = \frac{eG_F}{4\pi^2\sqrt{2}} m_{\nu} \sum_{i=1}^6 \bar{f}_M^{(i)}(q^2)$$

✓ dipole magnetic form factor



● $m_\nu \ll m_e \ll M_W$

light \checkmark

$$\mu_e = \frac{3eG_F}{8\sqrt{2}\pi^2} m_\nu$$

$$\mu_\nu = \frac{eG_F}{4\pi^2\sqrt{2}} m_\nu \frac{3}{4(1-a)^3} (2 - 7a + 6a^2 - 2a^2 \ln a - a^3) \quad a = \left(\frac{m_e}{M_W}\right)^2$$

Dvornikov,
Studenikin,
Phys.Rev.D 69
(2004) 073001;
JETP 99 (2004) 254

● $m_e \ll m_\nu \ll M_W$

intermediate \checkmark

Gabral-Rosetti,
Bernabeu,
Vidal, Zepeda,
Eur.Phys.J C 12
(2000) 633

$$\mu_\nu = \frac{3eG_F}{8\pi^2\sqrt{2}} m_\nu \left\{ 1 + \frac{5}{18} b \right\} \quad b = \left(\frac{m_\nu}{M_W}\right)^2$$

● $m_e \ll M_W \ll m_\nu$

$$\mu_\nu = \frac{eG_F}{8\pi^2\sqrt{2}} m_\nu$$

heavy \checkmark

$$\sim 10^{-19} \mu_B \left(\frac{m_\nu}{1\text{eV}}\right)$$


... μ_ν in case of mixing ...



Neutrino (beyond SM) dipole moments

P.Pal,
L.Wolfenstein
1982

(+ transition moments)



● Dirac neutrino

$$\left. \begin{matrix} \mu_{ij} \\ \epsilon_{ij} \end{matrix} \right\} = \frac{eG_F m_i}{8\sqrt{2}\pi^2} \left(1 \pm \frac{m_j}{m_i} \right) \sum_{l=e, \mu, \tau} f(r_l) U_{lj} U_{li}^*$$

$$r_l = \left(\frac{m_l}{m_W} \right)^2$$

- $m_e = 0.5 \text{ MeV}$
- $m_\mu = 105.7 \text{ MeV}$
- $m_\tau = 1.78 \text{ GeV}$
- $m_W = 80.2 \text{ GeV}$

● $m_i, m_j \ll m_l, m_W$

$$f(r_l) \approx \frac{3}{2} \left(1 - \frac{1}{2} r_l \right), \quad r_l \ll 1$$

transition moments vanish because unitarity of U implies that its rows or columns represent orthogonal vectors

● Majorana neutrino
only for

$$i \neq j$$

$$\mu_{ij}^M = 2\mu_{ij}^D \quad \text{and} \quad \epsilon_{ij}^M = 0$$

or

$$\mu_{ij}^M = 0 \quad \text{and} \quad \epsilon_{ij}^M = 2\epsilon_{ij}^D$$

● transition moments are suppressed, Glashow - Iliopoulos - Maiani cancellation,
● for diagonal moments there is no GIM cancellation

... depending on relative CP phase of ν_i and ν_j

The first nonzero contribution from **neutrino transition moments**

$$f_{r_l} \rightarrow -\cancel{\frac{3}{2}} + \frac{3}{4} \left(\frac{m_l}{m_W} \right)^2 \ll 1$$

GIM cancellation

$$\left. \begin{matrix} \mu_{ij} \\ \epsilon_{ij} \end{matrix} \right\} = \frac{3eG_F m_i}{32\sqrt{2}\pi^2} \left(1 \pm \frac{m_j}{m_i} \right) \left(\frac{m_\tau}{m_W} \right)^2 \sum_{l=e, \mu, \tau} \left(\frac{m_l}{m_\tau} \right)^2 U_{lj} U_{li}^*$$

$$\mu_B = \frac{e}{2m_e}$$

$$\left. \begin{matrix} \mu_{ij} \\ \epsilon_{ij} \end{matrix} \right\} = 4 \times 10^{-23} \mu_B \left(\frac{m_i \pm m_j}{1 \text{ eV}} \right) \sum_{l=e, \mu, \tau} \left(\frac{m_l}{m_\tau} \right)^2 U_{lj} U_{li}^*$$

... **neutrino radiative decay is very slow**

Dirac \checkmark **diagonal (i=j)** **magnetic moment**

$$\epsilon_{ii}^D = 0 \text{ for CP-invariant interactions}$$

$$\mu_{ii} = \frac{3eG_F m_i}{8\sqrt{2}\pi^2} \left(1 - \frac{1}{2} \sum_{l=e, \mu, \tau} r_l |U_{li}|^2 \right) \approx 3.2 \times 10^{-19} \left(\frac{m_i}{1 \text{ eV}} \right) \mu_B$$

$r_l = \left(\frac{m_l}{m_W} \right)^2$

$$\mu_{ii}^M = \epsilon_{ii}^M = 0$$

Lee, Shrock, Fujikawa, 1977

● no GIM cancellation

● μ_{ii}^D - to leading order - independent on U_{li} and $m_{l=e, \mu, \tau}$

$$\mu_e^2 = \sum_{i=1,2,3} |U_{ie}|^2 \mu_{ii}^2$$

...possibility to measure fundamental μ_{ii}^D


$\mu_{ii}^D = 0$ for **massless** \checkmark (in the absence of **right-handed charged currents**) \rightarrow

3.3 Naïve relationship between m_ν and μ_ν ■

... problem to get large μ_ν and still acceptable m_ν

If μ_ν is generated by physics beyond the SM at energy scale Λ ,

P. Vogel e.a., 2006




then

$$\mu_\nu \sim \frac{eG}{\Lambda}$$

...combination of constants and loop factors...

contribution to m_ν given by



, then

$$m_\nu \sim G\Lambda$$

$$m_\nu \sim \frac{\Lambda^2}{2m_e} \frac{\mu_\nu}{\mu_B} \sim \frac{\mu_\nu}{10^{-18} \mu_B} [\Lambda(\text{TeV})]^2 \text{ eV}$$


*Voloshin, 1988;
Barr, Freire,
Zee, 1990*

3.6 Neutrino magnetic moment in left-right symmetric models

$$SU_L(2) \times SU_R(2) \times U(1)$$

Gauge bosons $W_1 = W_L \cos \xi - W_R \sin \xi$
 mass states $W_2 = W_L \sin \xi + W_R \cos \xi$

with mixing angle ξ of gauge bosons $W_{L,R}$ with pure $(V \pm A)$ couplings



Kim, 1976; Marciano, Sanda, 1977
 Beg, Marciano, Ruderman, 1978

$$\mu_{\nu l} = \frac{eG_F}{2\sqrt{2}\pi^2} \left[m_l \left(1 - \frac{m_{W_1}^2}{m_{W_2}^2} \right) \sin 2\xi + \frac{3}{4} m_{\nu l} \left(1 + \frac{m_{W_1}^2}{m_{W_2}^2} \right) \right]$$

... charged lepton mass ...

... neutrino mass ...

Large magnetic moment $\mu_\nu = \tilde{\mu}_\nu (m_\nu, m_{\mathbf{e}^+}, m_{\mathbf{e}^-})$

- In the L-R symmetric models
($SU(2)_L \times SU(2)_R \times U(1)$)

Kim, 1976
Beg, Marciano,
Ruderman, 1978

- Voloshin, 1988
"On compatibility of small m_ν with large μ_ν of neutrino",
Sov.J.Nucl.Phys. 48 (1988) 512

K.Babu, Sidip Jana, M.Lindner,
"Large neutrino magnetic moments in the light
of recent experiments"
JHEP 10 (2020) 040

... there may be $SU(2)_\nu$
symmetry that forbids m_ν but not μ_ν

Z.Z.Xing, Y.L.Zhou,
"Enhanced electromagnetic transition dipole
moments and radiative decays of massive
neutrinos due to the seesaw-induced non-
unitary effects"

- Bar, Freire, Zee, 1990

- supersymmetry

considerable enhancement of μ_ν
to experimentally relevant range

Phys.Lett.B 715
(2012) 178

- extra dimensions

- model-independent constraint μ_ν

for BSM ($\Lambda \sim 1$ TeV) without fine tuning and under assumption

$\delta m_\nu \leq 1$ eV Bell,
Cirigliano,
Ramsey-Musolf,
Vogel, Wise,
2005

! $\mu_\nu^D \leq 10^{-15} \mu_B$

Dirac versus Majorana

$\mu_\nu^M \leq 10^{-14} \mu_B$!

... A remark on electric charge of ν ... Beyond Standard Model

ν neutrality $Q=0$ is attributed to

gauge invariance
+
anomaly cancellation constraints

imposed in SM of electroweak interactions

● ... General proof:

In SM:

$$Q = I_3 + \frac{Y}{2}$$

...from Gell-Mann - Nishijima ...

Foot, Joshi, Lew, Volkas, 1990;
Foot, Lew, Volkas, 1993;
Babu, Mohapatra, 1989, 1990
Foot, He (1991)

In SM (without ν_R) triangle anomalies cancellation constraints \rightarrow certain relations among particle hypercharges that is enough to fix all Y so that they, and consequently Q , are quantized

● $Q=0$ is proven also by direct calculation in SM within different gauges and methods

$Q=0$

● ... Strict requirements for Q quantization may disappear in extensions of standard $SU(2)_L \times U(1)_Y$ EW model if ν_R with $Y \neq 0$ are included: in the absence of Y quantization electric charges Q gets dequantized \rightarrow

Bardeen, Gastmans, Lautrup, 1972;
Cabral-Rosetti, Bernabeu, Vidal, Zepeda, 2000;
Beg, Marciano, Ruderman, 1978;
Marciano, Sirlin, 1980; Sakakibara, 1981;
● Dvornikov, Studenikin, 2004 (for SM in one-loop calculations)

millicharged ν

✓ charge radius and anapole moment

$$\Lambda_\mu(q) = \underbrace{f_Q(q^2)}_{\text{electric}} \gamma_\mu + \underbrace{f_M(q^2)}_{\text{magnetic}} i \sigma_{\mu\nu} q^\nu - \underbrace{f_E(q^2)}_{\text{dipole electric}} \sigma_{\mu\nu} q^\nu \gamma_5 + \underbrace{f_A(q^2)}_{\text{anapole}} (q^2 \gamma_\mu - q_\mu \not{q}) \gamma_5$$

- Although it is usually assumed that ✓ are electrically neutral (charge quant. Implies $Q \sim \frac{1}{3}e$), ✓ can be characterized by two ± charge distributions

$$f_Q(q^2) = f_Q(0) + q^2 \frac{df_Q}{dq^2}(0) + \dots, \text{ and } f_Q(q^2) \neq 0 \text{ for } q^2 \neq 0 \text{ even for electric charge } f_Q(0) = 0$$

✓ charge radius is introduced as

$$\langle r_\nu^2 \rangle = +6 \frac{df_Q}{dq^2}(0)$$

- for two-component massless left-handed Weyl spinors of SM

... it is often claimed for SM massless ✓ anapole moment

$$a_\nu = f_A(q^2) = \frac{1}{6} \langle r_\nu^2 \rangle \quad ???$$

... to be correct

Giunti, Studenikin
Rev.Mod.Phys.2015

$$\Lambda_{\text{SM}\mu}^{Q,A}(q) = (\gamma_\mu q^2 - q_\mu \not{q}) \mathbb{f}^{\text{SM}}(q^2)$$

$$\mathbb{f}^{\text{SM}}(q^2) = \tilde{f}_Q(q^2) - f_A(q^2) \xrightarrow{q^2 \rightarrow 0} \frac{\langle r^2 \rangle}{6} - a$$

... in SM charge radius and anapole moment are not defined separately ...

Interpretation of charge radius as an observable is rather delicate issue: $\langle r_\nu^2 \rangle$ represents a correction to tree-level electroweak scattering amplitude between ✓ and charged particles, which receives radiative corrections from several diagrams (including γ exchange) to be considered simultaneously → calculated CR is infinite and gauge dependent quantity. For ✓ with $m=0$, $\langle r_\nu^2 \rangle$ and a_ν can be defined (finite and gauge independent) from scattering cross section.

Bernabeu, Papavassiliou,
Vidal, Nucl.Phys. B 680
(2004) 450

??? For massive ✓ ???

The definition of the neutrino charge radius follows an analogy with the elastic electron scattering off a static spherically symmetric charged distribution of density $\rho(r)$ ($r = |\mathbf{x}|$), for which the differential cross section is determined [79–81] by the point particle cross section $\frac{d\sigma}{d\Omega}|_{point}$,

$$\frac{d\sigma}{d\Omega} = \frac{d\sigma}{d\Omega}|_{point} |f(q^2)|^2, \quad (90)$$

where the correspondent form factor $f(q^2)$ in the so-called *Breit frame*, in which $q_0 = 0$, can be expressed as

$$f(q^2) = \int \rho(r) e^{i\mathbf{q}\cdot\mathbf{x}} d^3x = 4\pi \int dr r^2 \rho(r) \frac{\sin(qr)}{qr}, \quad (91)$$

here $q = |\mathbf{q}|$. Thus, one has

$$\frac{df_Q}{dq^2} = \int \rho(r) \frac{qr \cos(qr) - \sin(qr)}{2q^{3/2}r} d^3x. \quad (92)$$

In the case of small q , we have $\lim_{q^2 \rightarrow 0} \frac{qr \cos(qr) - \sin(qr)}{2q^{3/2}r} = -\frac{r^2}{6}$ and

$$f(q^2) = 1 - |\mathbf{q}|^2 \frac{\langle r^2 \rangle}{6} + \dots \quad (93)$$


Therefore, the neutrino charge radius (in fact, it is the charge radius squared) is usually defined by

$$\langle r_\nu^2 \rangle = -6 \frac{df_Q(q^2)}{dq^2} \Big|_{q^2=0}. \quad (94)$$

Since the neutrino charge density is not a positively defined quantity, $\langle r_\nu^2 \rangle$ can be negative.

To obtain ν electroweak radius as **physical**
(finite, not divergent) **quantity**

Bernabeu,
Papavassiliou,
Vidal, 2004



$$\langle r_{\nu_i}^2 \rangle = \frac{G_F}{4\sqrt{2}\pi^2} \left[3 - 2 \log \left(\frac{m_i^2}{m_W^2} \right) \right] \quad i = e, \mu, \tau$$

$$\langle r_{\nu_e}^2 \rangle = 4 \times 10^{-33} \text{ cm}^2$$

...contribution to ν - e
scattering experiments
through (not the whole story,
off-diagonal charge radius)

Contribution of box diagram to

$$\nu_l + l' \rightarrow \nu_l + l'$$

$$g_V \rightarrow \frac{1}{2} + 2 \sin^2 \theta_W + \frac{2}{3} m_W^2 \langle r_{\nu_e}^2 \rangle \sin^2 \theta_W$$

- ... theoretical predictions and present experimental limits are in agreement within one order of magnitude...

Experimental constraints

(from studies of terrestrial and astrophysical ν fluxes)

on

μ_ν , q_ν

and

$\langle r_\nu^2 \rangle$

magnetic moment

millicharge

charge radius

Particle Data Group
Review of Particle Physics 2022, update 2023
and 2024

R.L.Workman et al. (Particle Data Group Collaboration),
Progress of Theoretical and Experimental Physics,
vol. 2022, no. 8, 083C01

S. Navas et al. (Particle Data Group Collaboration)
Phys. Rev. D 110 (2024) 030001

✓ magnetic moment

... most easily accepted are
dipole magnetic and electric moments

however most accessible for experimental
studies are charge radii $\langle r_{\nu}^2 \rangle$

Studies of ν - e scattering

- most sensitive method for experimental investigation of μ_ν

Cross-section:

$$\bullet \quad \frac{d\sigma}{dT}(\nu + e \rightarrow \nu + e) = \left(\frac{d\sigma}{dT}\right)_{\text{SM}} + \left(\frac{d\sigma}{dT}\right)_{\mu_\nu}$$

where the Standard Model contribution

$$\bullet \quad \left(\frac{d\sigma}{dT}\right)_{\text{SM}} = \frac{G_F^2 m_e}{2\pi} \left[(g_V + g_A)^2 + (g_V - g_A)^2 \left(1 - \frac{T}{E_\nu}\right)^2 + (g_A^2 - g_V^2) \frac{m_e T}{E_\nu^2} \right],$$

T is the electron recoil energy and

$$\bullet \quad \left(\frac{d\sigma}{dT}\right)_{\mu_\nu} = \frac{\pi \alpha_{em}^2}{m_e^2} \left[\frac{1 - T/E_\nu}{T} \right] \mu_\nu^2$$

$$\mu_\nu^2(\nu_l, L, E_\nu) = \sum_j \left| \sum_i U_{li} e^{-iE_i L} \mu_{ji} \right|^2$$

$$\mu_{ij} \rightarrow |\mu_{ij} - \epsilon_{ij}|$$

$$g_V = \begin{cases} 2 \sin^2 \theta_W + \frac{1}{2} & \text{for } \nu_e, \\ 2 \sin^2 \theta_W - \frac{1}{2} & \text{for } \nu_\mu, \nu_\tau, \end{cases} \quad g_A = \begin{cases} \frac{1}{2} & \text{for } \nu_e, \\ -\frac{1}{2} & \text{for } \nu_\mu, \nu_\tau \end{cases}$$

for anti-neutrinos
 $g_A \rightarrow -g_A$


• to incorporate charge radius: $g_V \rightarrow g_V + \frac{2}{3} M_W^2 \langle r^2 \rangle \sin^2 \theta_W$????

Magnetic moment contribution dominates at low electron recoil energies when

recoil energies when $\left(\frac{d\sigma}{dT}\right)_{\mu\nu} > \left(\frac{d\sigma}{dT}\right)_{SM}$ and

$$\frac{T}{m_e} < \frac{\pi^2 \alpha_{em}}{G_F^2 m_e^4} \mu_\nu^2$$

... the lower the smallest measurable electron recoil energy is, smaller values of μ_ν^2 can be probed in scattering experiments ...



... courtesy of A.Starostin...

Калининской атомной станции (Удомля, Тверская область)



GEMMA (2005 – 2012 - running) Germanium Experiment for Measurement of Magnetic Moment of Antineutrino

JINR (Dubna) + ITEP (Moscow) at Kalinin Nuclear Power Plant



World best experimental (reactor) limit

$$\mu_\nu < 2.9 \times 10^{-11} \mu_B$$

June 2012



A.Beda et al, in:

Special Issue on “Neutrino Physics”,
Advances in High Energy Physics (2012) 2012,
editors: J. Bernabeu, G. Fogli, A. McDonald, K. Nishikawa

... quite realistic prospects for future ...

● GEMMA-2 / ν GeN experiment

... searching for μ_ν and CE ν NS **unprecedentedly low threshold** $T \sim 200$ eV

$$\mu_\nu \sim (5 - 9) \times 10^{-12} \mu_B$$

2024 + to appear soon ?


● first results on CE ν NS: I.Alekseev et al., Phys.Rev.D 106 (2022) 5, L051101
coherent elastic neutrino–nucleus scattering

... claim that

ν - e cross section

should be increased by

Atomic Ionization effect:





?

H.Wong et al. (TEXONO Coll.),
arXiv: 1001.2074,
13 Jan 2010,
reported at

Neutrino 2010 Conference
(Athens, June 2010),
PRL 105 (2010)

061801

...new bounds ...



...much better limits on ν effective magnetic moment ...

H.Wong et al.,
(TEXONO Coll.),
arXiv: 1001.2074,
13 Jan 2010,

$$\mu_\nu < 1.3 \times 10^{-11} \mu_B$$




... atomic ionization effect accounted for ...

PRL 105 (2010)
061801

Neutrino 2010 Conference, Athens

$$\mu_\nu < 5.0 \times 10^{-12} \mu_B$$



... however .. 


... atomic ionization effect accounted for ...

A.Beda et al.
(GEMMA Coll.),
arXiv: 1005.2736,
16 May 2010

$$\mu_\nu < 3.2 \times 10^{-11} \mu_B$$



... ν -e scattering on free electrons ...
(without atomic ionization)



K.Kouzakov, A.Studenikin

- Magnetic neutrino scattering on atomic electrons revisited, **Phys.Lett. B** 105 (2011) 061801,
- Electromagnetic neutrino-atom collisions: The role of electron binding, **Nucl.Phys.** (Proc.Suppl.) 217 (2011) 353

K.Kouzakov, A.Studenikin, M.Voloshin

- Neutrino electromagnetic properties and new bounds on neutrino magnetic moments, **J.Phys.: Conf.Ser.** 375 (2012) 042045
- Neutrino-impact ionization of atoms in search for neutrino magnetic moment, **Phys.Rev.D** 83 (2011) 113001
- On neutrino-atom scattering in searches for neutrino magnetic Moments, **Nucl.Phys.B** (Proc.Supp.) 2011 (Proc. of Neutrino 2010 Conf.)
- Testing neutrino magnetic moment in ionization of atoms by neutrino impact, **JETP Lett.** 93 (2011) 699

M.Voloshin

- Neutrino scattering on atomic electrons in search for neutrino magnetic moment, **Phys.Rev.Lett.** 105 (2010) 201801

— No important effect of
Atomic Ionization on cross section in
 μ_ν experiments once all possible final
electronic states accounted for

...free electron approximation ...

M.Voloshin, 23 Aug 2010;

K.Kouzakov, A.Studenikin, 26 Nov 2010;

H.Wong et al, arXiv: 1001.2074 V3, 28 Nov 2010

K. Kouzakov, A. Studenikin,

“Theory of neutrino-atom collisions:
the history, present status, and BSM physics”,

in: Special issue

“Through Neutrino Eyes: The Search for New Physics”,
Adv. in High Energy Phys. 2014 (2014) 569409 (37pp)


editors: J. Bernabeu, G. Fogli, A. McDonald, K. Nishikawa



Limiting the effective magnetic moment of solar neutrinos with the Borexino detector

Livia Ludhova
on behalf of
the Borexino collaboration

IKP-2 FZ Jülich,
RWTH Aachen,
and JARA Institute, Germany



Phys. Rev. D 96 (2017) 091103

Limiting μ_ν with Borexino Phase-II solar neutrino data



BOREXINO Collaboration (2017)

NMM results from Phase 2



Data selection:

Fiducial volume: $R < 3.021$ m, $|z| < 1.67$ m
 Muon, ^{214}Bi - ^{214}Po , and noise suppression

Free fit parameters: solar- ν (pp, ^7Be) and backgrounds (^{85}Kr , ^{210}Po , ^{210}Bi , ^{11}C , external bgr.), **response parameters** (light yield, ^{210}Po position and width, ^{11}C edge (2×511 keV), 2 energy resolution parameters)

Constrained parameters: ^{14}C , pile up


Fixed parameters: pep-, CNO-, ^8B - ν rates

Systematics: treatment of pile-up, energy estimators, pep and CNO constraints with LZ and HZ SSM

Without radiochemical constraint
 $\mu_{\text{eff}} < 4.0 \times 10^{-11} \mu_{\text{B}}$ (90% C.L.)

With radiochemical constraint
 $\mu_{\text{eff}} < 2.6 \times 10^{-11} \mu_{\text{B}}$ (90% C.L.)
 adding systematics

$\mu_{\text{eff}} < 2.8 \times 10^{-11} \mu_{\text{B}}$ (90% C.L.)



Effective ν magnetic moment in experiments

(for neutrino produced as ν_l with energy E_ν and after traveling a distance L)

$$\mu_\nu^2(\nu_l, L, E_\nu) = \sum_j \left| \sum_i U_{li} e^{-iE_i L} \mu_{ji} \right|^2$$


where neutrino mixing matrix

$$\mu_{ij} \equiv |\beta_{ij} - \epsilon_{ij}|$$

magnetic and electric moments

Observable μ_ν is an effective parameter that depends on neutrino flavour composition at the detector.

Implications of μ_ν limits from different experiments (reactor, solar ^8B and ^7Be) are different.



... comprehensive analysis of ν - e scattering
with account for ν mixing and oscillations ...

PHYSICAL REVIEW D **95**, 055013 (2017)

Electromagnetic properties of massive neutrinos in low-energy elastic neutrino-electron scattering

Konstantin A. Kouzakov[†]

*Department of Nuclear Physics and Quantum Theory of Collisions, Faculty of Physics,
Lomonosov Moscow State University, Moscow 119991, Russia*

Alexander I. Studenikin[†]

*Department of Theoretical Physics, Faculty of Physics, Lomonosov Moscow State University,
Moscow 119991, Russia
and Joint Institute for Nuclear Research, Dubna 141980, Moscow Region, Russia
(Received 11 February 2017; published 14 March 2017)*

A thorough account of electromagnetic interactions of massive neutrinos in the theoretical formulation of low-energy elastic neutrino-electron scattering is given. The formalism of neutrino charge, magnetic, electric, and anapole form factors defined as matrices in the mass basis is employed under the assumption of three-neutrino mixing. The flavor change of neutrinos traveling from the source to the detector is taken into account and the role of the source-detector distance is inspected. The effects of neutrino flavor-transition millicharges and charge radii in the scattering experiments are pointed out.

● **Short-baselin case** $L \ll L_{kk'} = 2E_\nu / |\delta m_{kk'}^2| \rightarrow e^{-i(\delta m_{kk'}^2 / 2E_\nu)L} = 1$

● $P_{\nu_\ell \rightarrow \nu_\ell}(L, E_\nu) = \delta_{\ell e} \quad \mathcal{A}_{\nu_\ell \rightarrow \nu_{\ell'}}(L, E_\nu) \mathcal{A}_{\nu_{\ell'} \rightarrow \nu_{\ell''}}^*(L, E_\nu) = \delta_{\ell \ell'} \delta_{\ell \ell''}$

effect of \checkmark flavor change is insignificant
 $(\nu_\ell(L))$ is as in the source

$C_1 = (g_V + \delta_{\ell e} + \tilde{Q}_{\ell\ell})^2 + \sum_{\ell'=e,\mu,\tau} (1 - \delta_{\ell \ell'}) |\tilde{Q}_{\ell \ell'}|^2 \quad C_2 = (g_A + \delta_{\ell e})^2$

$C_3 = (g_V + \delta_{\ell e})(g_A + \delta_{\ell e}) + (g_A + \delta_{\ell e})\tilde{Q}_{\ell\ell}$

weak-electromagnetic interference term contains only
 flavour-diagonal **millicharges** and **charge radii**

● **Effective magnetic moment**

$|\mu_\nu(L, E_\nu)|^2 = \sum_{i=1}^3 \sum_{l, l'=1}^3 U_{\ell k}^* U_{\ell k'} (\mu_\nu)_{jk} (\mu_\nu)_{jk'}^* = \sum_{\ell'=e,\mu,\tau} |(\mu_\nu)_{\ell \ell'}|^2 \quad \text{where}$

$(\mu_\nu)_{\ell \ell} = \sum_{j,k=1}^3 U_{\ell k}^* U_{\ell' j} (\mu_\nu)_{jk}$

is the effective magnetic moment in flavor basis

● for GEMMA experiment ... ●

● Long-baseline case

$$L \gg L_{kj} = 2E_\nu / |\delta m_{kk'}^2|$$

$$\exp(-i\delta m_{kk'}^2 / 2E_\nu) = \delta_{kk'}$$

effect of decoherence

$$C_1 = g_V^2 + 2g_V P_{\nu_\ell \rightarrow \nu_e} + P_{\nu_\ell \rightarrow \nu_e} + \sum_{j,k=1}^3 |U_{\ell k}|^2 |\tilde{Q}_{jk}|^2 + 2g_V \sum_{j=1}^3 |U_{\ell j}|^2 \tilde{Q}_{jj} + 2 \sum_{j,k=1}^3 |U_{\ell k}|^2 \operatorname{Re} \left\{ U_{ej} U_{ek}^* \tilde{Q}_{jk} \right\}$$

$$C_2 = g_A^2 + 2g_A P_{\nu_\ell \rightarrow \nu_e} + P_{\nu_\ell \rightarrow \nu_e}$$

$$C_3 = g_V g_A + (g_V + g_A + 1) P_{\nu_\ell \rightarrow \nu_e} + g_A \sum_{j=1}^3 |U_{\ell j}|^2 \tilde{Q}_{jj} + 2 \sum_{j,k=1}^3 |U_{\ell k}|^2 U_{ej} U_{ek}^* \tilde{Q}_{jk}$$

where the flavour transition probability $P_{\nu_\ell \rightarrow \nu_e} = \sum_{k=1}^3 |U_{\ell k}|^2 |U_{ek}|^2$ does not depend on source-detector distance and ν energy

- Effective magnetic moment is independent of L and E

$$|\mu_\nu(L, E_\nu)|^2 = \sum_{j,k=1}^3 |U_{\ell k}|^2 |(\mu_\nu)_{jk}|^2$$

- for Borexino experiment ... ●

Experimental limits for different effective μ_ν

Method	Experiment	Limit	CL	Reference
Reactor $\bar{\nu}_e-e^-$	Krasnoyarsk	$\mu_{\nu_e} < 2.4 \times 10^{-10} \mu_B$	90%	Vidyakin <i>et al.</i> (1992)
	Rovno	$\mu_{\nu_e} < 1.9 \times 10^{-10} \mu_B$	95%	Derbin <i>et al.</i> (1993)
	MUNU	$\mu_{\nu_e} < 0.9 \times 10^{-10} \mu_B$	90%	Daraktchieva <i>et al.</i> (2005)
	TEXONO	$\mu_{\nu_e} < 7.4 \times 10^{-11} \mu_B$	90%	Wong <i>et al.</i> (2007)
	● GEMMA	$\mu_{\nu_e} < 2.9 \times 10^{-11} \mu_B$	90%	Beda <i>et al.</i> (2012)
Accelerator ν_e-e^-	LAMPF	$\mu_{\nu_e} < 10.8 \times 10^{-10} \mu_B$	90%	Allen <i>et al.</i> (1993)
Accelerator $(\nu_\mu, \bar{\nu}_\mu)-e^-$	BNL-E734	$\mu_{\nu_\mu} < 8.5 \times 10^{-10} \mu_B$	90%	Ahrens <i>et al.</i> (1990)
	LAMPF	$\mu_{\nu_\mu} < 7.4 \times 10^{-10} \mu_B$	90%	Allen <i>et al.</i> (1993)
	LSND	$\mu_{\nu_\mu} < 6.8 \times 10^{-10} \mu_B$	90%	Auerbach <i>et al.</i> (2001)
Accelerator $(\nu_\tau, \bar{\nu}_\tau)-e^-$	DONUT	$\mu_{\nu_\tau} < 3.9 \times 10^{-7} \mu_B$	90%	Schwienhorst <i>et al.</i> (2001)
Solar ν_e-e^-	Super-Kamiokande	$\mu_S(E_\nu \gtrsim 5 \text{ MeV}) < 1.1 \times 10^{-10} \mu_B$	90%	Liu <i>et al.</i> (2004)
	Borexino	$\mu_S(E_\nu \lesssim 1 \text{ MeV}) < 5.4 \times 10^{-11} \mu_B$	90%	Arpesella <i>et al.</i> (2008)

C. Giunti, A. Studenikin, Electromagnetic interactions of neutrinos:
A window to new physics, *Rev. Mod. Phys.* **87** (2015) 531

● new 2017 Borexino PRD: $\mu_\nu^{eff} < 2.8 \cdot 10^{-11} \mu_B$ at 90% c.l.

● Particle Data Group, 2014-2022 and 2024

Bounds on millicharge q_ν from μ_ν

2

(GEMMA Coll. data)

two not seen contributions:

ν - e cross-section

$$\left(\frac{d\sigma}{dT}\right)_{\nu-e} = \left(\frac{d\sigma}{dT}\right)_{SM} + \left(\frac{d\sigma}{dT}\right)_{\mu_\nu} + \left(\frac{d\sigma}{dT}\right)_{q_\nu}$$

$$\left(\frac{d\sigma}{dT}\right)_{\mu_\nu^a} \approx \pi\alpha^2 \frac{1}{m_e^2 T} \left(\frac{\mu_\nu^a}{\mu_B}\right)^2$$

$$\left(\frac{d\sigma}{dT}\right)_{q_\nu} \approx 2\pi\alpha \frac{1}{m_e T^2} q_\nu^2$$

Bounds on q_ν from ... unobserved effects of New Physics

$$R = \frac{\left(\frac{d\sigma}{dT}\right)_{q_\nu}}{\left(\frac{d\sigma}{dT}\right)_{\mu_\nu^a}} = \frac{2m_e}{T} \frac{\left(\frac{q_\nu}{e_0}\right)^2}{\left(\frac{\mu_\nu^a}{\mu_B}\right)^2} \ll 1$$



Studenikin, Europhys. Lett. 107 (2014) 210011

Particle Data Group, 2016-2024

Expected new constraints from GEMMA:

Constraints on q_ν

now $\mu_\nu < 2.9 \times 10^{-11} \mu_B$ ($T \sim 2.8$ keV)

● $|q_\nu| < 1.5 \times 10^{-12} e_0$

in ν Table of Particle Data Group since 2016

2023+ few years data taking

ν GeN experiment

... low threshold ...

$\mu_\nu \sim (5 - 9) \times 10^{-12} \mu_B$

$T \sim 200$ eV

● $|q_\nu| < 1.1 \times 10^{-13} e_0$

2

Experimental limits for different effective q_ν

=

C. Giunti, A. Studenikin, Electromagnetic interactions of neutrinos: a window to new physics, *Rev. Mod. Phys.* 87 (2015) 531

Limit	Method	Reference
$ q_{\nu_\tau} \lesssim 3 \times 10^{-4} e$	SLAC e^- beam dump	Davidson <i>et al.</i> (1991)
$ q_{\nu_\tau} \lesssim 4 \times 10^{-4} e$	BEBC beam dump	Babu <i>et al.</i> (1994)
$ q_\nu \lesssim 6 \times 10^{-14} e$	Solar cooling (plasmon decay)	Raffelt (1999a)
$ q_\nu \lesssim 2 \times 10^{-14} e$	Red giant cooling (plasmon decay)	Raffelt (1999a)
$ q_{\nu_e} \lesssim 3 \times 10^{-21} e$	● Neutrality of matter ●	Raffelt (1999a)
$ q_{\nu_e} \lesssim 3.7 \times 10^{-12} e$	Nuclear reactor	Gninenko <i>et al.</i> (2007)
$ q_{\nu_e} \lesssim 1.5 \times 10^{-12} e$	Nuclear reactor	Studenikin (2013)

A. Studenikin: New bounds on neutrino electric millicharge from limits on neutrino magnetic moment, *Eur.Phys.Lett.* 107 (2014) 2100

... since that C.Patrigiani et al (Particle Data Group), *The Review of Particle Physics 2016* Chinese Physics C 40 (2016) 100001



charge radii

... most accessible for experimental studies are charge radii $\langle r_{\nu}^2 \rangle$

Bernabeu, Papavassiliou, Vidal, 2004

... astrophysical bounds ???

... comprehensive analysis of ν - e scattering ...

PHYSICAL REVIEW D **95**, 055013 (2017)

Electromagnetic properties of massive neutrinos in low-energy elastic neutrino-electron scattering

Konstantin A. Kouzakov*

*Department of Nuclear Physics and Quantum Theory of Collisions, Faculty of Physics,
Lomonosov Moscow State University, Moscow 119991, Russia*

Alexander I. Studenikin[†]

*Department of Theoretical Physics, Faculty of Physics, Lomonosov Moscow State University,
Moscow 119991, Russia*

and Joint Institute for Nuclear Research, Dubna 141980, Moscow Region, Russia

(Received 11 February 2017; published 14 March 2017)

A thorough account of electromagnetic interactions of massive neutrinos in the theoretical formulation of low-energy elastic neutrino-electron scattering is given. The formalism of neutrino charge, magnetic, electric, and anapole form factors defined as matrices in the mass basis is employed under the assumption of three-neutrino mixing. The flavor change of neutrinos traveling from the source to the detector is taken into account and the role of the source-detector distance is inspected. The effects of neutrino flavor-transition millicharges and charge radii in the scattering experiments are pointed out.

DOI: 10.1103/PhysRevD.95.055013

... all experimental constraints on charge radius should be redone

Concluding remarks

Kouzakov, Studenikin

Phys. Rev. D 95 (2017) 055013

- cross section of ν - e is determined in terms of 3×3 matrices of ν electromagnetic form factors
- in **short-baseline** experiments one studies form factors in **flavour basis**
- **long-baseline** experiments more convenient to interpret in terms of fundamental form factors in **mass basis**
- ν millicharge when it is constrained in reactor short-baseline experiments (GEMMA, for instance) should be interpreted as

$$|e_{\nu e}| = \sqrt{|(e_{\nu})_{ee}|^2 + |(e_{\nu})_{\mu e}|^2 + |(e_{\nu})_{\tau e}|^2}$$

- ν charge radius in ν - e elastic scattering can't be considered as a shift $g_V \rightarrow g_V + \frac{2}{3} M_W^2 \langle r^2 \rangle \sin^2 \theta_W$, there are also contributions from flavor-transition charge radii

Generalized ν charge

Up to now we have used $\tilde{Q}_{jk} = \frac{2\sqrt{2}\pi\alpha}{G_F} \left[\frac{(e_\nu)_{jk}}{q^2} + \frac{1}{6} \langle r_\nu^2 \rangle_{jk} \right]$ in mass basis

Finally we have in flavour basis

$$\tilde{Q}_{\ell'\ell} = \sum_{j,k=1}^3 U_{\ell'j} U_{\ell k}^* \tilde{Q}_{jk} = \frac{2\sqrt{2}\pi\alpha}{G_F} \left[\frac{(e_\nu)_{\ell'\ell}}{q^2} + \frac{1}{6} \langle r_\nu^2 \rangle_{\ell'\ell} \right]$$

where

$$(e_\nu)_{\ell'\ell} = \sum_{j,k=1}^3 U_{\ell'j} U_{\ell k}^* (e_\nu)_{jk}$$

millicharge

in ν flavour basis

$$\langle r_\nu^2 \rangle_{\ell'\ell} = \sum_{j,k=1}^3 U_{\ell'j} U_{\ell k}^* \langle r_\nu^2 \rangle_{jk}$$

charge radius

Ch - It - Ru
collaboration

Neutrino charge radii from COHERENT elastic neutrino-nucleus scattering

M. Cadeddu*

*Dipartimento di Fisica, Università degli Studi di Cagliari,
and INFN, Sezione di Cagliari, Complesso Universitario di Monserrato—S.P.
per Sestu Km 0.700, 09042 Monserrato (Cagliari), Italy*

C. Giunti†

Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy

K. A. Kouzakov‡

*Department of Nuclear Physics and Quantum Theory of Collisions, Faculty of Physics,
Lomonosov Moscow State University, Moscow 119991, Russia*

Y. F. Li§

*Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China
and School of Physical Sciences, University of Chinese Academy of Sciences,
Beijing 100049, China*

A. I. Studenikin¶

*Department of Theoretical Physics, Faculty of Physics,
Lomonosov Moscow State University, Moscow 119991, Russia
and Joint Institute for Nuclear Research, Dubna 141980, Moscow Region, Russia*

Y. Y. Zhang¶

*Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China
and School of Physical Sciences, University of Chinese Academy of Sciences,
Beijing 100049, China* (Received 15 October 2018; published 26 December 2018)

Coherent elastic neutrino-nucleus scattering is a powerful probe of neutrino properties, in particular of the neutrino charge radii. We present the bounds on the neutrino charge radii obtained from the analysis of the data of the COHERENT experiment. We show that the time information of the COHERENT data allows us to restrict the allowed ranges of the neutrino charge radii, especially that of ν_μ . We also obtained for the first time bounds on the neutrino transition charge radii, which are quantities beyond the standard model.

DOI: 10.1103/PhysRevD.98.113010

$$(|\langle r_{\nu e\mu}^2 \rangle|, |\langle r_{\nu e\tau}^2 \rangle|, |\langle r_{\nu \mu\tau}^2 \rangle|) < (22, 38, 27) \times 10^{-32} \text{ cm}^2$$

K. Kouzakov, A. Studenikin, “Electromagnetic properties of massive neutrinos in low-energy elastic neutrino-electron scattering”
Phys. Rev. D 95 (2017) 055013

Physical Review D
– Highlights 2018 –
Editors' Suggestion

“Using data from the COHERENT experiment, the authors put bounds on electromagnetic charge radii, including the first bounds on transition charge radii. These results show promising prospects for current and upcoming ν -nucleus experiments”

Physical Review D – Highlights 2018 – Editors' Suggestion


29.12.2018

Physical Review D - Highlights

Editors' Suggestion

Neutrino charge radii from COHERENT elastic neutrino-nucleus scattering (/prd/abstract/10.1103/PhysRevD.98.113010)

M. Cadeddu, C. Giunti, K. A. Kouzakov, Y. F. Li, A. I. Studenikin, and Y. Y. Zhang
Phys. Rev. D **98**, 113010 (2018) – Published 26 December 2018



coherent ν scattering
due to charge radius

Using data from the COHERENT experiment, the authors put bounds on neutrino electromagnetic charge radii, including the first bounds on the transition charge radii. These results show promising prospects for current and upcoming neutrino-nucleus scattering experiments.

[Show Abstract +\(\)](#)

Particle Data Group,
Review of Particle Physics (2018-2024)

Experimental limits on ν charge radius $\langle r_\nu^2 \rangle$ =

C. Giunti, A. Studenikin, “Electromagnetic interactions of neutrinos: a window to new physics”, *Rev. Mod. Phys.* **87** (2015) 531

Method	Experiment	Limit (cm ²)	C.L.	Reference
Reactor $\bar{\nu}_e$ - e^-	Krasnoyarsk	$ \langle r_{\nu_e}^2 \rangle < 7.3 \times 10^{-32}$	90%	Vidyakin <i>et al.</i> (1992)
	TEXONO	$-4.2 \times 10^{-32} < \langle r_{\nu_e}^2 \rangle < 6.6 \times 10^{-32}$	90%	Deniz <i>et al.</i> (2010) ^a
Accelerator ν_e - e^-	LAMPF	$-7.12 \times 10^{-32} < \langle r_{\nu_e}^2 \rangle < 10.88 \times 10^{-32}$	90%	Allen <i>et al.</i> (1993) ^a
	LSND	$-5.94 \times 10^{-32} < \langle r_{\nu_e}^2 \rangle < 8.28 \times 10^{-32}$	90%	Auerbach <i>et al.</i> (2001) ^a
Accelerator ν_μ - e^-	BNL-E734	$-4.22 \times 10^{-32} < \langle r_{\nu_\mu}^2 \rangle < 0.48 \times 10^{-32}$	90%	Ahrens <i>et al.</i> (1990) ^a
	CHARM-II	$ \langle r_{\nu_\mu}^2 \rangle < 1.2 \times 10^{-32}$	90%	Vilain <i>et al.</i> (1995) ^a





... updated by the recent constraints
(effects of physics **Beyond Standard Model**)

$$(|\langle r_{\nu_{e\mu}}^2 \rangle|, |\langle r_{\nu_{e\tau}}^2 \rangle|, |\langle r_{\nu_{\mu\tau}}^2 \rangle| < (28, 30, 35) \times 10^{-32} \text{ cm}^2$$

M.Cadeddu, C. Giunti, K.Kouzakov,
Yu-Feng Li, A. Studenikin, Y.Y.Zhang,
Neutrino charge radii from COHERENT elastic neutrino-nucleus
scattering, *Phys.Rev.D* **98** (2018) 113010

Electromagnetic ν in
astrophysics and
bounds on μ_ν and q_ν


ν electromagnetic interactions



Neutrino radiative decay $\nu_i \rightarrow \nu_j + \gamma$

$$\mathcal{L}_{int} = \frac{1}{2} \bar{\psi}_i \sigma_{\mu\nu} (\mu_{if} + i\gamma_5 d_{if}) \psi_j F^{\mu\nu} + h.c.$$

$m_i > m_j$



$$\Lambda_{\mu}^{if}(q) = -i\sigma_{\mu\nu} q^{\nu} (\mu_{if} + i\gamma_5 d_{if})$$

Petkov 1977; Zatsepin, Smirnov 1978;
Bilenky, Petkov 1987; Pal, Wolfenstein 1982

Radiative decay rate

$$\Gamma_{\nu_i \rightarrow \nu_j + \gamma} = \frac{\mu_{eff}^2}{8\pi} \left(\frac{m_i^2 - m_j^2}{m_i^2} \right)^3 \approx 5 \left(\frac{\mu_{eff}}{\mu_B} \right)^2 \left(\frac{m_i^2 - m_j^2}{m_i^2} \right)^3 \left(\frac{m_i}{1 \text{ eV}} \right)^3 \text{ s}^{-1}$$

$$\mu_{eff}^2 = |\mu_{ij}|^2 + |\epsilon_{ij}|^2$$

$$\tau_{\nu_i \rightarrow \nu_j + \gamma}^{rf} \approx 0.19 \left(\frac{m_i^2}{m_i^2 - m_j^2} \right)^3 \left(\frac{\text{eV}}{m_i} \right)^3 \left(\frac{\mu_B}{\mu_{fi}^{eff}} \right)^2 \text{ s}$$

transition magnetic and electric moments
(for Dirac and Majorana ν)

• ν life time is indeed huge ...

Radiative decay has been constrained from absence of decay photons: ●

- 1) reactor $\bar{\nu}_e$ and solar ν_e fluxes,
- 2) SN 1987A ν burst (all flavours)
- 3) spectral distortion of CMBR

Raffelt 1999
Kolb, Turner 1990
Ressell, Turner 1990

Neutrino Cherenkov radiation $\nu_L(p) \rightarrow \nu_R(p') + \gamma(k)$

helicity flip process

ν transition amplitude due to μ_ν

$$M = \frac{\mu}{n} \overline{u^{(+)}(p')} \sigma_{\mu\nu} k^\mu u^{(-)}(p) \epsilon^\nu(k, \lambda)$$

Cherenkov process rate

$$\Gamma = \frac{1}{2(2\pi)^2 E} \int \frac{d^3 p'}{2E'} \frac{d^3 k}{2\omega} |M|^2 \delta^4(p - p' - k)$$

after integration


$$\Gamma = \frac{1}{16\pi E^2 v} \int n^2 d\omega d(\cos \theta) |M|^2 \delta\left(\cos \theta - \frac{2\omega E + (n^2 - 1)\omega^2}{2n\omega E v}\right)$$

photon emission angle

$$|\cos \theta| \leq 1$$

$$\cos \theta = \frac{1}{nv} \left(1 + (n^2 - 1) \frac{\omega}{2E} \right)$$

$$\Gamma = \frac{\mu^2}{4\pi E^2 v} \int_{\omega_{\min}}^{\omega_{\max}} \left\{ \left[\frac{(n^2 - 1)^2}{n^2} E^2 + (n^2 - 1) m_\nu^2 \right] \omega^2 - \frac{(n^2 - 1)^2}{n^2} E \omega^3 - \frac{(n^2 - 1)^3}{4n^2} \omega^4 \right\} d\omega$$



Cherenkov radiation

$$v = |\vec{p}|/E$$

Solar ν s with $\mu_\nu \sim 3 \times 10^{-11} \mu_B$ emit 5 γ per day in 1 Km³ water detector ●


Grimus & Neufeld, 1993

ν radiative decay and Cherenkov radiation in external environments

coherent forward elastic scattering on (electron) background also

generates $\nu_i \rightarrow \nu_j + \gamma$
 not suppressed by GIM

D'Olive, Nieves, Pal (1990)



Giunti et al (1992)

Cherenkov radiation by ν in magnetic field

Galtsov, Nikitina (1972)

B induces effective ν - γ vertex and modifies γ dispersion relation

Ioannian & Raffelt (1997)

(no need for BSM)

ν in medium acquire induce q as a consequence of weak interactions

Oraevsky, Semikoz, Smorodinsky (1986)

another mechanism of Cherenkov radiation in medium

Sawyer (1992)

D'Olive, Nieves, Pal (1996)

effect for $m_\nu = 0$ in SM (without physics BSM)

- other particular cases for $\nu_i \rightarrow \nu_j + \gamma$
 in em fields and matter

Skobelev (1976)


Borisov, Zhukosky, Ternov (1988)

Ternov (2016)



● New mechanism of electromagnetic radiation

"Spin light of neutrino"
in matter and
electromagnetic fields



- ... quasi-classical approach to ν spin evolution in external field

A. Egorov, A. Lobanov, A. Studenikin,
Phys.Lett. B 491 (2000) 137
Lobanov, Studenikin,
● Phys.Lett. B 515 (2001) 94
Phys.Lett. B 564 (2003) 27
Phys.Lett. B 601 (2004) 171
Studenikin, A.Ternov,
Phys.Lett. B 608 (2005) 107
A. Grigoriev, Studenikin, Ternov,
Phys.Lett. B 622 (2005) 199
Studenikin,
J.Phys.A: Math.Gen. 39 (2006) 6769
J.Phys.A: Math.Theor. 41 (2008) 16402
Grigoriev, A. Lokhov, Studenikin, Ternov,
Nuovo Cim. 35 C (2012) 57
Phys.Lett.B 718 (2012) 512

New mechanism of electromagnetic radiation

?

Why Spin Light

of neutrino

$SL\nu$

of electron

SLe

in matter

Analogies with:


* classical electrodynamics

an object with charge $Q=0$ and

magnetic moment $\vec{m} = \frac{1}{2} \sum_i e_i [\vec{r}_i \times \vec{v}_i] \neq 0$

$$\overset{\text{cl. el.}}{I} = \frac{2}{3} \ddot{\vec{m}}^2$$

← magnetic dipole radiation power



$$\frac{d\vec{S}}{dt} = 2\mu_B [\vec{S} \times \vec{B}] + 2\mu_B [\vec{S} \times \vec{G}]$$

electromagnetic
interaction with
e.m. field

weak interaction
with matter



spin evolution in presence of general external fields



M.Dvornikov, A.Studenikin,
JHEP 09 (2002) 016

General types non-derivative interaction with external fields

$$\begin{aligned}
-\mathcal{L} = & g_s s(x) \bar{\nu} \nu + g_p \pi(x) \bar{\nu} \gamma^5 \nu + g_v V^\mu(x) \bar{\nu} \gamma_\mu \nu + g_a A^\mu(x) \bar{\nu} \gamma_\mu \gamma^5 \nu + \\
& + \frac{g_t}{2} T^{\mu\nu} \bar{\nu} \sigma_{\mu\nu} \nu + \frac{g'_t}{2} \Pi^{\mu\nu} \bar{\nu} \sigma_{\mu\nu} \gamma^5 \nu,
\end{aligned}$$

scalar, pseudoscalar, vector, axial-vector, $s, \pi, V^\mu = (V^0, \vec{V}), A^\mu = (A^0, \vec{A}),$
 tensor and pseudotensor fields: $T_{\mu\nu} = (\vec{a}, \vec{b}), \Pi_{\mu\nu} = (\vec{c}, \vec{d})$

Relativistic equation (quasiclassical) for spin vector:

$$\begin{aligned}
\dot{\vec{\zeta}}_\nu = & 2g_a \left\{ A^0 [\vec{\zeta}_\nu \times \vec{\beta}] - \frac{m_\nu}{E_\nu} [\vec{\zeta}_\nu \times \vec{A}] - \frac{E_\nu}{E_\nu + m_\nu} (\vec{A} \vec{\beta}) [\vec{\zeta}_\nu \times \vec{\beta}] \right\} \\
& + 2g_t \left\{ [\vec{\zeta}_\nu \times \vec{b}] - \frac{E_\nu}{E_\nu + m_\nu} (\vec{\beta} \vec{b}) [\vec{\zeta}_\nu \times \vec{\beta}] + [\vec{\zeta}_\nu \times [\vec{a} \times \vec{\beta}]] \right\} + \\
& + 2ig'_t \left\{ [\vec{\zeta}_\nu \times \vec{c}] - \frac{E_\nu}{E_\nu + m_\nu} (\vec{\beta} \vec{c}) [\vec{\zeta}_\nu \times \vec{\beta}] - [\vec{\zeta}_\nu \times [\vec{d} \times \vec{\beta}]] \right\}.
\end{aligned}$$

● Neither S nor π nor V contributes to spin evolution

● Electromagnetic interaction

$$T_{\mu\nu} = F_{\mu\nu} = (\vec{E}, \vec{B})$$

● SM weak interaction


$$G_{\mu\nu} = (-\vec{P}, \vec{M}) \quad \begin{aligned} \vec{M} &= \gamma(A^0 \vec{\beta} - \vec{A}) \\ \vec{P} &= -\gamma[\vec{\beta} \times \vec{A}], \end{aligned}$$



... quantum theory of



Spin light of neutrino in matter



new mechanism of the electromagnetic process stimulated by the presence of matter, in which neutrino with **nonzero magnetic moment** emits light

A.Lobanov, A.Studenikin, Phys.Lett. B 564 (2003) 27,
Phys.Lett. B 601 (2004) 171

A.S., A.Ternov, Phys.Lett. B 608 (2005) 107

A.Grigoriev, A.S., A.Ternov, Phys.Lett. B 622 (2005) 199

A.S., J.Phys.A: Math.Theor. 41 (2008) 16402

A.S., J.Phys.A: Math.Gen. 39 (2006) 6769

A.Grigoriev, A.Lokhov, A.Ternov, A.Studenikin,
Phys. Lett. B 718 (2012) 512
JCAP 11 (2017) 024

«method of exact solutions»

Interaction of particles in external electromagnetic fields

(**Furry representation** in quantum electrodynamics)

Potential of electromagnetic field

$$A_\mu(x) = A_\mu^q(x) + A_\mu^{ext}(x)$$

quantized part
of potential

evolution operator

$$U_F(t_1, t_2) = Texp \left[-i \int_{t_1}^{t_2} j^\mu(x) A_\mu^q(x) dx \right]$$


charged particles current

$$j_\mu(x) = \frac{e}{2} [\bar{\Psi}_F \gamma_\mu, \Psi_F]$$

Dirac equation in external classical (nonquantized) field $A_\mu^{ext}(x)$

$$\left\{ \gamma^\mu \left(i\partial_\mu - eA_\mu^{ext}(x) \right) - m_e \right\} \Psi_F(x) = 0$$

- ...beyond perturbation series expansion, strong fields and non linear effects...



✓ in external
electromagnetic
fields

✓ in
dense
matter

...«method of exact solutions»...

Studenikin

- Quantum treatment of neutrino in background matter
J. Phys. A: Math. Gen. 39 (2006) 6769–6776
- Method of wave equations exact solutions in studies
of neutrinos and electron interactions in dense matter
J. Phys. A: Math. Theor. 41 (2008) 164047 (20 p)
- Neutrinos and electrons in background matter: A new
approach
Ann. Fond. de Broglie 31 (2006) 289-316

- ν quantum states in dense magnetized matter

... new effect of ...

Spin Light of ν
in matter



... phenomenological


ν energy quantization in rotating matter

consequences in astrophysics (pulsars)

ν in matter treated within
«method of exact solutions»

(Dirac equation with matter potential for ν)

Neutrino – photon coupling



broad neutrino lines
account for interaction
with environment

“Spin light of neutrino in matter”

$SL\nu$

- ... within the quantum treatment based on
method of exact solutions ...



Modified Dirac equation for neutrino in matter


Addition to the vacuum neutrino Lagrangian

$$\Delta L_{eff} = \Delta L_{eff}^{CC} + \Delta L_{eff}^{NC} = -f^\mu \left(\bar{\nu} \gamma_\mu \frac{1 + \gamma_5}{2} \nu \right)$$

matter
current

where $f^\mu = \frac{G_F}{\sqrt{2}} \left((1 + 4 \sin^2 \theta_W) j^\mu - \lambda^\mu \right)$

matter
polarization



$$\left\{ i \gamma_\mu \partial^\mu - \frac{1}{2} \gamma_\mu (1 + \gamma_5) f^\mu - m \right\} \Psi(x) = 0$$

A.Studenikin, A.Ternov, hep-ph/0410297;
Phys.Lett.B 608 (2005) 107

It is supposed that there is a macroscopic amount of electrons in the scale of a neutrino de Broglie wave length. Therefore, **the interaction of a neutrino with the matter (electrons) is coherent.**

L.Chang, R.Zia,'88; J.Panteleone,'91; K.Kiers, N.Weiss, M.Tytgat,'97-'98; P.Manheim,'88; D.Nötzold, G.Raffelt,'88; J.Nieves,'89; V.Oraevsky, V.Semikoz, Ya.Smorodinsky,89; W.Naxton, W-M.Zhang'91; M.Kachelriess,'98; A.Kusenko, M.Postma,'02.

This is the most general equation of motion of a neutrino in which the effective potential accounts for both the **charged** and **neutral-current** interactions with the background matter and also for the possible effects of the matter **motion** and **polarization.**

Quantum theory of spin light of neutrino




Quantum treatment of *spin light of neutrino* in matter shows that this process originates from the two subdivided phenomena:

the shift of the neutrino energy levels in the presence of the background matter, which is different for the two opposite **neutrino helicity states**,

$$E = \sqrt{\mathbf{p}^2 \left(1 - s\alpha \frac{m}{p}\right)^2 + m^2} + \alpha m$$

$$s = \pm 1$$



the radiation of the photon in the process of the neutrino transition from the **“excited” helicity state** to the **low-lying helicity state** in matter

A.Studenikin, A.Ternov, Phys.Lett.B 608 (2005) 107;

A.Grigoriev, A.Studenikin, A.Ternov, Phys.Lett.B 622 (2005) 199;
Grav. & Cosm. 14 (2005) 132;

neutrino-spin self-polarization effect in the matter



A.Lobanov, A.Studenikin, Phys.Lett.B 564 (2003) 27;
Phys.Lett.B 601 (2004) 171

≡ Spatial distribution of radiation power

From the angular distribution of


$$SL\nu$$

$$I = \mu^2 \int_0^\pi \omega^4 [(\tilde{\beta}\tilde{\beta}' + 1)(1 - y \cos \theta) - (\tilde{\beta} + \tilde{\beta}')(\cos \theta - y)] \frac{\sin \theta}{1 + \tilde{\beta}'y} d\theta$$



for $p/m = 5$ and $\alpha = 0.01$

$$n \approx 10^{35} \text{ cm}^{-3}$$



$$\cos \theta_{max} \approx 1 - \frac{2}{3} \alpha \frac{m}{p}$$

maximum in radiation power distribution

for $p/m = 10^3$ and $\alpha = 100$

$$n \approx 10^{39} \text{ cm}^{-3}$$

increase of matter density

projector-like distribution



cap-like distribution

It is possible to have

$$\tau = \frac{1}{\Gamma_{SL\nu}} \ll \text{age of the Universe ?}$$

For ultra-relativistic \checkmark

with momentum $p \sim 10^{20} \text{ eV}$

and magnetic moment $\mu \sim 10^{-10} \mu_B$

in very dense matter $n \sim 10^{40} \text{ cm}^{-3}$

from

$$\Gamma_{SL\nu} = 4\mu^2 \alpha^2 m_\nu^2 p$$

$$p \gg m_{\text{plasmon}}$$

also discussed by
A.Kuznetsov,
N.Mikheev,
IJMP A 2007

$$\alpha m_\nu = \frac{1}{2\sqrt{2}} G_F n (1 + \sin^2 \theta_W)$$

A.Lobanov, A.S., PLB 2003; PLB 2004

A.Grigoriev, A.S., PLB 2005

A.Grigoriev, A.S., A.Ternov, PLB 2005

A.Grigories, A.Lokhov, A.S., A.Ternov, PLB 2012

it follows that

$$\tau_{SL\nu} = \frac{1}{\Gamma} = 1.5 \times 10^{-8} \text{ s}$$

**A. Grigoriev, A. Lokhov,
A. Ternov, A. Studenikin**
**The effect of plasmon mass
on Spin Light of Neutrino
in dense matter**
**Phys. Lett. B 718
(2012) 512-515**




Figure 1: 3D representation of the radiation power distribution.




Figure 2: The two-dimensional cut along the symmetry axis. Relative units are used.

4. Conclusions

We developed a detailed evaluation of the spin light of neutrino in matter accounting for effects of the emitted plasmon mass. On the base of the exact solution of the modified Dirac equation for the neutrino wave function in the presence of the background matter the appearance of the threshold for the considered process is confirmed. The obtained exact and explicit threshold condition relation exhibit a rather complicated dependence on the matter density and neutrino mass. The dependence of the rate and power on the neutrino energy, matter density and the angular distribution of the $SL\nu$ is investigated in details. It is shown how the rate and power wash out when the threshold parameter $a = m_\gamma^2/4\tilde{n}p$ approaching unity. From the performed detailed analysis it is shown that the $SL\nu$ mechanism is practically insensitive to the emitted plasmon mass for very high densities of matter (even up to $n = 10^{41} \text{ cm}^{-3}$) for ultra-high energy neutrinos for a wide range of energies starting from $E = 1 \text{ TeV}$. This conclusion is of interest for astrophysical applications of $SL\nu$ radiation mechanism in light of the recently reported hints of $1 \div 10 \text{ PeV}$ neutrinos observed by IceCube [17].

Spin light of neutrino in astrophysical environments

**Alexander Grigoriev,^{b,c} Alexey Lokhov,^d Alexander Studenikin^{a,e,1}
and Alexei Ternov^c**

^aDepartment of Theoretical Physics, Moscow State University,
119992 Moscow, Russia

^bSkobeltsyn Institute of Nuclear Physics, Moscow State University,
119992 Moscow, Russia

^cDepartment of Theoretical Physics, Moscow Institute of Physics and Technology,
141701 Dolgoprudny, Russia

^dInstitute for Nuclear Research, Russian Academy of Sciences,
117312 Moscow, Russia

^eDzhelepov Laboratory of Nuclear Problems, Joint Institute for Nuclear Research,
141980 Dubna, Russia

E-mail: ax.grigoriev@mail.ru, lokhov.alex@gmail.com, studenik@srd.sinp.msu.ru,
ternov.ai@mipt.ru

Received May 23, 2017

Revised October 16, 2017

Accepted October 31, 2017

Published November 16, 2017

A.Grigoriev, A.Lokhov, A.Studenikin, A.Ternov, Spin light of neutrino in astrophysical environments. *J. Cosm. Astropart. Phys.* 11 (2017) 024

SLν in neutron matter of real astrophysical objects [4]

□ Plasma effects [5]

- Photon dispersion with plasmon mass in the degenerate electron gas:

$$\omega = \sqrt{k^2 + m_\gamma^2}$$

$$m_\gamma = \left(\frac{2\alpha}{\pi}\right)^{1/2} \mu_e \simeq 8.87 \times \left(\frac{n_e}{10^{37} \text{ cm}^{-3}}\right)^{1/3} \text{ MeV}$$

- Threshold condition for the SLν [10]: ($Y_e = n_\nu/n_n$)

$$\frac{m_\gamma^2 + 2 m_\gamma m_\nu}{4 \tilde{n} p} < 1$$

- **Neutron matter:** (antineutrinos act)

$$\tilde{n} = \frac{1}{2\sqrt{2}} G_F n_n \simeq 3.2 \times \left(\frac{n_n}{10^{38} \text{ cm}^{-3}}\right) \text{ eV,}$$

$$E > p_{th} \simeq 28.5 \times \frac{Y_e^{2/3}}{1 - Y_e} \left(\frac{10^{38} \text{ cm}^{-3}}{n_n}\right)^{1/3} \text{ TeV} \Rightarrow E_{th} \simeq 6.82 \text{ TeV.}$$

$$n_n = 10^{38} \text{ cm}^{-3}, \quad Y_e = 0.1$$

- Mean photon energy near the threshold: $\langle \omega \rangle = I/\Gamma \simeq p \simeq E_\nu$.

For most favorable conditions as low density of the charged matter component is needed as possible

□ W boson production $\bar{\nu}_e + e^- \rightarrow W^-$ [4]




Figure 2. The allowed range of electron antineutrino energies for the SLν in the matter of a neutron star depending on the neutron density. Solid line: the SLν process threshold without account for the $\bar{\nu}_e e$ -scattering; dash-dotted line: the SLν process threshold with account for the $\bar{\nu}_e e$ -scattering; dashed line: the threshold for the W boson production. (a) $Y_e = 0.09$; (b) $Y_e = 0.05$. The allowed regions are marked in green.

W-boson threshold energy $\epsilon_W = \frac{m_W^2}{4\mu_e} \simeq 5.77 \times \left(\frac{10^{38} \text{ cm}^{-3}}{Y_e n_n}\right)^{1/3} \text{ TeV}$

- Electron antineutrinos: s-channel interaction with matter through W-boson, importance of the propagator effects \Rightarrow correction to the effective potential of neutrino motion \rightarrow antineutrino energy shift up \rightarrow SLν is suppressed at $Y_e=0.1$, but allowed already for $Y_e=0.09$
- μ and τ antineutrinos: only t-channel interaction with matter through Z-boson, no propagator effects \Rightarrow the SLν is allowed if neutrino energy is greater than the W-boson threshold ϵ_W

Neutrino lifetime with respect to the SLν for most optimistic set of parameters:

$$\tau_{SL\nu} = 10^{-4} - 10^3 \text{ s, for } n_b = 10^{41} - 10^{38} \text{ cm}^{-3}$$

The SLν in short Gamma-Ray Bursts (SGRBs)

Factors for best SLν generation efficiency

- High neutrino energy and density
- High background neutral matter density
- Low density of the matter charged component
- Low temperature of the charged component
- Considerable extension of the medium

SLν radiation by ultra high-energy neutrino in the diffuse neutrino wind blown during neutron stars merger

Perego et al,
Mon.Not.Roy.Astron.Soc.
443 (2014) 3134



Matter characteristics[6]:

- neutrinos $n_\nu \sim 10^{32} \text{ cm}^{-3}$
 - electrons $Y_e = 0.01$
 - $T = 0.1 \text{ MeV}$
 - $\rho = 5 \times 10^3 \text{ g/cm}^3$
- $\Rightarrow n_e \simeq 3 \times 10^{25} \text{ cm}^{-3}$
 $m_\gamma \simeq 10^{-3} \text{ MeV}$
 $E_{th} \simeq 1 \text{ GeV}$

Radiation time

$$\tau_{SL\nu} \simeq 5.4 \times 10^{15} \left(\frac{10^{-11} \mu_B}{\mu}\right)^2 \left(\frac{10^{32} \text{ cm}^{-3}}{n_\nu}\right)^2 \left(\frac{1 \text{ PeV}}{E_\nu}\right) \text{ s}$$

Neutrino parameters:


$$\mu \simeq 2.9 \times 10^{-11} \mu_B$$

$$E_\nu \sim 10^{12} - 10^{18} \text{ eV}$$

$$\tau_{SL\nu} \simeq 6.4 \times (10^{11} - 10^{17}) \text{ s} = 2 \times (10^4 - 10^{10}) \text{ years}$$



Whiteboard 73643849-3731-2844



Neutrino spin operator and dispersion in moving matter



Alexander Grigoriev^{1,a}, Alexander Studenikin^{2,3,4,b}, Alexei Ternov^{1,c}

¹ Department of Theoretical Physics, Moscow Institute for Physics and Technology, 141700 Dolgoprudny, Russia
² Department of Physics, Moscow State University, 119992 Moscow, Russia
³ P.N. Lebedev Physical Institute of Russian Academy of Sciences, 125080 Moscow, Russia
⁴ Laboratory of Nuclear Problems, Joint Institute for Nuclear Research, 141980 Dubna, Russia
 Interdisciplinary Scientific and Educational School of Moscow University "Fundamental and Applied Physics Research", Moscow State University,

$\omega_{SL\nu}$ depends on $(\vec{V}_{matt} \vec{P}_\nu)$

Abstract We found the spin integral of motion for neutrinos propagating in moving and polarized matter. Contrary to all previous studies this is the exact spin operator commuting with the Hamiltonian for a neutrino in matter which moves in an arbitrary direction relative to the direction of neutrino propagation. The operator obtained opens up the possibility of consistent classification of neutrino states in such a medium and, as a consequence, a systematic description of the related physical phenomena. Using the operator, we obtain a dispersion relation for neutrinos in arbitrary moving matter and consider its particular cases.

$$\left\{ i\gamma_\mu \partial^\mu - \frac{1}{2}\gamma_\mu(1 + \gamma^5)f^\mu - m \right\} \Psi(x) = 0$$

$$\frac{1}{2} f^\mu = \tilde{n}_0 v^\mu, \quad \tilde{n}_0 = \frac{1}{2\sqrt{2}} G_F (1 + 4 \sin^2 \theta_W) n_0$$

$$n = \gamma n_0 \quad \gamma = 1/\sqrt{1 - v^2}$$

$$S = \gamma \left[\gamma^5 \gamma^0 m - \gamma^5 (\tilde{H} - (\tilde{p}v)) - m \gamma^0 (\Sigma v) \right]$$

$$E_{s=+1} = \sqrt{(p - \tilde{n})^2 + m^2} + \tilde{n},$$

$$E_{s=-1} = \sqrt{p^2 + 4\tilde{n}^2 v^2 + m^2} + 2\tilde{n}.$$

● Flavour oscillations $\nu_e \leftrightarrow \nu_\mu$ in moving matter

Wofenstein term

$$A \rightarrow A' = \sqrt{2} G_F n + 8 \frac{G_F^2 n^2 v^2}{p} \sin^2 \theta_W$$

Probability of oscillations

$$P = \frac{1}{2} \frac{\Delta^2 \sin^2 2\theta}{(\Delta \cos 2\theta - 2pA')^2 + \Delta^2 \sin^2 2\theta}$$

Resonance condition

$$\frac{\Delta}{2p} \cos 2\theta = A + 8 \frac{G_F^2 n^2 v^2}{p} \sin^2 \theta_W$$

two terms are of same order for


extra-dense matter $n \sim 10^{41} \text{ cm}^{-3}$ and $v \sim 0.9, P \sim 10 \text{ keV}$

Electromagnetic ν in
astrophysics and
bounds on μ_ν and g_ν

Astrophysical bounds on μ_ν

2 Astrophysical bound on μ_ν G.Raffelt, PRL 1990

comes from cooling (observed luminosity) of red giant stars by plasmon decay $\gamma^* \rightarrow \nu\nu$



$$L_{int} = \frac{1}{2} \sum_{a,b} \left(\mu_{a,b} \bar{\psi}_a \sigma_{\mu\nu} \psi_b + \epsilon_{a,b} \bar{\psi}_a \sigma_{\mu\nu} \gamma_5 \psi_b \right)$$

J.Bernstein et al 1963 ν : $\mu_\nu \sim 10^{-10} \mu_B$ solar

Matrix element

$$|M|^2 = M_{\alpha\beta} p^\alpha p^\beta, \quad M_{\alpha\beta} = 4\mu^2 (2k_\alpha k_\beta - 2k^2 \epsilon_\alpha^* \epsilon_\beta - k^2 g_{\alpha,\beta}), \quad \epsilon_\alpha k^\alpha = 0$$

Decay rate

$$\Gamma_{\gamma \rightarrow \nu\bar{\nu}} = \frac{\mu^2 (\omega^2 - k^2)^2}{24\pi \omega} = 0 \text{ in vacuum } \omega = k$$

In the classical limit γ^* - like a massive particle with $\omega^2 - k^2 = \omega_{pl}^2$

Energy-loss rate per unit volume

$$Q_\mu = g \int \frac{d^3k}{(2\pi)^3} \omega f_{BE} \Gamma_{\gamma \rightarrow \nu\bar{\nu}}$$

$$\mu^2 \rightarrow \sum_{a,b} (|\mu_{a,b}|^2 + |\epsilon_{a,b}|^2)$$


distribution function of plasmons

= Astrophysical bound on μ_ν

$$Q_\mu = g \int \frac{d^3k}{(2\pi)^3} \omega f_{BE} \Gamma_{\gamma \rightarrow \nu\bar{\nu}}$$

Magnetic moment **plasmon** decay enhances the Standard Model photo-neutrino cooling by photon polarization tensor

Energy-loss rate per unit volume



more fast star cooling

slightly reducing the core temperature

delay of helium ignition in low-mass red giants

(due to nonstandard ν losses)

astronomical observable

can be related to **luminosity** of stars before and after helium flash

... in order not to delay helium ignition in an unacceptable way
(a significant brightness increase is constraint by observations...)

... best
astrophysical
limit on ν
magnetic moment...

$$\mu \leq 3 \times 10^{-12} \mu_B$$

G.Raffelt, PRL 1990
D+M

$$\mu^2 \rightarrow \sum_{a,b} (|\mu_{a,b}|^2 + |\epsilon_{a,b}|^2)$$

... important for astrophysics consequence of

μ_ν is appearance ν_R

... examples 1-3 ...

1

a) helicity change in ν magnetic moment scattering on e (p, n)

(active) \Rightarrow (sterile)



effective μ_ν


$$\left(\frac{d\sigma}{dT}\right)_{\mu_\nu} = \frac{\pi\alpha_{em}^2}{m_e^2} \left[\frac{1 - T/E_\nu}{T}\right] \mu_\nu^2$$

$$\mu_\nu^2(\nu_l, L, E_\nu) = \sum_j \left| \sum_i U_{li} e^{-iE_i L} \mu_{ji} \right|^2$$

$\mu_{ij} \rightarrow |\mu_{ij} - i\epsilon_{ij}|$
electric dipole moment

b) spin (spin-flavor) precession in B_\perp

c) spin (spin-flavor) precession in transversal matter currents or polarization



2

... important for astrophysics consequence of $q_\nu \neq 0$

is ν deviation from a rectilinear trajectory

... example 4 ...

Astrophysics bounds on μ_ν

1) SN 1987A provides energy-loss limit on related to observed duration of ν signal

μ_ν

(also d_ν and transition moments)

...in magnetic moment scattering $\nu_e^L + e \rightarrow \nu_e^R + e$

due to change of helicity $\nu_L \Rightarrow \nu_R$

Dar, Nussinov & Rephaeli,
Goldman et al, Notzol, Voloshin,
Ayla et al, Balantekin et 1988

proto-neutron star formed in core-collapse SN can cool faster

since ν_R are sterile and not trapped in a core like ν_L for a few sec

- escaping ν_R will cool the core very efficient and fast (~ 1 s)

the observed 5-10 s pulse duration in Kamioka II and IMB

is in agreement with the standard model ν_L trapping ...

$$\mu_\nu^D \sim 10^{-12} \mu_B$$

... inconsistent with SN1987A observed cooling time

Barbieri, Mahapatra
Lattimer, Cooperstein,
1988
Raffelt, 1996

Astrophysics bounds on μ_ν

... example 5...

2) SN 1987A provides energy-loss limit on μ_ν related to observed ν energies

... helicity change in ν magnetic moment scattering $\nu_e^L + e \rightarrow \nu_e^R + e$
on e (p, n)

ν_R from inner SN core have larger energy than ν_L emitted from neutrino sphere

then $\nu_R \xleftrightarrow{B} \nu_L$ in galactic B and higher-energy ν_L would arrive to detector as a signal of SN 1987A

→ from absence of anomalous high-energy ν

Nötzold
1988

$$\mu_\nu^D \sim 10^{-12} \mu_B$$

Astrophysics bounds on μ_ν

$$\mu_\nu(\text{astro}) < 10^{-10} - 10^{-12} \mu_B$$

Mostly derived from consequences of helicity-state change in astrophysical medium:

- available degrees of freedom in BBN
- stellar cooling via plasmon decay
- cooling of SN1987a



Red Giant Lumin.
 $\mu_\nu \leq 3 \cdot 10^{-12} \mu_B$
G. Raffelt, D. Dearborn,
J. Silk, 1989.

Bounds depend on

- modeling of astrophysical system,
- on assumption on the neutrino properties.

Generic assumption:

- absence of other nonstandard interactions accept for μ_ν

A **global treatment** would be desirable, incorporating **oscillations** and **matter effects**, as well as the complications due to **interference** and **competitions** among **various channels**

Astrophysical bounds on q_ν

≡ Constraints on neutrino millicharge from red giants cooling

● Plasma process
(photon decay)


Interaction Lagrangian

Decay rate



$$L_{int} = -iq_\nu \bar{\psi}_\nu \gamma^\mu \psi_\nu A^\mu$$

$$\Gamma_{q_\nu} = \frac{q_\nu^2}{12\pi} \omega_{pl} \left(\frac{\omega_{pl}}{\omega} \right)$$



millicharge

Dobroliubov, Ignatiev 1990;
Babu, Volkas 1992;
Mohapatra, Nussinov 1992 ...

Delay of helium ignition in low-mass red giants due to nonstandard ν losses

$$q_\nu \leq 2 \times 10^{-14} e$$

...to avoid delay of helium ignition in low-mass red giants

Halt, Raffelt, Weiss, PRL1994

$$q_\nu \leq 3 \times 10^{-17} e$$

... absence of anomalous energy-dependent dispersion of SN1987A ν signal, most model independent

● $q_\nu \leq 3 \times 10^{-21} e$

... from "charge neutrality" of neutron...

- ... astrophysical bound on millicharge q_ν from

✓ energy quantization
in rotating
magnetized star

Grigoriev, Savochkin, Studenikin, Russ. Phys. J. 50 (2007) 845

Studenikin, J. Phys. A: Math. Theor. 41 (2008) 164047

Balantsev, Popov, Studenikin,

J. Phys. A: Math. Theor. 44 (2011) 255301

Balantsev, Studenikin, Tokarev, Phys. Part. Nucl. 43 (2012) 727

Phys. Atom. Nucl. 76 (2013) 489

- Studenikin, Tokarev, Nucl. Phys. B 884 (2014) 396

Millicharged ψ in rotating magnetized star

Balatsev, Tokarev, Studenikin,
Phys.Part.Nucl., 2012,

Phys.Atom.Nucl., Nucl.Phys. B, 2013,

- Studenikin, Tokarev, Nucl.Phys.B (2014)

Modified Dirac equation for ψ wave function

$$\left(\gamma_\mu (p^\mu + q_0 A^\mu) - \frac{1}{2} \gamma_\mu (c_l + \gamma_5) f^\mu - \frac{i}{2} \mu \sigma_{\mu\nu} F^{\mu\nu} - m \right) \Psi(x) = 0$$

external magnetic field

$$V_m = \frac{1}{2} \gamma_\mu (c_l + \gamma_5) f^\mu$$


matter potential

$$c_l = 1$$

rotating matter

$$f^\mu = -G n_n (1, -\epsilon y \omega, \epsilon x \omega, 0)$$

matter density



rotation
angular
frequency

- ✓ energy is quantized in rotating and magnetized star

A.Studenikin, I.Tokarev,
Nucl.Phys.B (2014)

$$G = \frac{G_F}{\sqrt{2}}$$

$$p_0 = \sqrt{p_3^2 + 2N|2Gn_n\omega - \epsilon q_\nu B| + m^2} - Gn_n - q\phi$$

$N = 0, 1, 2, \dots$
integer number

matter
rotation
frequency

millicharge

scalar potential
of electric field

- ! ✓ energy is quantized in rotating matter like electron energy in magnetic field (Landau energy levels):

- $$p_0^{(e)} = \sqrt{m_e^2 + p_3^2 + 2\gamma N}, \quad \gamma = eB, \quad N = 0, 1, 2, \dots$$

In quasi-classical approach

✓ quantum states in rotating matter
✓ motion in circular orbits

$$R = \int_0^\infty \Psi_L^\dagger r \Psi_L dr = \sqrt{\frac{2N}{|2Gn_n\omega - \epsilon q_0 B|}} \quad N=1,2,3 \dots$$

due to effective Lorentz force

$$\mathbf{F}_{eff} = q_{eff} \mathbf{E}_{eff} + q_{eff} [\boldsymbol{\beta} \times \mathbf{B}_{eff}]$$

A. Studenikin,
J.Phys.A: Math.Theor.
41(2008) 164047

$$q_{eff} \mathbf{E}_{eff} = q_m \mathbf{E}_m + q_0 \mathbf{E}$$

$$q_{eff} \mathbf{B}_{eff} = |q_m B_m + q_0 B| \mathbf{e}_z$$

matter
density

matter
rotation
frequency

where

$$q_m = -G, \quad \mathbf{E}_m = -\nabla n_n, \quad \mathbf{B}_m = 2n_n \boldsymbol{\omega}$$

matter induced “charge”, “electric” field, “magnetic” fields

... we predict :

A.Studenikin, I.Tokarev,
Nucl.Phys.B (2014)

$$E \sim 1 \text{ eV}$$

1) low-energy ν are trapped in circular orbits inside rotating neutron stars

$$R = \sqrt{\frac{2N}{Gn\omega}} < R_{NS} = 10 \text{ km}$$

$$\begin{aligned} R_{NS} &= 10 \text{ km} \\ n &= 10^{37} \text{ cm}^{-3} \\ \omega &= 2\pi \times 10^3 \text{ s}^{-1} \end{aligned}$$

2) rotating neutron stars as

filters for low-energy relic ν ?

$$T_\nu \sim 10^{-4} \text{ eV}$$

● Millicharged ν as star rotation engine

- Single ν generates feedback force with projection on rotation plane

- $F = (q_0 B + 2Gn_n \omega) \sin \theta$

single ν torque

- $M_0(t) = \sqrt{1 - \frac{r^2(t)\Omega^2 \sin^2 \theta}{4}} F r(t) \sin \theta$



total N_ν torque

$$M(t) = \frac{N_\nu}{4\pi} \int M_0(t) \sin \theta d\theta d\varphi$$

- Should effect initial star rotation
(shift of star angular velocity)

$$|\Delta\omega| = \frac{5N_\nu}{6M_S} (q_0 B + 2Gn_n \omega_0)$$

$$\Delta\omega = \omega - \omega_0$$



A.Studenikin,
I.Tokarev,
Nucl.Phys.B (2014)

• ν Star Turning mechanism (ν ST)

Studenikin, Tokarev, Nucl. Phys. B 884 (2014) 396

Escaping millicharged ν s move on curved orbits inside magnetized rotating star and feedback of effective Lorentz force should effect initial star rotation

- New astrophysical constraint on ν millicharge

$$\frac{|\Delta\omega|}{\omega_0} = 7.6\varepsilon \times 10^{18} \left(\frac{P_0}{10 \text{ s}} \right) \left(\frac{N_\nu}{10^{58}} \right) \left(\frac{1.4M_\odot}{M_S} \right) \left(\frac{B}{10^{14}G} \right)$$

- $|\Delta\omega| < \omega_0$! ...to avoid contradiction of ν ST impact with observational data on pulsars ...

$$q_0 < 1.3 \times 10^{-19} e_0$$

.. best astrophysical bound ... !

Main steps in ν oscillations

67 years!
early history of
 ν oscillations

① $\nu_e \xleftrightarrow{\text{vac}} \bar{\nu}_e$, B. Pontecorvo, 1957

② $\nu_e \xleftrightarrow{\text{vac}} \nu_\mu$, Z. Maki, M. Nakagawa, S. Sakata, 1962

③ $\nu_e \xleftrightarrow{\text{matter, } g = \text{const}} \nu_\mu$, L. Wolfenstein, 1978

④ $\nu_e \xleftrightarrow{\text{matter, } g \neq \text{const}} \nu_\mu$, S. Mikheev, A. Smirnov, 1985

• resonances in ν flavour oscillations \Rightarrow
MSW-effect, solution for ν_\odot -problem

⑤ $\nu_{eL} \xleftrightarrow{B_\perp} \nu_{eR}$, A. Cisneros, 1971
M. Voloshin, M. Vysotsky, L. Okun, 1986, ν_\odot

⑥ $\nu_{eL} \xleftrightarrow{B_\perp} \nu_{eR}, \nu_{\mu R}$, E. Akhmedov, 1988
C.-S. Lim & W. Marciano, 1988

• resonances in ν spin (spin-flavour) oscillations in matter



Bruno Pontecorvo
1913-1993

only in **B_\perp**
and
matter at rest



Бруно Понтекорво

Bruno Pontecorvo,

«Mesonium and anti-mesonium»,

Sov.Phys.JETP 6 (1957) 429

Zh.Eksp.Teor.Fiz. 33 (1957) 549-551:

«It was assumed above that there exists a conservation law for the neutrino charge, according to which a neutrino cannot change into an antineutrino in any approximation. This law has not yet been established; evidently it has been merely shown that the neutrino and antineutrino are not identical particles.

if

$m_\nu \neq 0$
then

$\nu \leftrightarrow \bar{\nu}$

MIXING

In vacuum

If the two-component neutrino theory should turn out to be incorrect ... and if the conservation law of neutrino charge would not apply, then in principle neutrino – antineutrino transitions could take place in vacuo»





Bruno Pontecorvo,
«Inverse β processes and
nonconservation of leptonic charge»,
JINR Preprint P-95, Dubna, 1957,
3 pages :

67 years of mixing
and oscillations !

«Neutrinos in vacuum can transform themselves into antineutrino and vice versa. **This means that neutrino and antineutrino are particle mixtures... So, for example, a beam of neutral leptons from a reactor which at first consists mainly of antineutrinos will change its composition and at a certain distance R from the reactor will be composed of neutrino and antineutrino in equal quantities**».

Бруно Понтекорво

● Staff member at
Faculty of Physics of
Moscow State University,
1966 - 1986

if $m_\nu \neq 0$
then $\nu \leftrightarrow \bar{\nu}$
oscillations
in vacuum

✓ spin and spin-flavour oscillations in B_{\perp}

$$\nu_{eL} \longleftrightarrow \nu_{\mu R}$$

$$B = |\mathbf{B}_{\perp}| e^{i\phi(t)}$$

● ... twisting magnetic field ...

● $P_{\nu_L \nu_R} = \sin^2 \beta \sin^2 \Omega z$ $\sin^2 \beta = \frac{(\mu_{e\mu} B)^2}{(\mu_{e\mu} B)^2 + \left(\frac{\Delta_{LR}}{4E}\right)^2}$


$$\Omega^2 = (\mu_{e\mu} B)^2 + \left(\frac{\Delta_{LR}}{4E}\right)^2$$

$$\Delta_{LR} = \frac{\Delta m^2}{2} (\cos 2\theta + 1) - 2EV_{\nu_e} + 2E\dot{\phi}$$

● Resonance amplification of oscillations in matter:

Akhmedov, 1988
Lim, Marciano

$$\Delta_{LR} \rightarrow 0$$



$$\sin^2 \beta \rightarrow 1$$

... similar to
MSW effect

New developments in ν spin and flavour oscillation

... new astrophysical probes of ν

1

generation of ν spin (flavour) oscillations by interaction with transversal matter current \mathbf{j}_\perp

P.Pustoshny, Studenikin,

Neutrino spin and spin-flavour oscillations in transversal matter currents with standard and non-standard interactions

● Phys. Rev. D98 (2018) 113009

Studenikin, Neutrino in electromagnetic fields and moving matter

● Phys.Atom.Nucl. 67 (2004) 993-1002

2

inherent interplay of ν spin and flavour oscillations in \mathbf{B}

A. Popov, Studenikin,

Neutrino eigenstates and flavour, spin and spin-flavor oscillations in a constant magnetic field

● Eur. Phys. J. C79 (2019) 144

①

ν

Neutrino spin $\nu_e^L \Leftarrow (j_\perp) \Rightarrow \nu_e^R$ and

spin-flavour $\nu_e^L \Leftarrow (j_\perp) \Rightarrow \nu_\mu^R$

oscillations engendered

by transversal matter current j_\perp
! without ~~(μ, B)~~

A. Studenikin,

Neutrino in electromagnetic fields and moving matter,

Phys. Atom. Nucl. 67 (2004) 993-1002

P. Pustoshny, A. Studenikin,

Neutrino spin and spin-flavour oscillations in

transversal matter currents with standard

and non-standard interactions, Phys. Rev. D98 (2018) 113009

ELEMENTARY PARTICLES AND FIELDS

Theory

Phys.Atom.Nucl. 67 (2004) 993-1002

Neutrino in Electromagnetic Fields and Moving Media

A. I. Studenikin*

Moscow State University, Vorob'evy gory, Moscow, 119899 Russia

Received March 26, 2003; in final form, August 12, 2003

The possible emergence of neutrino-spin oscillations (for example, $\nu_{eL} \leftrightarrow \nu_{eR}$) owing to neutrino interaction with matter under the condition that there exists a nonzero transverse current component or matter polarization (that is, $\mathbf{M}_{0\perp} \neq 0$) is the most important new effect that follows from the investigation of neutrino-spin oscillations in Section 4. So far, it has been assumed that neutrino-spin oscillations may arise only in the case where there exists a nonzero transverse magnetic field in the neutrino rest frame.

Consider ^{spin}
^{spin-flavour}

$$\nu_{eL} \rightarrow \nu_{eR}, \quad \nu_{eL} \rightarrow \nu_{\mu R}$$

$$P(\nu_i \rightarrow \nu_j) = \sin^2(2\theta_{\text{eff}}) \sin^2 \frac{\pi x}{L_{\text{eff}}}, \quad i \neq j$$

$$L_{\text{eff}} = \frac{2\pi}{\sqrt{E_{\text{eff}}^2 + \Delta_{\text{eff}}^2}}$$

$$\sin^2 2\theta_{\text{eff}} = \frac{E_{\text{eff}}^2}{E_{\text{eff}}^2 + \Delta_{\text{eff}}^2}, \quad \Delta_{\text{eff}}^2 = \frac{\mu}{\gamma_\nu} |\mathbf{M}_{0\parallel} + \mathbf{B}_{0\parallel}|, \quad E_{\text{eff}} = \mu \left| \mathbf{B}_\perp + \frac{1}{\gamma_\nu} \mathbf{M}_{0\perp} \right|$$

A. Studenikin,
"Neutrinos in electromagnetic
fields and moving media",
Phys. Atom. Nucl. 67 (2004)

• transversal current \mathbf{j}

$$\vec{M}_0 = \gamma_\nu \rho n_e \left(\vec{\beta}_\nu (1 - \vec{\beta}_\nu \vec{v}_e) - \frac{1}{\gamma_\nu} \vec{v}_{e\perp} \right)$$

(Annotations: $\vec{\beta}_\nu$ is speed of ν ; \vec{v}_e is transversal speed of matter; ρ is matter density; $\vec{v}_{e\perp}$ is transversal speed of matter.)

$$\gamma_\nu = \frac{E_\nu}{m_\nu}$$

where

$$\rho = \frac{G_F}{2\mu_\nu \sqrt{2}} (1 + 4 \sin^2 \theta_W)$$


... the effect of ν helicity

$$\nu_{eL} \rightarrow \nu_{eR}, \quad \nu_{eL} \rightarrow \nu_{\mu R}$$

conversions and oscillations induced by transversal matter currents has been confirmed in studies of ν propagation in astrophysical media:

- J. Serreau and C. Volpe, Neutrino-antineutrino correlations in dense anisotropic media, Phys. Rev. D90 (2014) 125040
- V. Cirigliano, G. M. Fuller, and A. Vlasenko, A new spin on neutrino quantum kinetics Phys. Lett. B747 (2015) 27
- A. Kartavtsev, G. Raffelt, and H. Vogel, ● Neutrino propagation in media: flavor-, helicity-, and pair correlations, Phys. Rev. D91 (2015) 125020 ...

— Neutrino spin (spin-flavour) oscillations in transversal matter currents ... quantum treatment ...

- \checkmark spin evolution effective Hamiltonian in moving matter ? transversal and longitudinal currents 
- two flavor \checkmark with two helicities: $\nu_f = (\nu_e^+, \nu_e^-, \nu_\mu^+, \nu_\mu^-)^T$
- \checkmark interaction with matter composed of neutrons: $n = \frac{n_0}{\sqrt{1-v^2}}$ neutron number density in laboratory reference frame

$\mathbf{v} = (v_1, v_2, v_3)$ velocity of matter

- $L_{\text{int}} = -f^\mu \sum_l \bar{\nu}_l(x) \gamma_\mu \frac{1 + \gamma_5}{2} \nu_l(x) = -f^\mu \sum_i \bar{\nu}_i(x) \gamma_\mu \frac{1 + \gamma_5}{2} \nu_i(x)$ $l = e, \text{ or } \mu$
 $i = 1, 2$

$$f^\mu = -\frac{G_F}{2\sqrt{2}} j_n^\mu$$

$$\begin{aligned} \nu_e^\pm &= \nu_1^\pm \cos \theta + \nu_2^\pm \sin \theta, \\ \nu_\mu^\pm &= -\nu_1^\pm \sin \theta + \nu_2^\pm \cos \theta \end{aligned}$$

\checkmark flavour and mass states

- $j_n^\mu = n(1, \mathbf{v})$

P. Pustoshny, A. Studenikin,
Phys. Rev. D98 (2018) 113009

✓ (2 flavours × 2 helicities) evolution equation

$$i \frac{d}{dt} \nu_f^s = \left(\underbrace{H_0}_{\text{vacuum}} + \underbrace{\Delta H_0^{SM}}_{\text{matter at rest}} + \underbrace{\Delta H_{j_{||}+j_{\perp}}^{SM}}_{\text{moving matter}} + \underbrace{\Delta H_{B_{||}+B_{\perp}}^{SM}}_{\mathbf{B}} + \underbrace{\Delta H_0^{NSI}}_{\text{matter at rest}} + \underbrace{\Delta H_{j_{||}+j_{\perp}}^{NSI}}_{\text{moving matter}} \right) \nu_f^s$$

Standard Model Non-Standard Interactions

Resonant amplification of ✓ oscillations:


- $\nu_e^L \Leftarrow (j_{\perp}) \Rightarrow \nu_e^R$ by longitudinal matter current $j_{||}$
- $\nu_e^L \Leftarrow (j_{\perp}) \Rightarrow \nu_e^R$ by longitudinal $\mathbf{B}_{||}$
- $\nu_e^L \Leftarrow (j_{\perp}) \Rightarrow \nu_{\mu}^R$ by matter-at-rest effect
- $\nu_e^L \Leftarrow (j_{\perp}^{NSI}) \Rightarrow \nu_{\mu}^R$ by matter-at-rest effect

$$\nu_e^L \leftarrow (j_\perp) \Rightarrow \nu_e^R$$

a model of short GRB

$$D \sim 20 \text{ km}$$

$$d \sim 20 \text{ km}$$



- Consider v escaping central neutron star with inclination angle α from accretion

disk: $B_{||} = B \sin \alpha \sim \frac{1}{2} B$

- Toroidal bulk of rotating dense matter with $\omega = 10^3 \text{ s}^{-1}$

- transversal velocity of matter

$$v_\perp = \omega D = 0.067 \text{ and } \gamma_n = 1.002$$

$$E_{eff} = \left(\frac{\eta}{\gamma}\right)_{ee} \tilde{G} n v_\perp = \frac{\cos^2 \theta}{\gamma_{11}} \tilde{G} n v_\perp \approx \tilde{G} n_0 \frac{\gamma_n}{\gamma_\nu} v_\perp$$

$$\Delta_{eff} = \left| \left(\frac{\mu}{\gamma}\right)_{ee} B_{||} + \eta_{ee} \tilde{G} n \beta \right| \approx \left| \frac{\mu_{11}}{\gamma_\nu} B_{||} - \tilde{G} n_0 \gamma_n \right|$$

$$B_{||} \beta = -1$$

$$E_{eff} \geq \Delta_{eff}$$

resonance condition

$$\left| \frac{\mu_{11} B_{||}}{\tilde{G} n_0 \gamma_n} - \gamma_\nu \right| \leq 1$$

• Perego et al,
Mon.Not.Roy.Astron.Soc.
443 (2014) 3134
• Grigoriev, Lokhov,
Studenikin, Ternov,
JCAP 1711 (2017) 024

Resonance amplification of spin-flavor oscillations
(in the absence of \mathbf{j}_{\parallel})

$$\nu_e^L \Leftrightarrow (j_{\perp}, B_{\perp}) \Rightarrow \nu_{\mu}^R$$

$$\vec{B} = \vec{B}_{\perp} + \vec{B}_{\parallel} \rightarrow \mathbf{0}$$

Criterion – oscillations are important:

$$\sin^2 2\theta_{\text{eff}} = \frac{E_{\text{eff}}^2}{E_{\text{eff}}^2 + \Delta_{\text{eff}}^2} \geq \frac{1}{2}$$

$$E_{\text{eff}} = \left| \mu_{e\mu} B_{\perp} + \left(\frac{\eta}{\gamma} \right)_{e\mu} \tilde{G} n v_{\perp} \right| \geq \left| \Delta M - \frac{1}{2} \left(\frac{\mu_{11}}{\gamma_{11}} + \frac{\mu_{22}}{\gamma_{22}} \right) B_{\parallel} - \tilde{G} n (1 - v\beta) \right|$$

neglecting $\vec{B} = \vec{B}_{\perp} + \vec{B}_{\parallel} \rightarrow \mathbf{0}$:

$$L_{\text{eff}} = \frac{\pi}{\left(\frac{\eta}{\gamma} \right)_{e\mu} \tilde{G} n v_{\perp}} \quad \left(\frac{\eta}{\gamma} \right)_{e\mu} \approx \frac{\sin 2\theta}{\gamma_{\nu}}$$

$$\left| \left(\frac{\eta}{\gamma} \right)_{e\mu} \tilde{G} n v_{\perp} \right| \geq \left| \Delta M - \tilde{G} n (1 - v\beta) \right|$$



$$\tilde{G} n \sim \Delta M$$

•

$$\Delta m^2 = 7.37 \times 10^{-5} \text{ eV}^2$$

$$\tilde{G} = \frac{G_F}{2\sqrt{2}} = 0.4 \times 10^{-23} \text{ eV}^{-2}$$

$$\sin^2 \theta = 0.297$$

$$p_0^{\nu} = 10^6 \text{ eV}$$

$$\Rightarrow \Delta M = 0.75 \times 10^{-11} \text{ eV}$$

$$n_0 \sim \frac{\Delta M}{\tilde{G}} = 10^{12} \text{ eV}^3 \approx 10^{26} \text{ cm}^{-3}$$

$$L_{\text{eff}} = \frac{\pi}{\left(\frac{\eta}{\gamma} \right)_{e\mu} \tilde{G} n v_{\perp}} \approx 5 \times 10^{11} \text{ km}$$

• $L_{\text{eff}} \approx 10 \text{ km}$ (within short GRB) if $n_0 \approx 5 \times 10^{36} \text{ cm}^{-3}$ •

2



A.Popov, A.Studenikin, Eur. Phys. J. C79 (2019) 144

“Neutrino eigenstates and
flavour, spin and spin-flavour oscillations
in a constant magnetic field”

- $\nu_e^L \leftrightarrow \nu_\mu^L$ $\nu_e^L \leftrightarrow \nu_e^R$ $\nu_e^L \leftrightarrow \nu_\mu^R$

Consider two flavour ν with two helicities as superposition of helicity mass states $\nu_i^{L(R)}$

Popov, Studenikin, Eur. Phys. J. C79 (2019) 144

$\nu_e^{L(R)} = \nu_1^{L(R)} \cos \theta + \nu_2^{L(R)} \sin \theta,$
 $\nu_\mu^{L(R)} = -\nu_1^{L(R)} \sin \theta + \nu_2^{L(R)} \cos \theta$ however, $\nu_i^{L(R)}$ are not stationary states in magnetic field $\mathbf{B} = (B_\perp, 0, B_\parallel)$

$\nu_i^L(t) = c_i^+ \nu_i^+(t) + c_i^- \nu_i^-(t),$
 $\nu_i^R(t) = d_i^+ \nu_i^+(t) + d_i^- \nu_i^-(t)$ $\leftarrow \nu_i^{-(+)}$ stationary states in \mathbf{B}

• Dirac equation $(\gamma_\mu p^\mu - m_i - \mu_i \Sigma \mathbf{B}) \nu_i^s(p) = 0$ in a constant \mathbf{B}

$\hat{H}_i \nu_i^s = E \nu_i^s$ $\hat{H}_i = \gamma_0 \boldsymbol{\gamma} \mathbf{p} + \mu_i \gamma_0 \Sigma \mathbf{B} + m_i \gamma_0$ ($s = \pm 1$) $\mu_{ij}(i \neq j) = 0$

ν spin operator that commutes with \hat{H}_i : “bra-ket” products

$\hat{S}_i = \frac{1}{N} \left[\Sigma \mathbf{B} - \frac{i}{m_i} \gamma_0 \gamma_5 [\Sigma \times \mathbf{p}] \mathbf{B} \right]$ $\hat{S}_i |\nu_i^s\rangle = s |\nu_i^s\rangle, s = \pm 1$ $\langle \nu_i^s | \nu_k^{s'} \rangle = \delta_{ik} \delta_{ss'}$

$\frac{1}{N} = \frac{m_i}{\sqrt{m_i^2 \mathbf{B}^2 + p^2 B_\perp^2}}$

• ν energy spectrum

$E_i^s = \sqrt{m_i^2 + p^2 + \mu_i^2 \mathbf{B}^2 + 2\mu_i s \sqrt{m_i^2 \mathbf{B}^2 + p^2 B_\perp^2}}$

Probabilities of ν oscillations (flavour, spin and spin-flavour)

$\nu_e^L \leftrightarrow \nu_\mu^L$ $P_{\nu_e^L \rightarrow \nu_\mu^L}(t) = |\langle \nu_\mu^L | \nu_e^L(t) \rangle|^2$ $\mu_\pm = \frac{1}{2}(\mu_1 \pm \mu_2)$ magnetic moments of ν mass states

flavour

$$P_{\nu_e^L \rightarrow \nu_\mu^L}(t) = \sin^2 2\theta \left\{ \cos(\mu_1 B_\perp t) \cos(\mu_2 B_\perp t) \sin^2 \frac{\Delta m^2}{4p} t + \sin^2(\mu_+ B_\perp t) \sin^2(\mu_- B_\perp t) \right\}$$

spin

$$P_{\nu_e^L \rightarrow \nu_e^R} = \left\{ \sin(\mu_+ B_\perp t) \cos(\mu_- B_\perp t) + \cos 2\theta \sin(\mu_- B_\perp t) \cos(\mu_+ B_\perp t) \right\}^2 - \sin^2 2\theta \sin(\mu_1 B_\perp t) \sin(\mu_2 B_\perp t) \sin^2 \frac{\Delta m^2}{4p} t.$$

spin-flavour

$$P_{\nu_e^L \rightarrow \nu_\mu^R}(t) = \sin^2 2\theta \left\{ \sin^2 \mu_- B_\perp t \cos^2(\mu_+ B_\perp t) + \sin(\mu_1 B_\perp t) \sin(\mu_2 B_\perp t) \sin^2 \frac{\Delta m^2}{4p} t \right\}$$

... interplay of oscillations on vacuum and on magnetic frequencies


$$\omega_{vac} = \frac{\Delta m^2}{4p}$$

$$\omega_B = \mu B_\perp$$

• For the case $\mu_1 = \mu_2$, probability of flavour oscillations

$$P_{\nu_e^L \rightarrow \nu_\mu^L} = \left(1 - \sin^2(\mu B_\perp t)\right) \sin^2 2\theta \sin^2 \frac{\Delta m^2}{4p} t = \left(1 - P_{\nu_e^L \rightarrow \nu_e^R}^{cust}\right) P_{\nu_e^L \rightarrow \nu_\mu^L}^{cust}$$

flavour no spin oscillations




... amplitude of flavour oscillations on vacuum frequency $\omega_{vac} = \frac{\Delta m^2}{4p}$ is modulated by magnetic frequency $\omega_B = \mu B_\perp$

Fig. 1 The probability of the neutrino flavour oscillations $\nu_e^L \rightarrow \nu_\mu^L$ in the transversal magnetic field $B_\perp = 10^{16} G$ for the neutrino energy $p = 1 MeV$, $\Delta m^2 = 7 \times 10^{-5} eV^2$ and magnetic moments $\mu_1 = \mu_2 = 10^{-20} \mu_B$.

For the case $\mu_1 = \mu_2$, probability of spin oscillations

$$P_{\nu_e^L \rightarrow \nu_e^R} = \left[1 - \sin^2 2\theta \sin^2 \left(\frac{\Delta m^2}{4p} t \right) \right] \sin^2(\mu B_{\perp} t) = \left(1 - P_{\nu_e^L \rightarrow \nu_{\mu}^L}^{cust} \right) P_{\nu_e^L \rightarrow \nu_e^R}^{cust}$$

spin no flavour oscillations



... amplitude of spin oscillations on magnetic frequency $\omega_B = \mu B_{\perp}$
 is modulated by
 vacuum $\omega_{vac} = \frac{\Delta m^2}{4p}$
frequency


A. Popov, A.S.,
 Eur. Phys. J. C
 79 (2019) 144

Fig. 2 The probability of the neutrino spin oscillations $\nu_e^L \rightarrow \nu_e^R$ in the transversal magnetic field $B_{\perp} = 10^{16} \text{ G}$ for the neutrino energy $p = 1 \text{ MeV}$, $\Delta m^2 = 7 \times 10^{-5} \text{ eV}^2$ and magnetic moments $\mu_1 = \mu_2 = 10^{-20} \mu_B$.

• For the case $\mu_1 = \mu_2$, probability of **spin-flavour** oscillations

$$P_{\nu_e^L \rightarrow \nu_\mu^R} = \sin^2(\mu B_\perp t) \sin^2 2\theta \sin^2 \frac{\Delta m^2}{4p} t = P_{\nu_e^L \rightarrow \nu_e^R}^{cust} P_{\nu_e^L \rightarrow \nu_\mu^L}^{cust}$$

spin-flavour



$$P_{\nu_e^L \rightarrow \nu_\mu^L}^{cust} = \sin^2 2\theta \sin^2 \frac{\Delta m^2}{4p} t$$

$$P_{\nu_e^L \rightarrow \nu_e^R}^{cust} = \sin^2(\mu B_\perp t)$$

... interplay of oscillations
 on vacuum $\omega_{vac} = \frac{\Delta m^2}{4p}$
 and
 on magnetic $\omega_B = \mu B_\perp$
 frequencies

... in literature usually:

- $P_{\nu_e^L \rightarrow \nu_\mu^R} = \sin^2(\mu_{e\mu} B_\perp t) = 0$
 $\mu_{e\mu} = \frac{1}{2}(\mu_2 - \mu_1) \sin 2\theta$
 $\mu_1 = \mu_2, \mu_{ij} = 0, i \neq j$
- ... M. Dvornikov, J. Maalampi, Phys. Lett. B 657 (2007) 217

Fig. 3 The probability of the neutrino spin flavour oscillations $\nu_e^L \rightarrow \nu_\mu^R$ in the transversal magnetic field $B_\perp = 10^{16} G$ for the neutrino energy $p = 1 MeV$, $\Delta m^2 = 7 \times 10^{-5} eV^2$ and magnetic moments $\mu_1 = \mu_2 = 10^{-20} \mu_B$.

- For completeness: ν survival $\nu_e^L \leftrightarrow \nu_e^L$ probability

... depends on μ_ν and \mathbf{B}

$$P_{\nu_e^L \rightarrow \nu_e^L}(t) = \left\{ \cos(\mu_+ B_\perp t) \cos(\mu_- B_\perp t) - \cos 2\theta \sin(\mu_+ B_\perp t) \sin(\mu_- B_\perp t) \right\}^2 - \sin^2 2\theta \cos(\mu_1 B_\perp t) \cos(\mu_2 B_\perp t) \sin^2 \frac{\Delta m^2}{4p} t$$

\sum of all probabilities (as it should be...):

$$P_{\nu_e^L \rightarrow \nu_\mu^L} + P_{\nu_e^L \rightarrow \nu_e^R} + P_{\nu_e^L \rightarrow \nu_\mu^R} + P_{\nu_e^L \rightarrow \nu_e^L} = 1$$

A. Popov, A.S., Eur. Phys. J. C79 (2019) 144

the discovered correspondence between flavour and spin oscillations in \mathbf{B} can be important in studies of ν propagation in astrophysical environments

3 New effect in \checkmark flavor oscillation in moving matter =

$$\nu_e^L \leftarrow (j_{||}, j_{\perp}) \Rightarrow \nu_{\mu}^L \quad j_{\perp} = n\mathbf{v}_{\perp}$$

longitudinal matter current transversal matter current invariant number density

Studenikin,
Physics of Particles and Nuclei 55 (2024) 1444
+ arXiv: 1912.12491

Equal role of j_{\perp} and B_{\perp} in generation of

$$\nu_e^L \leftarrow (j_{\perp}, B_{\perp}) \Rightarrow \nu_e^R \quad \text{spin oscillations}$$

$$\nu_e^L \leftarrow (j_{\perp}, B_{\perp}) \Rightarrow \nu_{\mu}^R \quad \text{spin-flavour}$$

Probability of \checkmark flavor oscillations $\nu_e^L \leftarrow (j_{||}, j_{\perp}) \Rightarrow \nu_{\mu}^L$ in moving matter

$$P_{\nu_e^L \rightarrow \nu_{\mu}^L}^{(j_{||}+j_{\perp})}(t) = \left(1 - P_{\nu_e^L \rightarrow \nu_e^R}^{(j_{\perp})} - P_{\nu_e^L \rightarrow \nu_{\mu}^R}^{(j_{\perp})} \right) P_{\nu_e^L \rightarrow \nu_{\mu}^L}^{(j_{||})}$$

$$P_{\nu_e^L \rightarrow \nu_{\mu}^L}^{(j_{||})}(t) = \sin^2 2\theta_{eff} \sin^2 \omega_{eff} t, \quad \omega_{eff} = \frac{\Delta m_{eff}^2}{4p_0^{\nu}}$$

probability of flavor oscillations in $j_{||}$

probability of spin survival (not spin flip)

$$P_{\nu_e^L \rightarrow \nu_e^R}^{j_{\perp}}(t) = \frac{\left(\frac{\eta}{\gamma}\right)_{ee}^2 v_{\perp}^2}{\left(\frac{\eta}{\gamma}\right)_{ee}^2 v_{\perp}^2 + (1 - v\beta)^2} \sin^2 \omega_{ee}^{j_{\perp}} t$$

spin oscillations in j_{\perp}

$$P_{\nu_e^L \rightarrow \nu_{\mu}^R}^{j_{\perp}}(t) = \frac{\left(\frac{\eta}{\gamma}\right)_{e\mu}^2 v_{\perp}^2}{\left(\frac{\eta}{\gamma}\right)_{e\mu}^2 v_{\perp}^2 + \left(\frac{\Delta M}{\tilde{G}n} - (1 - v\beta)\right)^2} \sin^2 \omega_{e\mu}^{j_{\perp}} t$$

spin-flavor oscillations in j_{\perp}

$$\omega_{ee}^{j_{\perp}} = \tilde{G}n \sqrt{\left(\frac{\eta}{\gamma}\right)_{ee}^2 v_{\perp}^2 + (1 - v\beta)^2}$$

... is modulated by two "matter" frequencies ...

$$\omega_{e\mu}^{j_{\perp}} = \tilde{G}n \sqrt{\left(\frac{\eta}{\gamma}\right)_{e\mu}^2 v_{\perp}^2 + \left(\frac{\Delta M}{\tilde{G}n} - (1 - v\beta)\right)^2}$$

$$\left(\frac{\eta}{\gamma}\right)_{ee} = \frac{\cos^2 \theta}{\gamma_{11}} + \frac{\sin^2 \theta}{\gamma_{22}} \quad \gamma_{\alpha\alpha'}^{-1} = \frac{1}{2}(\gamma_{\alpha}^{-1} + \gamma_{\alpha'}^{-1}) \quad \gamma_{\alpha}^{-1} = \frac{m_{\alpha}}{E_{\alpha}}$$

$$\left(\frac{\eta}{\gamma}\right)_{e\mu} = \frac{\sin 2\theta}{\tilde{\gamma}_{21}} \quad \tilde{\gamma}_{\alpha\alpha'}^{-1} = \frac{1}{2}(\gamma_{\alpha}^{-1} - \gamma_{\alpha'}^{-1})$$

Manifestations of nonzero Majorana CP -violating phases in oscillations of supernova neutrinos

Artem Popov^{*}

Department of Theoretical Physics, Moscow State University, Moscow 119991, Russia

Alexander Studenikin[†]Department of Theoretical Physics, Moscow State University, Moscow 119991, Russia
and Joint Institute for Nuclear Research, Dubna 141980, Russia
 (Received 14 February 2021; accepted 18 May 2021; published 22 June 2021)

We investigate effects of nonzero Dirac and Majorana CP -violating phases on neutrino-antineutrino oscillations in a magnetic field of astrophysical environments. It is shown that in the presence of strong magnetic fields and dense matter, nonzero CP phases can induce new resonances in the oscillations channels $\nu_e \leftrightarrow \bar{\nu}_e$, $\nu_e \leftrightarrow \bar{\nu}_\mu$, and $\nu_e \leftrightarrow \bar{\nu}_\tau$. We also consider all other possible oscillation channels with ν_μ and ν_τ in the initial state. The resonances can potentially lead to significant phenomena in neutrino oscillations accessible for observation in experiments. In particular, we show that neutrino-antineutrino oscillations combined with Majorana-type CP violation can affect the $\bar{\nu}_e/\nu_e$ ratio for neutrinos coming from the supernovae explosion. This effect is more prominent for the normal neutrino mass ordering. The detection of supernovae neutrino fluxes in the future experiments, such as JUNO, DUNE, and Hyper-Kamiokande, can give an insight into the nature of CP violation and, consequently, provides a tool for distinguishing the Dirac or Majorana nature of neutrinos.

DOI: 10.1103/PhysRevD.103.115027

I. INTRODUCTION

CP symmetry implies that the equations of motion of a system remain invariant under the CP transformation, that is a combination of charge conjugation (C) and parity inversion (P). In 1964, with the discovery of the neutral kaon decay [1], it was confirmed that CP is not an underlying symmetry of the electroweak interactions theory, thus opening a vast field of research in CP violation. Currently, CP violation is a topic of intense studies in particle physics that also has important implications in cosmology. In 1967, Sakharov proved that the existence of CP violation is a necessary condition for generation of the baryon asymmetry through baryogenesis in the early Universe [2]. A review of possible baryogenesis scenarios can be found in [3].

Today we have solid understanding of CP violation in the quark sector, that appears due to the complex phase

in the Cabibbo-Kobayashi-Maskawa matrix parametrization. Its magnitude is expressed by the Jarlskog invariant $\mathcal{J}_{CKM} = (3.18 \pm 0.15) \times 10^{-5}$ [4], which seems to be excessively small to engender baryogenesis at the electroweak phase transition scale [3]. However, in addition to experimentally confirmed CP violation in the quark sector, CP violation in the lepton (neutrino) sector hypothetically exists (see [5] for a review). Leptonic CP violation is extremely difficult to observe due to weakness of neutrino interactions. In 2019, a first breakthrough happened when NO ν A [6] and T2K [7] collaborations reported constraints on the Dirac CP -violating phase in neutrino oscillations. Hopefully, future gigantic neutrino experiments, such as DUNE [8] and Hyper-Kamiokande [9], also JUNO [10] with detection of the atmospheric neutrinos, will have a good chance significantly improve this results. Note that leptonic CP violation plays an important role in baryogenesis through leptogenesis scenarios [11].

The CP -violation pattern in the neutrino sector depends on whether neutrino is a Dirac or Majorana particle. The Pontecorvo-Maki-Nakagawa-Sakata (PMNS) neutrino mixing matrix in the most common parametrization has the following form

*ar.popov@physics.msu.ru

†studenik@srj.sinp.msu.ru

A. Popov, A. Studenikin

Phys. Rev. D **103** (2021) 115027

... the role of Majorana CP -violating phases in neutrino oscillations

$$\nu_e \leftrightarrow \bar{\nu}_{e,\mu,\tau}$$

in strong \mathbf{B} and dense matter of supernovae for two mass hierarchies

... Majorana CP phases induce new resonances

... a tool for distinguishing Dirac-Majorana nature of ν



Neutrino quantum decoherence engendered by neutrino radiative decay

Konstantin Stankevich*

*Department of Theoretical Physics, Faculty of Physics, Lomonosov Moscow State University,
119992 Moscow, Russia*

Alexander Studenikin†

*Department of Theoretical Physics, Faculty of Physics, Lomonosov Moscow State University,
Moscow 119991, Russia
and Dzhelapov Laboratory of Nuclear Problems, Joint Institute for Nuclear Research,
Dubna 141980, Moscow Region, Russia*

 (Received 26 November 2019; accepted 6 February 2020; published 5 March 2020)

A new theoretical framework, based on the quantum field theory of open systems applied to neutrinos, has been developed to describe the neutrino evolution in external environments accounting for the effect of the neutrino quantum decoherence. The developed new approach enables one to obtain the explicit expressions of the decoherence and relaxation parameters that account for a particular process, in which the neutrino participates, and also for the characteristics of an external environment and of the neutrino itself, including the neutrino energy. We have used this approach to consider a new mechanism of the neutrino quantum decoherence engendered by the neutrino radiative decay to photons and dark photons in an astrophysical environment. The importance of the performed studies is highlighted by the prospects of the forthcoming new large volume neutrino detectors that will provide new frontier in high-statistics measurements of neutrino fluxes from supernovae.

DOI: 10.1103/PhysRevD.101.056004

I. INTRODUCTION

Half a century ago Gribov and Pontecorvo derived [1] the first analytical expression for the neutrino oscillation probability that has opened a new era in the theoretical and experimental studies of the neutrino oscillation phenomenon. The neutrino oscillation patterns can be modified by neutrino interactions with external environments including electromagnetic fields that can influence neutrinos in the case neutrinos have nonzero electromagnetic properties [2]. The phenomenon of neutrino oscillations can proceed only in the case of the coherent superposition of neutrino mass states. An external environment can modify a neutrino evolution in a way that conditions for the coherent superposition of neutrino mass states are violated. Such a violation is called quantum decoherence of neutrino states and leads to the suppression of flavor neutrino oscillations. It should be noted that the quantum neutrino decoherence differs from the standard neutrino decoherence that appears

due to separation of neutrino wave packets, the effect that is not considered below.

The quantum neutrino decoherence has attracted a growing interest during the last 15 years. Within reasonable amount of the performed studies, the method based on the Lindblad master equation [3,4] for describing neutrino evolution has been used. This approach is usually considered as the most general one that gives a possibility to study neutrino quantum decoherence as a consequence of standard and nonstandard interactions of a neutrino system with an external environment [5–15].

The Lindblad master equation can be written in the following form (see, for instance, [13]):

$$\frac{\partial \rho_\nu(t)}{\partial t} = -i[H_S, \rho_\nu(t)] + D[\rho_\nu], \quad (1)$$

where ρ_ν is the density matrix that describes the neutrino evolution, H_S is the Hamiltonian, and the dissipation term (or dissipator) is given by

$$D[\rho_\nu(t)] = \frac{1}{2} \sum_{k=1}^{N^2-1} [V_k, \rho_\nu V_k^\dagger] + [V_k \rho_\nu, V_k^\dagger], \quad (2)$$

where V_k are dissipative operators that arise from interaction between the neutrino system and the external

Stankevich, Studenikin, Neutrino quantum decoherence engendered by neutrino radiative decay Phys.Rev.D 101 (2020) 056004

...  radiative decay as a source of
quantum decoherence in extreme
astrophysical environments



... observable consequences
for SN 
(JUNO, DUNE, Hyper-Kamiokande)

*kl.stankevich@physics.msu.ru
†studenik@srd.sinp.msu.ru

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.



1 Electromagnetic Properties of ν




C.Giunti, A.Studenikin, "electromagnetic interactions: A window to new physics", Rev.Mod.Phys, 2015

MSU Alexander Studenikin NCPM

Studenikin, "Overview of ν electromagnetic properties 2022", arXiv:2301.06071;

"Electromagnetic properties of neutrino 2023", Physics of Particles and Nuclei 55 (2024) 1144



1 ν EP theory - ν vertex function

matrices in ν mass eigenstates space

$$\Lambda_\mu^{if}(q) = f_Q^{if}(q^2)\gamma_\mu + f_M^{if}(q^2)i\sigma_{\mu\nu}q^\nu + f_E^{if}(q^2)\sigma_{\mu\nu}q^\nu\gamma_5 + f_A^{if}(q^2)(q^2\gamma_\mu - q_\mu\not{q})\gamma_5,$$

form factors $f_X^{if}(q^2)$ at $q^2=0$ static EP of ν

electric charge magnetic moment electric moment anapole moment

Dirac ν Majorana

q_{if}	$q=0$	} CPT + charge conservation
$\mu_{if} \neq 0$	$\mu_{if}^{(i \neq f)}$	
ϵ_{if}	$\epsilon_{if}^{(i \neq f)}$	
a_{if}	a_{if}	

Hermiticity and discrete symmetries of EM current put constraints on form factors

$$\mu_{jj}^D = \frac{3e_0 G_F m_j}{8\sqrt{2}\pi^2} \approx 3.2 \times 10^{-19} \mu_B \left(\frac{m_j}{1 \text{ eV}} \right)$$

Fujikawa & Shrock, 1980

- much greater values are Beyond Minimally Extended SM
- transition moments $\mu_{i \neq f}, \epsilon_{i \neq f}$ are GIM suppressed

3 ν EMP experimental bounds

$$\mu_\nu^{eff} < 2.8 \times 10^{-11} \mu_B$$

GEMMA 2012 Borexino 2017 ~ XENON1T 2020 astrophys., Raffelt ea 1988, 2020 Arcoa Dias ea 2015

$$q_\nu < \begin{cases} \sim 10^{-12} \\ \sim 10^{-19} \\ \sim 10^{-21} \end{cases} e_0$$

reactor ν scattering AS '14, Chen ea '14 AS '14 (astrophysics) neutrality of matter

• charge rad. $\langle r_\nu^2 \rangle$ is most accessible for exp. observations •

2 ν electromagnetic properties: **Future prospects**

● new constraint on μ_ν (and q_ν) from GEMMA-2/ ν GeN and Borexino (?)

● XENON1T an excess in electronic recoil events in < 7 keV (2-3 keV) over known backgrnds $\Rightarrow \mu_\nu \in (1.4, 2.9) \times 10^{-11} \mu_B$ E. Aprile et al, XENON1T coll., *Phys. Rev. D* 102 (2020) 072004

● XENONnT \Rightarrow XENON1T signal from transition neutrino magnetic moments, O. Miranda, D. Papoulias, M. Tórtola, J. W. F. Valle, *Phys. Lett. B* 808 (2020) 135685

● new improved limit from stellar evolution data for global cluster ω -Centauri $\Rightarrow \mu_\nu < 2.2 \times 10^{-12} \mu_B$ S. Arceo-Diaz, K.-P. Schroder, K. Zuber, D. Jack, *Astropart. Phys.* 70 (2015) 1

● new improved limit $\mu_\nu < 1.2 \times 10^{-12} \mu_B$ F. Capozzi, G. Raffelt, *Phys. Rev. D* 102 (2020) 083007

comes from improved new calibrations of tip of red-giant branch which allows one to constrain novel energy losses

● new setup to observe coherent elastic neutrino-atom scattering using electron antineutrinos from tritium decay and a liquid helium target \Rightarrow upper limit

$\mu_\nu < 7 \times 10^{-13} \mu_B$ M. Cadeddu, F. Dordei, C. Giunti, K. Kouzakov, E. Picciau, A. Studenikin, Potentialities of a low-energy detector based on 4 He evaporation to observe atomic effects in coherent neutrino scattering and physics perspectives, *Phys. Rev. D* 100 (2019) no. 7, 073014

⇒ **XENONnT Collaboration** E. Aprile et al. (XENON Coll.), Search for New Physics in electronic recoil data from XENONnT, *Phys.Rev.Lett.* 122 (2022) 161805

● Upgraded experiment has managed to rule out the so-called XENON1T excess by using a new larger liquid xenon (LXe)

Implications of first LUX-ZEPLIN and XENONnT results

A.Khan, Light new physics and neutrino electromagnetic interactions in XENONnT, *Phys.Lett.B* 837 (2023) 137650

M.Atzori Corona, W.Bonivento, M.Cadeddu, N.Cargioli, F.Dordei, New constraint on neutrino magnetic moment from LZ dark matter search results, *Phys.Rev.D* 107 (2023) 053001

K.ShivaSankar, A.Majumdar, D.Papoulias, H.Prajapati, R. Srivastava, *Phys.Lett. B* 839 (2023) 137742

C.Giunti, C.Ternes, Testing neutrino electromagnetic properties at current and future dark matter experiments, *Phys. Rev. D* 108 (2023) 095044

J. Aalbers et al., A next-generation liquid xenon observatory for dark matter and neutrino physics, *J.Phys.G* 50 (2023) 1, 013001

! ... new stringent upper limits

! ... new stringent upper limits on

$$\mu_\nu \sim \text{few} \times 10^{-12} \mu_B$$

$$q_\nu \sim 10^{-13} e_0$$

! **2024 + soon ...**

! **vGeN experiment ... low threshold ...** $T \sim 200 \text{ eV}$

$$\mu_\nu \sim (5 - 9) \times 10^{-12} \mu_B$$

I.Alekseev et al., First results of the vGeN experiment on coherent elastic neutrino-nucleus scattering, *Phys.Rev.D* 106 (2022) 5, L051101



A.Studenikin, *Europhys. Lett.* 107 (2014) 210011

$$|q_\nu| < 1.1 \times 10^{-13} e_0$$

- Initial interest in effect of ν spin oscillations in B on supernovae ν fluxes

- H.Nunokawa, R.Tomas, J. W. F. Valle,
Type-II supernovae and neutrino magnetic moment,
Astropart.Phys. 11 (1999) 317

has recently increased again

(new large-volume ν detectors, DUNE, Hyper-Kamiokande and JUNO, open up new possibilities in precise determination of flavour ratios in supernovae ν fluxes)

- V.Brdar, A.de Gouvêa, Y.-Y Li, P. Machado,
The neutrino magnetic moment portal and supernovae: New constraints
and multimessenger opportunities, *Phys.Rev. D* 107 (2023) 073005

- E.Wang, Resonant spin-flavor precession of sterile neutrinos, *JCAP* 05 (2024) 056

- S.Jana, Y.Porto, New resonances of supernova neutrinos in magnetic fields,
Phys. Rev. Lett. 132 (2024) 101005

probe non-negligible

μ_ν

potentially down to $\sim 10^{-15} \mu_B$

Resonances of Supernova Neutrinos in Twisting Magnetic Fields


Sudip Jana^{1,*} and Yago Porto^{2,†}

¹*Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany*

²*Instituto de Física Gleb Wataghin - UNICAMP, 13083-859, Campinas, São Paulo, Brazil*

(Received 28 March 2023; revised 20 June 2023; accepted 14 February 2024; published 5 March 2024)

We investigate the effect of resonant spin conversion of the neutrinos induced by the geometrical phase in a twisting magnetic field. We find that the geometrical phase originating from the rotation of the transverse magnetic field along the neutrino trajectory can trigger a resonant spin conversion of Dirac neutrinos inside the supernova, even if there were no such transitions in the fixed-direction field case. We have shown that, even though resonant spin conversion is too weak to affect solar neutrinos, it could have a remarkable consequence on supernova neutronization bursts where very intense magnetic fields are quite likely. We demonstrate how the flavor composition at Earth can be used as a probe to establish the presence of non-negligible magnetic moments, potentially down to $10^{-15} \mu_B$ in upcoming neutrino experiments like the Deep Underground Neutrino Experiment and the Hyper-Kamiokande. Possible implications are analyzed.



The SATURNE Collaboration



RUSSIAN FEDERAL NUCLEAR CENTER
ALL-RUSSIAN RESEARCH INSTITUTE OF EXPERIMENTAL PHYSICS



PRODUCTION
ASSOCIATION
MAYAK




LOMONOSOV MOSCOW
STATE UNIVERSITY



JOINT INSTITUTE
FOR NUCLEAR RESEARCH



IPM RAS



Ioffe
Physical-
Technical
Institute

The search for coherent elastic neutrino-atom scattering and neutrino magnetic moment in Sarov

Sarov Tritium Neutrino Experiment = SATURNE Collaboration





Matteo Cadeddu (INFN, Cagliari), Francesca Derdei (INFN, Cagliari), Carlo Giunti (INFN, Turin),
Konstantin Kouzakov (MSU), Bayarto Lubsandorzhev (INR RAS), Oleg Moskalev, (VNIIEF, Sarov),
Ivan Stepantsov (MSU), Alexander Studenikin (MSU), Vladimir Trofimov (JINR),
Maxim Vyalkov (MSU), Arkady Yukhimchuk (VNIIEF, Sarov)



The **Sarov Tritium Neutrino Experiment (SATURNE)** is part of the research program of the **National Center for Physics and Mathematics** founded in 2021

Architectural and urban planning of NCPM in Sarov



The main goals of the experiment are

- first observation of coherent elastic neutrino-atom scattering ($CE\nu AS$)
- search for neutrino magnetic moment

using a high-intensity tritium neutrino source: at least 1 kg, possibly up to 4 kg of T_2

CE ν AS vs CE ν NS

CE ν NS: Coherent Elastic Neutrino-Nucleus Scattering

predicted by D. Z. Freedman, PRD 9 (1974) 1389;

V. B. Kopeliovich & L. L. Frankfurt, ZhETF Pis. Red. 19, No. 4 (1974) 236

observed by D. Akimov et al. (COHERENT Collab.), Science 357 (2017) 1123

CE ν NS

▶ $|\vec{q}|R_{\text{nuc}} \ll 1$


\vec{q} is the momentum transfer

R_{nuc} is the nuclear radius

CE ν AS

▶ $|\vec{q}|R_{\text{atom}} \ll 1$

R_{atom} is the atomic radius



CE ν NS: $E_{\nu} \lesssim 1/R_{\text{nuc}} \sim 200A^{-1/3}$ MeV

CE ν AS: $E_{\nu} \lesssim 1/R_{\text{atom}} \sim 1 - 10$ keV

Potentialities of a low-energy detector based on ^4He evaporation to observe atomic effects in coherent neutrino scattering and physics perspectives

M. Cadeddu,^{1,*} F. Dordei,^{2,†} C. Giunti,^{3,‡} K. A. Kouzakov,^{4,§} E. Picciau,^{1,||} and A. I. Studenik,^{5,6,¶}

¹Università degli studi di Cagliari and Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Cagliari, Complesso Universitario di Monserrato—S.P. per Sestu Km 0.700, 09042 Monserrato (Cagliari), Italy

²Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Cagliari, Complesso Universitario di Monserrato—S.P. per Sestu Km 0.700, 09042 Monserrato (Cagliari), Italy

³Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy

⁴Department of Nuclear Physics and Quantum Theory of Collisions, Faculty of Physics,

Lomonosov Moscow State University, Moscow 119991, Russia

⁵Department of Theoretical Physics, Faculty of Physics, Lomonosov Moscow State University,

Moscow 119991, Russia

⁶Joint Institute for Nuclear Research, Dubna 141980, Moscow Region, Russia

ⓧ (Received 14 July 2019; published 29 October 2019)

We propose an experimental setup to observe coherent elastic neutrino-atom scattering (CEvAS) using electron antineutrinos from tritium decay and a liquid helium target. In this scattering process with the whole atom, that has not been observed so far, the electrons tend to screen the weak charge of the nucleus as seen by the electron antineutrino probe. The interference between the nucleus and the electron cloud produces a sharp dip in the recoil spectrum at atomic recoil energies of about 9 meV, reducing sizably the number of expected events with respect to the coherent elastic neutrino-nucleus scattering case. We estimate that with a 60 g tritium source surrounded by 500 kg of liquid helium in a cylindrical tank, one could observe the existence of CEvAS processes at 3σ in 5 yr of data taking. Keeping the same amount of helium and the same data-taking period, we test the sensitivity to the Weinberg angle and a possible neutrino magnetic moment for three different scenarios: 60, 160, and 500 g of tritium. In the latter scenario, the Standard Model (SM) value of the Weinberg angle can be measured with a statistical uncertainty of $\sin^2\theta_W^{SM} - 0.015$. This would represent the lowest-energy measurement of $\sin^2\theta_W$, with the advantage of being not affected by the uncertainties on the neutron form factor of the nucleus as the current lowest-energy determination. Finally, we study the sensitivity of this apparatus to a possible electron neutrino magnetic moment and we find that using 60 g of tritium it is possible to set an upper limit of about $7 \times 10^{-13} \mu_B$ at 90% C.L., that is more than one order of magnitude smaller than the current experimental limit.


DOI: 10.1103/PhysRevD.100.073014

I. INTRODUCTION

Coherent elastic neutrino-nucleus scattering (CEvNS) has been recently observed by the COHERENT experiment [1,2], after many decades from its prediction [3–5].

This observation triggered a lot of attention from the scientific community and unlocked a new and powerful tool to study many and diverse physical phenomena: nuclear physics [6,7], neutrino properties [8–10], physics beyond the Standard Model (SM) [11–17], and electroweak interactions [18,19]. The experimental challenge related to the CEvNS observation is due to the fact that in order to meet the coherence requirement $qR \ll 1$ [20], where $q = |\vec{q}|$ is the three-momentum transfer and R is the nuclear radius, one has to detect very small nuclear recoil energies E_R , lower than a few keV.

At even lower momentum transfers, such that $qR_{\text{atom}} \ll 1$, where R_{atom} is the radius of the target atom including the electron shells, the reaction can be viewed as taking place on the atom as a whole [21]. This effect should be visible for $qR_{\text{atom}} \sim 1$, i.e., for momentum



In our paper we have proposed an experimental setup to observe coherent elastic neutrino-atom scattering (CEvAS) using electron antineutrinos from tritium decay and a supefluid ^4He target.

In this scattering process with the whole atom, that has not been observed so far, the electrons tend to screen the weak charge of the nucleus as seen by the electron antineutrino probe.

$$\mu_\nu \sim 10^{-13} \mu_B$$

supefluid ^4He target technology (HeRALD) for direct detection of sub-GeV DM has been recently proposed in: S.Hartel et al., Phys.Rev.D 100 (2019) 9, 092007

*matteo.cadeddu@ca.infn.it
 †francesca.dordei@cern.ch
 ‡carlo.giunti@to.infn.it
 §kouzakov@gmail.com
 ||emmanuele.picciau@ca.infn.it
 ¶studenik@srdsinp.msu.ru


Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

He-4 atomic-recoil spectrum with tritium $\bar{\nu}_e$


M. Cadeddu, F. Dordei, C. Giunti, K. Kouzakov, E. Picciau, A. Studenikin, *PRD* **100** (2019) 073014

$$\frac{d\sigma_{\text{SM}}}{dT_R} = \frac{G_F^2 m}{\pi} \left[Z\left(\frac{1}{2} - 2\sin^2\theta_W\right) - \frac{1}{2}N + Z\left(\frac{1}{2} + 2\sin^2\theta_W\right)F_{\text{el}}(q^2) \right]^2 \left(1 - \frac{mT_R}{2E_\nu^2} \right)$$

$$\frac{d\sigma_{\mu\nu}}{dT_R} = \frac{\pi\alpha^2 Z^2}{m_e^2} |\mu_\nu|^2 \left(\frac{1}{T_R} - \frac{1}{E_\nu} \right) [1 - F_{\text{el}}(q^2)]^2 \quad \text{with} \quad q^2 = 2mT_R$$



500 kg of helium
60 g of tritium
5 yrs of taking data



Summary and outlook

The Sarov tritium neutrino experiment aims at

- (i) first observation of **coherent elastic neutrino-atom scattering**
to test SM neutrino interactions at unprecedentedly low energies
- (ii) search for **neutrino magnetic moment**

A high-intensity tritium neutrino source is being prepared

- at least **1 kg, 10 MCi** (possibly up to **4 kg, 40 MCi**)

A 1000-L He II detector is being developed (to be **ready by 2027**)

- observation of **CEvAS (2032)**

- sensitivity to $\mu_\nu \sim (2-4) \times 10^{-13} \mu_B$ (**2032**)

A 4-kg Si detector is being developed (to be **ready by 2026**)

- sensitivity to $\mu_\nu \sim (1-1.5) \times 10^{-12} \mu_B$ (**2027**)

A 14-kg SrI₂(Eu) detector is being developed (to be **ready by 2025**)

- sensitivity to $\mu_\nu \sim (1.5-2) \times 10^{-12} \mu_B$ (**2026**)



LOMONOSOV MOSCOW
STATE UNIVERSITY

The SATURNE Collaboration



National Center
FOR PHYSICS AND MATHEMATICS

M. Cadeddu, F. Dordei, C. Giunti, K. Kouzakov, E. Picciau, A. Studenikin,

- *Potentialities of a low-energy detector based on 4He evaporation to observe atomic effects in coherent neutrino scattering and physics Perspectives*, *Phys.Rev.D* **100** (2019) 073014

- *New process in superfluid 4He detectors: The coherent elastic neutrino-atom scattering*, *PoS ICHEP2020* (2021) 211

A.Yukhimchuk, K.Kouzakov, A.Studenikin et al.,

- *Physics of hydrogen isotopes*, *PhysMath J.* **1** (2023) p. 5-19,
DOI: 10.56304/S2949609823010057

Presentations on behalf of SATURNE Coll.

- XXI Lomonosov Conference on Elementary Particle Physics, *August 2023, Moscow*
- First African Conference on High Energy Physics, *October 2023, Rabat, Marocco*
- XXII International Seminar on High-Energy Physics “Quarks 2024”,
May 2024, Pereslavl Zalesky, Russia
- LXXIV International Conference “Nucleus-2024”, *July 2024, Dubna*

Supported by the Russian Science Foundation project #24-12-00084



Национальный центр
Физики и Математики

National Centre for
Physics and Mathematics

XVI International School on Neutrino Physics and Astrophysics September 23-27, 2024, Sarov, Russia

<http://school.lomcon.ru>



Moscow State University
Branch and Techno Park Sarov

Unified Organizing Committee of Scientific Schools of NCPM

- D. Bisikalo (NCPM) – Co-chair
- S. Davydenko (NCPM) – Secretary
- M. Devyatkin (RFNC-VNIIEF)
- V. Goloviznin (MSU branch in Sarov)
- A. Vasiliev (NCPM)
- V. Il'gisonis (ROSATOM Corp.)
- A. Kovalishin (RFC “Kurchatov Institute”)
- V. Kveder (Russian Academy of Sciences)
- S. Nedelko (JINR)
- O. Olkhov (ROSATOM Corp.)
- A. Saveliev-Trofimov (MSU branch in Sarov)
- B. Sharkov (JINR & VNIIEF & NCPM)
- A. Sergeev (NCPM) – Chair
- B. A. Soloviev (Techno Park Sarov)
- V. Voevodin (MSU branch in Sarov)
- V. Soloviev (RFNC-VNIIEF)
- O. Vorontsova (RFNC-VNIIEF)

Working Group of Organizing Committee

- T. Aksentyeva (ROSATOM Corp.)
- F. Lazarev (MSU)
- A. Lichkunov (MSU)
- A. Popov (MSU)
- A. Purtova (MSU)
- K. Stankevich (MSU)
- T. Stavnichaya (Techno Park Sarov)
- M. Vyalkov (MSU & NCPM)

International Advisory Committee

- I. Bozovic Jelisavcic (Univ. of Belgrade)
- C. Giunti (INFN, Turin)
- D. Gorbunov (INR RAS)
- Y.-F. Li (IHEP CAS, Beijing)
- V. Neznamov (VNIIEF)
- M. Rebelo (CFTP & IST, Lisbon)
- V. Shevchenko (MEPhI)
- A. Studenikin (MSU & NCPM) – Co-chair
- I. Tkachev (INR RAS)
- A. Yukhimchuk (VNIIEF & NCPM) – Co-chair

Programme Committee

- E. Boos (MSU)
- D. Bisikalo (NCPM)
- S. Davydenko (NCPM)
- K. Kouzakov (MSU & NCPM) – Scientific Secretary
- A. Popov (MSU)
- A. Studenikin (MSU & NCPM) – Co-chair
- A. Yukhimchuk (VNIIEF & NCPM) – Co-chair

Organizing Committee at Faculty of Physics of MSU

- K. Kouzakov (MSU) – Scientific Secretary
- A. Popov (MSU)
- K. Stankevich (MSU)
- A. Studenikin (MSU) – Chair

for contacts:

[Alexander Studenikin lomcon@phys.msu.ru](mailto:Alexander.Studenikin@lomcon.ru)
[Arkady Yukhimchuk arkad@triton.vniief.ru](mailto:Arkady.Yukhimchuk@triton.vniief.ru)



РОСАТОМ



РОСАТОМ
РФЯЦ-ВНИИЭФ

<http://school.lomcon.ru>

online participation
by request
for connection to

studenik@srd.sinp.msu.ru

Konstantin Kouzakov (MSU)

on behalf of SATURNE Coll.

“Neutrino interactions with
atomic systems and the
SATURNE project”

September 27, 2024

A scenic view of a coastal town with a white church on a small island, surrounded by blue water and mountains. The church is a white building with a tiled roof and a bell tower. In the foreground, there is a dock with several small boats. The background shows a large body of water and mountains under a clear sky.

Thank you!

... about 200 speakers and 400 participants



Dedicated to the 270th Anniversary of
Moscow State University

TWENTY-SECOND LOMONOSOV CONFERENCE ON ELEMENTARY PARTICLE PHYSICS

Moscow, August 21 - 27, 2025



Tests of Standard Model & Beyond
Theories Beyond Standard Model
Neutrino Physics
Astroparticle Physics
Electroweak Theory
Gravitation and Cosmology
Developments in QCD
Heavy Quark Physics
Physics at Future Accelerators

XVII International
School on Neutrino
Physics and Astrophysics
August 22 - 25, 2025

Under the patronage of
V. Sadovnichy
Rector of MSU

Organizing Committee

International Advisory Committee

- V. Belokurov (MSU)
- E. Boos (MSU)
- I. Bozovic Jelisavcic (Univ. of Belgrade)
- S. Choudhury (IISc, Bangalore)
- M. Danilov (Lebedev Phys. Inst.)
- Z. Djuric (Argonne Nat. Lab.)
- A. Dolgov (Novosibirsk & Ferrara Univ.)
- A. Fedyanin (MSU)
- C. Giunti (INFN, Turin)
- M. Itkis (JINR)
- L. Kravchuk (INR, Moscow)
- M. Libanov (INR, Moscow)
- M. Malek (Univ. of Sheffield)
- V. Matveev (JINR)
- K.K. Phua (World Scientific Publ. Co., Singapore)
- K. Postnov (MSU)
- A. Sergeev (NCPM, Sarov)
- B. Shirkov (JINR, VNIIEF & NCPM, Sarov)
- V. Shevchenko (MEPhI)
- J. Silk (Univ. of Oxford)
- A. Skrinsky (INP, Novosibirsk)
- V. Soloviev (VNIIEF & NCPM, Sarov)
- P. Spillantini (INFN, Florence)
- G. Trubnikov (JINR)
- Z.-Z. Xing (IHEP, Beijing)
- L. Zivkovic (Inst. of Physics, Belgrade)
- V. Bednyakov (JINR)
- A. Bondar (INP, Novosibirsk)
- A. Egorov (ICAS)
- D. Galtsov (MSU)
- A. Isaev (MSU & JINR)
- A. Kataev (INR, Moscow)
- K. Kouzakov (MSU & NCPM)
- Yu. Kudenko (INR, Moscow)
- F. Lazarev (MSU)
- A. Lichkunov (MSU)
- O. Moskalev (VNIIEF & NCPM, Sarov)
- D. Naumov (JINR)
- A. Nikishov (Lebedev Phys. Inst.)
- A. Popov (MSU)
- Yu. Popov (MSU)
- A. Purtova (MSU)
- V. Ritus (Lebedev Phys. Inst.)
- G. Rubtsov (INR, Moscow)
- V. Savrin (MSU)
- V. Shakhov (MSU)
- K. Stankevich (MSU)
- A. Studenikin (MSU & NCPM, Sarov)
- A. Vasilev (Rosatom & NCPM, Sarov)
- M. Vyalkov (NCPM & MSU branch in Sarov)
- E. Yakushev (JINR)
- A. Yukhimchuk (VNIIEF & NCPM, Sarov)



Organizers and sponsors

- Faculty of Physics, MSU
- Joint Institute for Nuclear Research (Dubna)
- Institute for Nuclear Research (Moscow)
- Skobeltzyn Institute of Nuclear Physics, MSU
- Bruno Pontecorvo Laboratory on Neutrino Physics and Astrophysics (MSU)



INTERREGIONAL CENTRE FOR ADVANCED STUDIES

For contacts: Alexander Studenikin, Chairman
Scientific Secretaries: Konstantin Stankevich, Fedor Lazarev, Artem Popov and Anastasia Purtova
e-mail: lomcon@phys.msu.ru



... registration will be opened
in September 2024 ...
<http://lomcon.ru>

... Backup slides ...

✓ exhibits unexpected properties (puzzles)

W. Pauli, 1930

- neutral “neutron” \Rightarrow ✓ E. Fermi, 1933
- probably $m_\nu \neq 0$! ?

↪ Pauli himself wrote to Baade:

“Today I did something a physicist should never do. I predicted something which will never be observed experimentally...”

H. Bethe, R. Peierls,

«The 'neutrino'»

Nature 133 (1934) 532

- «There is no practically possible way of observing the neutrino»



... puzzles ...


- ... what about electromagnetic properties of ν ?

Coherent elastic ν - nucleous scattering (CE ν NS)



Predicted in 1974 (Freedman)

constrains
on
fundamental
physics



Observations: COHERENT (2017 – CsI detector,
2020 – Ar detector)
Dresden-II reactor
(2022 – Ge detector)



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: May 14, 2019
REVISED: June 21, 2019
ACCEPTED: July 9, 2019
PUBLISHED: July 17, 2019

Probing neutrino transition magnetic moments with coherent elastic neutrino-nucleus scattering

O.G. Miranda,^a D.K. Papoulias,^b M. Tórtola^b and J.W.F. Valle^b

^aDepartamento de Física, Centro de Investigación y de Estudios Avanzados del IPN, Apartado Postal 14-740 07000 Mexico, Distrito Federal, Mexico

^bAHEP Group, Institut de Física Corpuscular — CSIC/Universitat de València, Parc Científic de Paterna, C/Catedrático José Beltrán 2, E-46980 Paterna, Valencia, Spain

E-mail: omr@fis.cinvestav.mx, dipapou@ific.uv.es, mariam@ific.uv.es, valle@ific.uv.es

ABSTRACT: We explore the potential of current and next generation of coherent elastic neutrino-nucleus scattering (CE ν NS) experiments in probing neutrino electromagnetic interactions. On the basis of a thorough statistical analysis, we determine the sensitivities on each component of the Majorana neutrino transition magnetic moment (TMM), $|\Delta_i|$, that follow from low-energy neutrino-nucleus experiments. We derive the sensitivity to neutrino TMM from the first CE ν NS measurement by the COHERENT experiment, at the Spallation Neutron Source. We also present results for the next phases of COHERENT using HPGe, LAr and NaI[Tl] detectors and for reactor neutrino experiments such as CONUS, CONNIE, MINER, TEXONO and RED100. The role of the CP violating phases in each case is also briefly discussed. We conclude that future CE ν NS experiments with low-threshold capabilities can improve current TMM limits obtained from Borexino data.

JHEP07(2019)103

- Neutrino, electroweak, and nuclear physics from COHERENT ... with refined quenching factor, Cadeddu, Dordei, Giunti, Li, Zhang, PRD 2020

COHERENT data have been used for different purposes:

- nuclear neutron distributions
Cadeddu, Giunti, Li, Zhang
PRL 2018
- weak mixing angle
Cadeddu & Dordei, PRD 2019
Huang & Chen 2019
- ν electromagnetic properties
Papoulias & Kosmas PRD 2018
- ν non-standard interactions
Coloma, Gonzalez-Garcia,
Maltoni, Schwetz PRD 2017
Liao & Marfatia PLB 2017

Global map of CE ν NS experiments

Experiments

- Stopped-pion beams
 $E_\nu \sim$ few tens of MeV
- Nuclear reactors
 $E_\nu \sim$ few MeV



● Future/Planned

[C. Bonifazi, Neutrino 2022]

CE ν AS: Coherent Elastic Neutrino-Atom Scattering

Yu. V. Gaponov and V. N. Tikhonov,


Elastic scattering of low energy neutrinos by atomic systems,

Yad. Fiz. (USSR) 26 (1977) 594 (in Russian).

Abstract. Elastic scattering of low energy neutrinos by atomic systems is treated. For the V variant of weak interaction scattering on the total system (on electrons, protons and neutrons) is coherent; for the A variant neutrino scatters coherently on using simple atomic systems. The result for an arbitrary atom is presented. The analysis shows that at neutrino energies $\lesssim 10$ keV a region of coherent optical neutrino phenomena exists where the neutrino elastic scattering by an atom as a whole dominates.

So far there is no corresponding experimental observation. An experimental study of CE ν AS could provide a unique test of the SM neutrino interactions at very low energies.

Schematic representation of detector proposed to observe $CE\nu AS$ processes





- We consider a detector setup such that the tritium source is surrounded with a cylindrical superfluid-helium tank.
- This configuration allows us to maximize the geometrical acceptance, while allowing us to have a top flat surface where helium atoms could evaporate after a recoil.


${}^4\text{He}$ volume $\sim 1 \text{ m}^3$, tritium source activity: 1 kg (10 MCi) with the possibility to increase up to 4 kg (40 MCi)

- The recoil of a helium atom after the scattering with an electron antineutrino coming from the tritium source produces phonons and rotons which, upon arrival at the top surface, cause helium atoms to be released by quantum evaporation. An array of bolometers on the top surface detects the number of helium atoms evaporated.

Experimental scheme: tritium source, tank with liquid helium, dilution refrigerator, passive shielding, bolometer array




The tritium source design includes copper tubular elements



with titanium tritide TiT_2


- ✓ flux: $F_{\text{max}} \sim 1 \times 10^{14} \text{ cm}^2 \text{ s}^{-1}$ (1 kg)
- ✓ flux: $F_{\text{max}} \sim 2 \times 10^{14} \text{ cm}^2 \text{ s}^{-1}$ (4 kg)

He II detector concept to study CE ν AS



Tritium neutrino source
(1-4 kg, 10-40 MCi)

- Tubular copper elements with TiT₂



Helium II detector (1000 L)

- Liquid He-4 at 40-60 mK
- Array of 1000 TESs (transition edge sensors)
- 1000-channel SQUID readout

Expected results after 5 years of data collection

Number of CE ν AS events within SM: 60 for 1 kg of T₂
and 200 for 4 kg of T₂

Sensitivity to neutrino magnetic moment: $\mu_\nu \sim (2-4) \times 10^{-13} \mu_B$ at 90% C.L.

Expected number of $CE\nu AS$ events for 5 years data taking

$$N_1 = 58,9 \quad (10 \text{ MCi}), \quad N_2 = 195,2 \quad (40 \text{ MCi}),$$

... sensitivity to $CE\nu AS$ of at least 5σ

$$\text{test } \mu_\nu \sim 3.8 \times 10^{-13} \mu_B \quad (1 \text{ kg of tritium})$$

at 90% C.L.


$$\text{test } \mu_\nu \sim 2.3 \times 10^{-13} \mu_B \quad (4 \text{ kg of tritium})$$

Important determining factors that ensure success of project are

- many years of experience in working with radioactive tritium at VNIIEF Sarov
- already available infrastructure and personal
- confirmed availability of 1 kg of tritium at the first stage of project (up to 4 kg of tritium at the next stage)



Low-background neutrino laboratory in Sarov @ All-Russian Research Institute of Experimental Physics, RFNC



The overburden of ~18 m.w.e. stops the soft and hadronic components of cosmic radiation